



Wind Speed Response of Sap Flow in Five Subtropical Trees Based on Wind Tunnel Experiments

Sophie Laplace^{1*}, Tomonori Kume¹, Chia-Ren Chu²
and Hikaru Komatsu³

¹Graduate School of Forestry and Resource Conservation, National Taiwan University, Taipei, Taiwan.

²Department of Civil Engineering, National Central University, Zhongli, Taiwan.

³The Hakubi Center for Advanced Research, Kyoto University, Kyoto, Japan.

Authors' contributions

This work was carried out in collaboration between all authors. Author SL performed the experiments and statistical analyses and drafted the manuscript. Authors TK and CRC designed the study and wrote the protocol. Authors SL and HK performed the literature searches. All authors read and approved the final manuscript.

Research Article

Received 18th March 2013

Accepted 7th June 2013

Published 31st July 2013

ABSTRACT

Aims: We evaluated the responses of tree sap flow to wind speeds in coniferous and broad-leaved plants under steady and unsteady wind conditions.

Study Design: We performed sap flow and micro-meteorological measurements on two conifers, *Chamaecyparis obtusa* var. *formosana* and *Araucaria cunninghamii*, and three broadleaved species, *Swietenia mahagoni*, *Michelia formosana* and *Plumeria acutifolia* in a wind tunnel.

Place and Duration of Study: Civil Engineering Department, National Central University, Zhongli, Taiwan, China, between May and July 2011.

Methodology: In a wind tunnel, wind speed was increased incrementally from 0 to 2, 4, 6, and 8 m s⁻¹, then decreased from 8 to 0 m s⁻¹. To examine how sap flow responded to unsteady wind, we estimated the time constant of each individual for each step change in wind speed. Moreover, we examined differences in transpiration rate and leaf conductance among individuals under stable wind speeds.

Results: The time constant was generally about 30 min in both conifers and broadleaved

*Corresponding author: Email: felttie2000@yahoo.com.tw;

individuals. Interestingly, under steady winds, transpiration rate showed two different response patterns to increased wind speed: linear and saturated. The two patterns may be a consequence of different stomatal conductance values, but not of different leaf shapes.

Conclusion: Our results suggested that neither hydraulic system nor leaf shape differences between coniferous and broad-leaved trees was an ultimate factor affecting the transient response of tree sap flux and transpiration to wind speed, and that stomatal conductance played an important role in transpiration regulation in response to wind speed.

Keywords: *Transient response; transpiration rate; stomatal conductance; conifers; broadleaved tree species.*

ABBREVIATIONS

a, n	regression constants (Granier, 1985); $a = 1.123, n = 1.19 \times 10^2$
ΔT_m	maximum temperature difference between heated and non-heated probes at $V = 0$ ($^{\circ}\text{C}$)
ΔT	temperature difference between heated and non heated probes at a given V ($^{\circ}\text{C}$)
E	rate of leaf transpiration ($\text{g m}^{-2} \text{s}^{-1}$)
g_a	boundary layer conductance (m s^{-1})
g_s	stomatal conductance (m s^{-1})
g_w	leaf conductance for vapor transport ($\text{mol m}^{-2} \text{s}^{-1}$)
LA	total leaf area (m^2)
Q	whole water use amount ($\text{g m}^{-1} \text{s}^{-1}$); $Q = Fd \times As$, where $Fd = V \times 1.0 \times 10^6$ g m^{-3} and As is the stem sectional area at sensor height (m^2)
r_a	boundary layer resistance ($\text{m}^2 \text{s mol}^{-1}$)
r_w	leaf resistance for vapor transport ($\text{m}^2 \text{s mol}^{-1}$)
T_c	time constant (min)
U	wind speed (m s^{-1})
V	sap flow ($\mu\text{m s}^{-1}$)
w_i, w_a	mole or volume fractions of water vapor in the intercellular spaces and air, respectively (g mol^{-1}) (w_i is the saturated water vapor at a given leaf temperature)

1. INTRODUCTION

In recent years, climate change has considerably impacted the hydrology of terrestrial ecosystems, and hence water resources [1,2]. Plant transpiration is an important hydrological process that regulates water, energy, and carbon dioxide fluxes in terrestrial ecosystems [3]. Therefore, understanding transpiration is essential to predict potential ecosystem responses to climate change. Transpiration largely depends on five environmental factors: wind, radiation, temperature, humidity, and soil moisture. Wind has especially substantial effects on transpiration [4].

To understand how wind affects transpiration, both steady and unsteady wind-speed conditions must be considered, but traditionally, only steady wind conditions have been tested in controlled-environment wind tunnels [5,6,7]. Dixon and Grace [8] clarified that wind

could significantly affect transpiration, mainly by changing the boundary layer resistance (r_a) between leaves and air. The boundary layer is strongly regulated by leaf shape [6,9]; the relatively rough leaf surfaces of coniferous needle leaves can have smaller r_a values than the relatively large, smooth leaves of broad-leaf trees. Thus, coniferous and broad-leaf trees might show systematic differences in transpiration responses to wind speed. However, few wind-tunnel studies have compared how wind speed impacts transpiration in coniferous and broad-leaf trees.

Wind typically varies substantially over time. Variable wind speeds cause unsteady sap flow because of internal water-dynamics responses in plants. Sap flow (i.e., water uptake) has been reported to lag behind leaf transpiration [10,11]. The time lag is regulated by hydraulic resistance and capacitance in plants and is often characterized using a time constant parameter (T_c) defined as the time taken for sap flow to reach 63.2% of its steady velocity after a step change in transpiration [11]. Coniferous and broad-leaf trees use different xylem cells to transport water (tracheids and vessel elements, respectively) which might affect T_c . However, few studies have examined T_c in a variety of species because of difficulty in measuring it *in situ*. Previously, T_c of conifers was derived from a combination of field measurements and models [12,13]. In contrast, Chu et al. (2009) [14] successfully used a wind tunnel and sap flux measurements to determine T_c in a broad-leaf individual; however, wind tunnel experiments are usually limited to potted individuals of small size. Previous studies performed in wind tunnels estimated transpiration using weighing methods [6,7,8] that are unsuitable for unsteady wind conditions.

The present study aimed to clarify the response of tree sap flow and, hence, transpiration to wind speed in coniferous and broad-leaved plants in a wind tunnel. We measured the T_c of sap flow in five subtropical species, including broad-leaf and coniferous species, using step changes in wind speed. Moreover, differences in transpiration rate and leaf conductance between species were also evaluated under stable wind speeds.

2. MATERIALS AND METHODS

2.1 Wind Tunnel

Wind speed experiments were carried out in the open-circuit suction-type wind tunnel of the Department of Civil Engineering, National Central University, Zhongli, Taiwan (Fig. 1a). The total tunnel length was 30 m, and the test section was 18.5 m long, 3.0 m wide, and 2.1 m high. Constant wind speeds of 1–20 m s⁻¹ could be set by adjusting the rotation speed of a fan of 1.8 m diameter, with steady wind speed in the test section reached within 1–2 min. Wind speeds of 0, 2, 4, 6, and 8 m s⁻¹ were used in these experiments.



Fig. 1. Experimental set-up. a) External view of wind tunnel; b) *Chamaecyparis obtusa* var. *formosa*; c) *Araucaria cunninghamii*; d) *Swietenia mahagoni*; e) *Michelia formosana*; and f) *Plumeria acutifolia*

2.2 Trees Species

Five subtropical forest tree species from Taiwan were studied (Fig. 1b–f), including two conifers, *Chamaecyparis obtusa* var. *formosana* (Cupressaceae) and *Araucaria cunninghamii* (Araucariaceae), and three broad-leaved species, *Swietenia mahagoni* (Meliaceae), *Michelia formosana* (Magnoliaceae), and *Plumeria acutifolia* (Apocynaceae). Three specimens of each species were used. Trunk diameter at a height of 8 cm ranged from 2.0–3.0 cm in *C. obtusa* var. *formosana*, *A. cunninghamii*, and *P. acutifolia*; in *S. mahagoni* and *M. formosana*, trunk diameter at a height of 3 cm ranged from 1.0–1.5 cm (Table 1). Individuals of all species ranged in height from 80–130 cm, except *A. cunninghamii*, which were 160–180 cm tall.

Table 1. Tree characteristics

Species	Leaf shape	Tree height (cm)	Sensor height (cm)	Diameter* (cm)	LA (m ²)	g _s ** (m s ⁻¹)
<i>C. obtusa</i>	needle	1 128	18	1.74	0.73	6.11
		2 138		1.99	0.81	
		3 132		17.8	0.82	
<i>A. cunninghamii</i>	needle	1 183	18	3.28	0.76	19.62
		2 178		2.61	0.71	
		3 161		1.80	0.53	
<i>S. mahagoni</i>	broad leaf	1 102	13	1.48	0.14	35.79
		2 95		1.31	0.15	
		3 112		1.17	0.1	
<i>M. formosana</i>	broad leaf	1 90	13	1.15	0.03	822.45
		2 86		1.15	0.01	
		3 88		1.13	0.05	
<i>P. acutifolia</i>	broad leaf	1 120	18	3.29	0.49	12.09
		2 114		3.20	0.49	
		3 113		2.71	0.48	

*: Diameter at the height that sap flow sensors were installed.

** : g_s was derived from g_w under the condition of U = 6 m s⁻¹ (See Section 2.7)

2.3 Sap Flow Measurements: the Granier Sensors

To measure sap flow, we used a dual-probe sap flow measuring system [15,17], commonly referred to as the thermal dissipation probe (TDP) method. Sensors consisted of two cylindrical probes 10 mm long and 2 mm diameter inserted radially into the stem 10 cm apart. The upper probe included a heating element wound around a steel needle enclosing a T-type thermocouple (copper-constantan); it was continuously heated at constant power (0.15 W). The lower probe was unheated to measure the temperature of the wood tissue. Both thermocouples were connected to a data logger (CR1000, Campbell Scientific, Logan, UT, USA), and the temperature difference at the ends of the copper probe wires was recorded. Sap flux density (V) was calculated as:

$$V = 119 \times \left[\frac{\Delta T_m - \Delta T}{\Delta T} \right]^{1.231} \quad (1)$$

where: V is the sap flux density (g m⁻² s⁻¹); ΔT_m is the maximum temperature difference between heated and unheated probes at $V = 0$ (K); ΔT is the temperature difference between the heated and unheated probes at a given V (K), respectively. For each experiment, we estimated the maximum temperature difference using measurements taking during the night (18:00 to 6:00) preceding experiments.

The sensors were installed 18 cm high into stems of *C. obtusa* var. *formosana*, *A. cunninghamii*, and *P. acutifolia* and 13 cm high into *S. mahagoni* and *M. formosana*. The sensors were covered with aluminium to insulate them from radiation and potential cooling effects of the wind.

2.4 Meteorological Measurements

A hygrometric transmitter (HD9008TR, Delta Ohm, Padua, Italy) was placed near samples to measure air temperature and relative humidity. A soil moisture sensor (EC10, Decagon Devices, Pullman, WA, USA) was inserted in the soil of one sample per species. A quantum sensor (LI-190SZ, LI-COR Scientific, Lincoln, NE, USA) was installed on the ground upwind of the samples to measure photosynthetically active radiation. These meteorological data were recorded during all experiments using a data logger (CR10X, Campbell Scientific). Copper-constantan thermocouples (Type copper-constantan, Omega Engineering, Stamford, CT, USA) were used to measure the temperature difference between tree leaf surfaces and the air. One thermocouple tip was in constant contact with a leaf blade, avoiding the midrib; the other was in contact with the air. Leaf temperature was then calculated using the temperature difference and the measured air temperature inside the wind tunnel. Two to three pairs of thermocouples were installed on each tree in all experiments.

2.5 Procedure of Wind Speed Experiment

Three individuals of the one species were positioned in the test section of the wind tunnel 30 cm from one other in a row perpendicular to the wind direction. Five halogen lamps above them operated at the same intensity during all experiments. The potting soil was saturated before each experiment and covered with plastic film to prevent surface evaporation. The wind speed was increased stepwise from 0 m s⁻¹ to 2.0, 4.0, 6.0, and 8.0 m s⁻¹, then decreased from 8.0 to 0 m s⁻¹. Each wind speed was maintained for 1–3 h and changed only after sap flow had stabilized. The experiment was repeated for each tree species.

2.6 Time Constant (T_c)

To explain the unsteady water flow through plants, resistance-capacitance models derived from the SPAC (Soil Plant Atmosphere Continuum) concept have been used to determine how water transport lags behind transpiration [11]. In the SPAC model, T_c can be determined by the following simplified equation [13,14]:

$$V(t) = V_s \left[1 + \left(\frac{1}{\beta} - 1 \right) \exp \left(-\frac{t}{T_c} \right) \right] \quad (2)$$

where: $V(t)$ is the sap flux density (g m⁻² s⁻¹); V_s is the steady sap flux density after the change ($t = \infty$) (g m⁻² s⁻¹); T_c is the time constant (s); and $\beta = (R_c + R_s)/R_c$ where R_s and R_c are axial hydraulic resistance and tissue resistance associated with storage and withdrawal of water in xylem tissue, respectively.

Based on the least-square method, T_c and β were determined from the measured V_s and $V(t)$ at each wind speed treatment.

Because broad-leaf vessels have better water conductivity than conifer tracheids [16,19], the time response of sap flux is expected to be faster in broad-leaf plants. Thus, we hypothesized that T_c would be smaller in the broad-leaf trees than in the conifers.

2.7 Transpiration Rate (E) and Leaf Conductance (g_w)

The transpiration rate E ($\text{g m}^{-2} \text{s}^{-1}$) at the leaf area basis was calculated with the following equation:

$$E = \frac{Q}{LA} \quad (3)$$

where: Q is the sap flux (total water use: g s^{-1}) and LA is the total leaf area (m^2). Total water use $Q = V \times As$, where As is the sectional area of the stem at the height of sensors (m^2). After the experiments, we scanned each leaf with an area meter (Li-3000 scanning head with LI-3050C Transparent Belt Conveyor Accessory, LI-COR Scientific) and obtained the accumulated LA for all individuals. In this study, we assumed that a 1-cm-long sensor could cover the radial variation in V across the cross-section because of the small stem radius (< 1.6 cm) in our sample trees.

Flick's law describes the relationship between transpiration and leaf conductance as follows [17]:

$$E = g_w(w_i - w_a) = (w_i - w_a)/r_w \quad (4)$$

where: E is the rate of leaf transpiration ($\text{g m}^{-2} \text{s}^{-1}$); g_w is the leaf conductance for vapor transport ($\text{mol m}^{-2} \text{s}^{-1}$); w_i and w_a are the mole or volume fractions of water vapor in the intercellular spaces and in air, respectively (g mol^{-1}) (w_i is the saturated water vapor at the given leaf temperature); and r_w is the leaf resistance for vapor transport ($\text{m}^2 \text{s mol}^{-1}$).

The total leaf resistance for water vapour transfer, r_w , has two components: stomatal resistance r_s and boundary layer resistance r_a . The inverse of stomatal resistance is stomatal conductance, which is influenced by stomatal density, size, and aperture [18]. Boundary layer conductance g_a varies with leaf size and shape and also wind speed [9,20].

Based on Eq.(4), we calculated g_w using each V_s and meteorological data (humidity and air and leaf temperatures) at the time of V_s . Leaf conductance can be described as $1/g_w = 1/g_s + 1/g_a$, and because g_a increases with increasing wind speed, we can assume that g_w is equivalent to g_s at higher wind speeds ($U \geq 6 \text{ m s}^{-1}$).

Because conifer and broad-leaf leaves differ morphologically in both shape and texture (i.e., needle leaves with rough surface vs. broad leaves with smooth surfaces) [21] and because leaf size affects transpiration rate, we hypothesized that the transpiration rate of broad-leaf plants would be more responsive to wind speed.

3. RESULTS AND DISCUSSION

3.1 Changes in Sap Flow (V)

Between wind speeds of $0-8 \text{ m s}^{-1}$, the standard deviations of temperature and relative humidity were about 1.8°C and 3.2% , respectively. Sap flow V increased with wind speed (Fig. 2). At lower wind speeds ($U = 2 \text{ m s}^{-1}$), more time (about 125 min) was needed to reach steady flow than at higher wind speeds ($U = 6-8 \text{ m s}^{-1}$; about 40 min). A similar increase in V with wind speed was also found by Chu et al. [14]. Other studies showed a decline in transpiration with increased wind speed [6,8]. The Penman-Monteith equation [19] predicted

that transpiration would decrease at higher wind speeds occur under high available-energy conditions of $> 400 \text{ W m}^{-2}$, and that transpiration would increase with wind speed under low available-energy conditions [8]. Thus, our experiments and those of Chu et al. [14] may have been conducted under lower available-energy conditions (short wave radiation of $\sim 70 \text{ W m}^{-2}$; note that available energy is normally less than short wave radiation (12)) than in other studies. Here, higher wind speeds increased boundary layer conductance (g_a) possibly increasing transpiration. On the other hand, increased g_a also increases heat transfer (sensible heat), which decreased the leaf-air temperature differences (see eq.(4)), resulting in less transpiration under high available-energy conditions [8].




Fig. 2. *Michelia formosana* sap flow V ($\text{g m}^{-2} \text{ s}^{-1}$) at different wind speeds (U) between 0 and 8 m s^{-1} . Different symbols indicate V of different individuals

3.2 Time Constant (T_c)

The T_c of *C. obtusa* var *formosa* ranged from 8.0 min (when U increased from $6\text{--}8 \text{ m s}^{-1}$) to 77.9 min ($0\text{--}2 \text{ m s}^{-1}$), with a mean of 32.8 min. For *A. cunninghamii*, T_c ranged between 16.5 min ($6\text{--}8 \text{ m s}^{-1}$) and 29.4 min ($0\text{--}2 \text{ m s}^{-1}$), with a mean of 25.1 min. For *S. mahagoni*, T_c averaged 33.4 min and ranged between 9.7 min ($4\text{--}6 \text{ m s}^{-1}$) and 91.7 min ($0\text{--}2 \text{ m s}^{-1}$). For *M. formosana*, T_c ranged between 7.3 min ($6\text{--}8 \text{ m s}^{-1}$) and 16.6 min ($0\text{--}2 \text{ m s}^{-1}$) with a mean of 10.5 min. For *P. acutifolia*, T_c averaged 35.0 min and ranged from 12.8 min ($4\text{--}6 \text{ m s}^{-1}$) to 46.1 min ($0\text{--}2 \text{ m s}^{-1}$).

Fig. 3 shows that T_c of *C. obtusa* var. *formosana*, *M. formosana*, and *S. mahagoni* decreased with increasing wind speed. This means that when the wind speed was higher with a larger steady V , sap flow stabilized faster. In contrast, *M. formosana* and *A. cunninghamii* showed relatively conservative T_c values in response to changes in wind speed. The decreased T_c was likely to be found in the individuals with saturated trends of E

in response to the wind speed (Fig. 4). Here, T_c was defined as the multiple of total hydraulic resistance and capacitance in the plant hydraulic pathway [13,14]. In some previous studies, T_c values were implicitly assumed to be constant given that there was no severe soil water deficit [12,13]. In contrast, our study suggested that T_c in some species could change in response to wind speed and/or steady V , even when soil moisture was saturated, although the derived results were possibly due to an experimental artefact.

In our study, coniferous tree samples had T_c values of 16.5–32.8 min and broad-leaved tree samples had T_c values of 10.5–35.0 min. Although we expected that broad-leaved trees with vessels might have shorter T_c than conifers with tracheids based on previous studies [12,16], a difference was not apparent in this study. Also, at the lower wind speed of 2 m s^{-1} , the T_c value of the coniferous and broad-leaved trees were 29.4–77.9 and 16.6–91.7, respectively. At the higher wind speed of 8 m s^{-1} , the T_c values were 7.2–9.3 and 9.4–11.7, respectively.




Fig. 3. Time constant (T_c) of each tree species for each wind speed change

3.3 Transpiration Rate (E), Wind Speed (U), and Leaf Conductance (g_w)

Fig. 4 shows the increasing in normalized E in response to U . In our samples, we found two different trends: linear curves in *S. mahagoni*, *M. formosana*, and *A. cunninghamii*, and saturated curves in *C. obtusa* var. *formosana*, and *P. acutifolia*. Because g_a is strongly affected by leaf shape [6,9,21], we expected the correlations between E and U to differ between coniferous and broad-leaved trees. However, the linear pattern was seen in one coniferous and two broad-leaved tree species, and the saturated pattern occurred in one coniferous and one broad-leaved tree sample.

Previously, Jones [11] calculated the relationships between E , g_a , and g_s using the Penman–Monteith equation and showed that E could increase with g_a under larger- g_s condition but not under smaller- g_s condition. When g_s is large, E is regulated by g_a and hence U , while, E is limited by g_s regardless of g_a when g_s is small. Therefore, a linear increase in response to U would occur in individuals with relatively large g_s and a saturated response to U would be expected in individuals with relatively small g_s . In our study, estimated g_s for *S. mahagoni*, *M. formosana*, and *A. cunninghamii* (linear trends) was larger than those of *C. obtusa* var. *formosana* and *P. acutifolia* (saturated trends), in concordance with the hypothesis. Over all, this study clarified that g_s could be an important factor determining the effects of U on E in addition to leaf shape.




Fig. 4. Normalized transpiration rate (E) of five tree species based on changes in wind speed (U). Normalized E was calculated using maximum E in each individual during the whole experiment period

4. CONCLUSION

To clarify how steady and unsteady wind speeds affect sap flow, we investigated the T_c of sap flow under unsteady U and the relationship between E derived from steady sap flow, U , and g_s in two conifers and three broad-leaved species using a wind tunnel.

We expected to observe different patterns of water dynamics between conifers (with tracheids) and broad-leaved trees (with vessels). However, our wind tunnel experiments showed that the different xylem physiologies yielded approximately the same T_c values (about 10–30 min). Our samples also demonstrated two trends of E in response to increases

in U , but they were independent of leaf shape (i.e., needles vs. broad leaves). *Swietenia mahagoni*, *M. formosana*, and *A. cunninghamii*, with higher estimated g_s values, showed linear correlations between E and U and probably g_a . In contrast, the E values of *C. obtusa* var. *formosa* and *P. acutifolia*, with smaller g_s values, showed saturated curves, probably because E was limited by g_s .

We concluded that in addition to leaf shape, g_s could be a dominant factor controlling how wind speed affects E . However, we could not readily distinguish g_a and g_s in this experiment. Further experiments using plaster-coated branches (Landsberg and Ludlow, 1970) should enable us to separate g_a and g_s more precisely and to understand the mechanism by which wind speed impacts g_a , g_s , and E .

ACKNOWLEDGEMENTS

We thank Mr. Tsung-Chiang Liu, Mr. Yu-Feng Chang (National Central University), Ms. Mei-Li Chen, Ms. I-Fang Hsieh, and Mr. Yan-Fu Lin (National Taiwan University) for their help with the wind tunnel experiments. Samples were provided by the Experimental Forest of National Taiwan University. Our experiments were supported by the National Science Council of the Republic of China (100-2313-B-002-033-MY3).

COMPETING INTERESTS

Authors declare that no competing interests exist.

REFERENCES

1. Dibike YB, Coulibaly P. Hydrologic impact of climate change in the Saguenay watershed: comparison of downscaling methods and hydrologic models. *J Hydrol.* 2005;307(1):145-163.
2. Ferguson IM, Maxwell RM. Human impacts on terrestrial hydrology: climate change versus pumping and irrigation. *Environ Res Letters.* 2012;7(4):044022.
3. Thornthwaite CW and Mather JR. The Role of Evapotranspiration in Climate. *Theoretical Applied Climatol.* 1951;3(1):16-39.
4. Wooley JT. Mechanisms by which wind influences transpiration. *Plant Physiol.* 1961;36(1):112-114.
5. Bidwell A. Observations on the biology of nymphs of *Povilla adusta* NAVAS (Ephemeroptera: Polymitarchidae) in lake Kainji, Nigeria. *Hydrobiologia.* 1979;67:161-172.
6. Drake BG, Raschke K, Salisbury FB. Temperature and transpiration resistances of xanthium leaves as affected by air temperature, humidity, and wind speed. *Plant Physiol.* 1970;46(2):324-330.
7. Grace J. The effect of wind on grasses. I. Cuticular and stomatal transpiration. *Journal of Experimental Bot.* 1974;25:542-551.
8. Dixon M, Grace J. Effects of wind on the transpiration of young trees. *Ann Bot.* 1984;53:811-819.
9. Schuepp PH. Leaf boundary layers: Tansley Review No. 59. *New Phytol.* 1993;125:477-507.
10. Landsberg JJ, Ludlow MM. A technique for determining resistance to mass transfer through the boundary layers of plants with complex structure. *J Appl Ecol.* 1976;7:187-192.

11. Phillips NG, Scholz FG, Bucci SJ, Goldstein G, Meinzer FC. Using branch and basal trunk sap flow measurements to estimate whole-plant water capacitance: comment on Burgess and Dawson (2008). *Plant Soil*. 2009;315:315-324.
12. Jones HG. *Plants and microclimate*. 2nd ed. Cambridge University Press; 1992.
13. Phillips NG, Nagchaudhuri A, Oren R, Katul G. Time constant for water transport in loblolly pine trees estimated from time series of evaporative demand and stem sapflow. *Trees-Structure and Function*. 1997;11:412-419.
14. Phillips NG, Oren R, Licata J, Linder S. Time series diagnosis of tree hydraulic characteristics. *Tree Physiol*. 2004;24:879-890.
15. Chu CR, Hsieh CI, Wu SY, Phillips NG. Transient response of sap flow to wind speed. *J Exp Bot*. 2009;60(1):249-255.
16. Granier A. Une nouvelle méthode pour la mesure du flux de sève brute dans le tronc des arbres. *Ann Sci for*. 1985;42(2):81-88. French.
17. James SA, Clearwater MJ, Meinzer FC, Goldstein G. Variable length heat dissipation sensors for the measurements of sap flow in trees with deep sapwood. *Tree Physiol*. 2002; 22:277–283.
18. Perämäki M. A physical analysis of sap flow dynamics in trees. Academic dissertation. The Finnish Society of Forest Science; 2005.
19. Tyree M, Ewers FW. The hydraulic architecture of trees and other woody plants. *New Phytol*. 1991;199:345-360.
20. Lambers H, Stuart Chapin III F, Pons TL. *Plant physiological ecology*. 2nd ed. Springer Science; 2008.
21. Martin TA, Hinckley TM, Meinzer FC, Sprugel DG. Boundary layer conductance, leaf temperature and transpiration of *Abies amabilis* branches. *Tree Physiol*. 1999;19:435-443.
22. Monteith JL, Unsworth MH. *Principles of environmental physics*. 3rd ed. New York, 291: Edward Arnold; 1990.
23. Nobel PS. *Physiochemical and environmental plant physiology*. 4th ed. San Diego, 635: Academic Press; 1991.
24. Bond BJ, Meinzer FC, Brooks JR. How trees influence the hydrological cycle in forest ecosystems. In: Wood PJ, Hannah DM, Sadler JP, editors. *Ecohydrology: Past, Present and Future*. Chichester: John Wiley & Sons; 2007.

© 2013 Laplace et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the origin al work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=251&id=10&aid=1784>