



Antibacterial Activities of Aqueous and Hydroethanolic Extracts of *Anogeissus leiocarpus* on the *In vitro* Growth of two Multiresistant Strains of *Salmonella typhimurium* Isolated from Broilers Chickens

**Anzoumana Lassinan Ouattara ^{a*}, Ouattara Abou ^a,
Ouattara Karamoko ^{b,c}, Golly Koffi Julien ^d
and Coulibaly Adama ^b**

^a Department of Biochemistry-Microbiology (UFR Agroforestry), Jean Lorougnon Guede University of Daloa, P.O. Box 150, Daloa, Côte d'Ivoire.

^b Biochemical Pharmacodynamics Laboratory (UFR Biosciences), Félix Houphouët Boigny University of Cocody-Abidjan, P.O. Box 582, Abidjan, Côte d'Ivoire.

^c UFR Agriculture Halieutic Resources and Agro-Industry. (UFR-ARHAI), San Pedro University, P.O.Box V1800 San Pedro, Côte d'Ivoire.

^d Institut Pasteur of Côte d'Ivoire (IPCI), 01 P.O. Box 490 Abidjan 01, Côte-d'Ivoire.

Authors' contribution

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/MRJI/2023/v33i71394

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106063>

Original Research Article

Received: 14/07/2023

Accepted: 18/09/2023

Published: 23/09/2023

*Corresponding author: E-mail: yankon2003@yahoo.fr;

ABSTRACT

Poultry products have quickly become one of the most consumed sources of protein in the world. However, poultry farmers face enormous economic losses caused by the emergence of multi-resistant bacteria. In addition, the consumption of eggs and poultry meat also causes several infectious diseases within populations as well as poisonings and therefore a public health problem.

Aims: The objective of our study was to evaluate *in vitro* the antibacterial properties of aqueous and 70% hydroethanolic extracts of the leaves of *Anogeissus leiocarpus* on two strains of multiresistant *Salmonella typhimurium* isolated from the faeces of broilers chickens.

Methodology: The agar well method was used to test the susceptibility of bacterial strains while the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) were determined by the liquid dilution method.

Results: On the two multiresistant *Salmonella typhimurium* germs, the aqueous and 70% hydroethanolic extracts of *Anogeissus leiocarpus* gave zones of inhibition diameter between 13-22 mm while the minimum bactericidal concentrations (MBC) were ranged between 12.5 mg/ml and 25 mg/ml. The results of the phytochemical analysis of the extracts studied reveal the presence of several families of secondary metabolites including phenolic compounds, cardiac glycosides, saponins, alkaloids and then sterols and terpenes. However, these secondary metabolites are distributed differently within the aqueous and 70% hydroethanolic extracts.

Conclusion: In view of the results obtained in the present work, this plant could be used as a phytomedicine to combat pathologies linked to *Salmonella typhimurium* in broiler chickens and as an antibacterial agent in food products that may harbor strains of *Salmonella typhimurium*.

Keywords: *Anogeissus leiocarpus*; extracts; salmonella typhimurium; multiresistant.

1. INTRODUCTION

Poultry products have rapidly become one of the most consumed sources of protein in the world, making them very important for the global food supply [1]. It should represent about half of meat production in the next ten years. This growth will be mainly due to developing countries, which will contribute about 73% of the additional production [2]. Indeed, the increase in poultry production is particularly visible in developing countries, due to the status of poultry as cheap meat [3]. To this, it should be added that poultry farming, which was of the traditional or semi-industrial type, has rapidly evolved with the installation of numerous industrial farms following the establishment of public supervisory structures. However, the spectacular development of poultry farming is not without difficulties. Indeed, most breeders are not professionals and do not necessarily master the application of the basic rules of hygiene in poultry farms. This has favored the emergence of many diseases on these farms. These diseases decimate farms and cause considerable economic losses for breeders. In addition to the economic losses generated among poultry farmers, the consumption of eggs and poultry meat causes several infectious diseases within populations, thus creating poisonings and

therefore a public health problem. These pathologies are of all kinds: parasitic [4], viral [5], fungal [6] and especially bacterial [7]. Among bacterial diseases, salmonellosis is one of the most damaging diseases in poultry farming [7]. Therefore, antibiotic therapy based on an adequate diagnosis as well as prophylaxis, still remain the only means of combating this disease. However, these sanitary practices encounter many difficulties. In fact, in recent years, there has been an increasing appearance of antimicrobial resistance in salmonellosis [8]. Faced with these worrying therapeutic failures, the use of phytomedicines offers itself as a solution to be explored. *Anogeissus leiocarpus* is an important evergreen tree of the family Combretaceae, widely distributed in Africa [9,10] and is well known in African traditional medicine to treat skin sore, alopecia, headache, bronchitis, diarrhea, bruises, dysmenorrhea, constipation, dysentery, fever, cataract, anti-cold, inflammation, asthma, tumor, liver diseases and venereal diseases [11]. Extracts from this plant have shown antibacterial activities against several strains [12,13,14]. It is in this perspective that we evaluated the therapeutic effects of aqueous and hydroethanolic extracts of the leaves of *Anogeissus leiocarpus*, on two strains of multiresistant *Salmonella typhimurium* of avian origin.

2. MATERIALS AND METHODS

2.1 Plant Material

It consists of leaves of *Anogeissus leiocarpus*. They were collected in the village of Lataha in the Korhogo region (North of Côte d'Ivoire) in March 2022 and authenticated by the National Floristic Center of the Félix HOUPHOUËT-BOIGNY University of Cocody-Abidjan.

2.2 Bacterial Strains

Two bacterial strains of *Salmonella typhimurium* of avian origin were the subject of this study. They were provided by the Microbiology Unit of the Biotechnology Laboratory, of the UFR Biosciences of the Félix Houphouët-Boigny University of Côte d'Ivoire (Table 1).

2.3 Preparation of Aqueous and Hydroethanolic Extracts of *Anogeissus leiocarpus*

The leaves of *Anogeissus leiocarpus* were washed, cut and dried in the shade for two weeks. Once dried, these plant elements were ground. Thus, 100 g of this powder were diluted in 1 liter of distilled water. The mixture was homogenized at ambient laboratory temperature using a magnetic stirrer for 24 hours. The homogenate obtained was successively filtered twice on cotton wool and then once on Whatman paper (3 mm). The volume of the filtrate obtained is reduced using a Med Center Venticell type oven at 50°C to give a powder which constitutes the total aqueous extract [15]. The same operation was carried out using 70% ethanol instead of distilled water, to obtain the 70% hydroethanolic extract [16]. The two plant extracts obtained were stored in the refrigerator for the rest of the tests.

2.4 Preparation of the Bacterial Suspension

The bacterial inoculum was prepared by homogenizing two (2) colonies of each *Salmonella Typhimurium* Multi-R in 10 ml of Mueller-Hinton broth. After an incubation of approximately 3 hours at 37°C, 0.1 mL of the broth is added to 10 ml of sterile Mueller-Hinton broth to constitute the inoculum at 10⁶ bacteria/ml with a turbidity of 0.5 Mac Farland [15].

2.5 Antibacterial Susceptibility Screening of Aqueous and Hydroethanolic Extracts

This test is based on the determination of growth inhibition zones. The method used was that of punch wells on Mueller-Hinton agar. Thus, as in the classic production of an antibiogram, the agar is inoculated by flooding (inoculum at 10⁶ bacteria/ml). Then we make wells 6 mm in diameter. Then each well was filled with 80 µl of plant extract at a concentration of 200 mg/ml, taking care to separate two holes of at least 20 mm [17,18]. After a pre-diffusion of 45 minutes at room temperature under the hood, the whole was incubated in an oven at 37°C for 24 hours. After incubation, the action of each extract was assessed by measuring a growth inhibition zone (absence of colonies) around the well.

2.6 Determination of Minimum Inhibitory and Bactericidal Concentrations of Aqueous and Hydroethanolic Extracts

First, a range of concentrations of each plant extract ranging from 25 to 0.39 mg/ml was prepared by the double dilution method in eight (8) test tubes. Then, 1 ml of each concentration of plant extract prepared is mixed with 9 ml of the bacterial inoculum whose bacterial load is estimated at 10⁶ bacteria/ml. The Minimum Inhibitory Concentration (MIC) of the plant extract corresponds to the concentration of the first tube in which there is no growth visible to the naked eye of the bacteria tested after incubation for 24 hours at 37°C. To determine the Minimum Bactericidal Concentrations (MBC), the surface of a new Mueller-Hinton agar poured into a Petri dish was inoculated with 0.1 ml of the contents of the tubes having a concentration greater than or equal to the MIC using a calibrated loop. At the same time, double dilutions from the original suspension are carried out up to the 10⁻⁴ dilution. Then, these dilutions and the stock suspension are streaked onto Mueller-Hinton agar in another Petri dish. Seeding was done in parallel 5 cm streaks. All the plates containing the agars were incubated at 37°C for 24 hours. The MBC corresponds to the concentration of the extract presenting a number of colonies less than or equal to that of the 10⁻⁴ dilution. This MBC corresponds to the smallest concentration which allows at most 0.01% of the germs of the starting suspension to survive in 24 hours [15].

Table 1. Origin and antibiotic profiles of the *Salmonella typhimurium* strains studied

Salmonella strains tested	Origin	Antibiotic profiles
<i>Salmonella typhimurium</i> N°1	broilers chickens faeces	AMP-CHL-STR-SUL-TE-AUG-CTX-CIP-NA
<i>Salmonella typhimurium</i> N°2	broilers chickens faeces	AMP-CHL-STP-SUL-TE

AMP : ampicillin, CHL : chloramphénicol, STR : streptomycin, SUL : sulfamides, TE : tétracyclin, AUG : amoxicillin- clavulanic acid, CTX : cefotaxim, CIP : ciprofloxacin, NA : nalidixic acid

Table 2. Diameters of inhibition of extracts of *Anogeissus leiocarpus* on the strains of *Salmonella typhimurium* studied

Salmonella strains tested	Origin	Antibiotic profiles	Aqueous extract (200 mg/ml)	70% hydroethanolic extract (200 mg/ml)
<i>Salmonella typhimurium</i> N°1	broilers chickens faeces	AMP-CHL-STR-SUL-TE-AUG-CTX-CIP-NA	13±1 mm	16±0.5 mm
<i>Salmonella typhimurium</i> N°2	broilers chickens faeces	AMP-CHL-STP-SUL-TE	19±1 mm	22±0.5 mm

AMP: ampicillin, CHL : chloramphénicol, STR : streptomycin, SUL : sulfamides, TE : tétracyclin, AUG : amoxicillin- clavulanic acid, CTX : cefotaxim, CIP : ciprofloxacin, NA : nalidixic acid

2.7 Phytochemical Screening of to Aqueous and Hydroethanolic Extracts

The phytochemical analysis of the different extracts of *Anogeissus leiocarpus* have been based on the coloration and precipitation tests [19]. To better estimate the quantity of chemical constituents in the extracts, scores ranging from 0-3 (absent or present) have been allocated. Thus, the absence was symbolized by a score of 0, the presence in small quantities by a score of 1, the presence in average quantity score 2 and finally, the abundance by a score of 3.

3. RESULTS AND DISCUSSION

3.1 Antibacterial Susceptibility Screening of Extracts

Table 2 presents the values of the diameters of the growth inhibition zones of *Salmonella typhimurium* germs tested with aqueous and 70% hydroethanolic extracts of *Anogeissus leiocarpus*. It appears that each of the extracts prepared has a well-defined activity on the growth of the bacterial strains studied. The inhibition diameters are between 13 and 22 mm. Thus, the aqueous extract gave an inhibition

diameter of 13 mm on *Salmonella typhimurium* N°1 and 19 mm on the strain of *Salmonella typhimurium* N°2. As for the 70% hydroethanolic extract of *Anogeissus leiocarpus*, the inhibition diameters varied from 16 mm (*Salmonella typhimurium* N°1) to 22 mm (*Salmonella typhimurium* N°2). These first tests related to the determination of the zones of inhibition show the sensitivity of the bacteria studied to the plant extracts prepared because an extract is judged to be active if it induces a zone of inhibition greater than or equal to 10 mm [20].

Similar studies were conducted on aqueous and ethanolic extracts of *Tinospora cordifolia*, *Ziziphus mauritiana*, *Swertia chirata*, *Erythrina blakei* and *Asparagus densiflorus* to inhibit the growth of a strain of *Salmonella typhimurium* [21]. The results of the work of these authors showed that the diameters of the zones of inhibition of the aqueous extracts of these plants on the *in vitro* growth of the strain studied varied between 8.7 and 10.3 mm while those of the ethanolic extracts were between 5.2 and 7.7 mm. Also, a study showed that *Rhazya stricta* leaf extracts had varying effects on growth inhibition against eight (8) strains of *S. typhimurium* [22]. Thus, among the plant extracts studied by these authors, the hydroalcoholic extract was the most effective in inhibiting the growth of strains of *S.*

typhimurium with an average diameter of 12.25 mm. This extract was followed by the ethanolic extract with an inhibition diameter of 12.12 mm at different concentrations. Others authors showed that aqueous extracts at a concentration of 400 mg/ml from *Cymbopogon citratus*mm, *Psidium guajava* and *Anacardium occidentale* recorded inhibition diameters of 11, 14 and 17 mm respectively on a strain of *S. typhimurium* [23].

3.2 Antibacterial Parameters of Extracts

The values of the antibacterial parameters (MIC and MBC) as well as those of the MBC/MIC ratios determined are reported in Table 3. On the strain of *Salmonella typhimurium* N°1, the aqueous extract of *Anogeissus leiocarpus* recorded a minimum inhibitory concentration (MIC) which is equal to the minimum bactericidal concentration (MIC = MBC = 25 mg/ml). On this same strain of *Salmonella typhimurium* N°1, the 70% hydroethanolic extract of *A. leiocarpus* gave a minimum bactericidal concentration of 25 mg/ml which was twice the minimum inhibitory concentration (12.5 mg/ml).

In the presence of the strain of *Salmonella typhimurium* N°2, the MIC of the aqueous extract of *A. leiocarpus* was 12.5 mg/ml with also a MBC of 12.5 mg/ml. On the other hand, the 70% hydroethanolic extract of *A. leiocarpus* gave a minimum inhibitory concentration of 6.25 mg/ml and a MBC of 12.5 mg/ml on this germ of *Salmonella typhimurium* N°2. However, the MBC/MIC ratio of each extract prepared with respect to the two strains of *Salmonella typhimurium* studied was equal to 1 or 2. These ratios thus concluded that the plant extracts prepared were bactericidal

for the two *Salmonella typhimurium* germs which is the subject of study. Indeed, an extract is said to be bactericidal if the MBC/MIC ratio ≤ 2 [24].

Furthermore, we note from this study that the two aqueous and 70% hydroethanolic extracts gave MBCs of 25 mg/ml on *Salmonella typhimurium* N°1 against MBCs of 12.5 mg/ml on *Salmonella typhimurium* N°2 (Fig. 1). These results could be explained by the intrinsic resistance presented by each strain of *Salmonella typhimurium*. These results confirm those of [18] who proved that the antimicrobial activities are related to the origin of the extract and the strain tested as well as the nature of the solvent used.

Evaluation of the antimicrobial activities of the hydroethanolic extract of *Satureja sahendica* against 20 isolates of *S. typhimurium* showed that the values of MIC and MBC ranged from 0.29 to 4.68 mg/ml and 75-150 mg/ml respectively [25].

A study was carried out on an ethanolic extract of *Anogeissus leiocarpus* obtained by Soxhlet extraction on three β -lactamase-producing bacteria, namely *E. coli*, *K. pneumoniae*, *P. aeruginosa* and *S. typhi* [26]. Inhibition diameters ranging from 15.67 mm to 17.00 mm were obtained on bacteria tested during this study by the method of impregnated discs. This study also showed that the minimum inhibitory concentration was between 66.67 mg/l and 16.67 mg/l while the minimum bactericidal and bacteriostatic concentrations were between 266.67 mg/l and 106.67 mg/l and 266.67 mg/l to 80.00 mg/l respectively. In their study, the

Table 3. Antibacterial parameters of the extracts of *Anogeissus leiocarpus* on the strains of *Salmonella typhimurium* studied

Salmonella strains tested	Antibiotic profiles	Extracts of <i>Anogeissus leiocarpus</i>	Antibacterial Parameters		Ratios MBC/MIC	Effect of the extract
			MIC (mg/ml)	MBC (mg/ml)		
<i>Salmonella typhimurium</i> N°1	AMP-CHL-	Aqueous extract	25	25	1	bactericidal
	STR-SUL-TE-AUG-CTX-CIP-NA	70% hydroethanolic extract	12,5	25	2	bactericidal
<i>Salmonella typhimurium</i> N°2	AMP-CHL-	Aqueous extract	12,5	12,5	1	bactericidal
	STP-SUL-TE	70% hydroethanolic extract	6,25	12,5	2	bactericidal

AMP : ampicillin, CHL : chloramphénicol, STR : streptomycin, SUL : sulfamides, TE : tétracyclin, AUG : amoxicillin- clavulanic acid, CTX : cefotaxim, CIP : ciprofloxacine, NA : nalidixic acid, MIC : Minimum inhibitory concentration, MBC : Minimum bactericidal concentration




Fig. 1. Compared antibacterial effects of *Anogeissus leiocarpus* extracts.




Fig. 2. Phytochemical analysis of *Anogeissus leiocarpus* extracts

MBC/MIC ratio showed a bacteriostatic effect for *E. coli* and *S. typhi* and bactericidal for *K. pneumoniae* and *P. aeruginosa*. These results further show that the antibacterial activity of a plant extract depends on several factors. These factors may take into account the nature of the plant extract, the nature of the strains studied and the methodology used.

3.3 Phytochemical Analysis

The results of the phytochemical analysis of the extracts studied reveal the presence of several

families of secondary metabolites including phenolic compounds, cardiac glycosides, saponins, alkaloids and then sterols and terpenes. However, these secondary metabolites are distributed differently within the aqueous and 70% hydroethanolic extracts. Thus, in the present study, the aqueous extract of *Anogeissus leiocarpus* does not contain catechin tannins and sterols and terpenes. The 70% hydroethanolic extract also does not contain catechin tannins in this study. Also, the 70% hydroethanolic extract of *Anogeissus leiocarpus*

contains more phenolic compounds compared to the aqueous extract (Fig. 2).

The study by Dahiru *et al.* (2021) from an ethanolic extract of *Anogeissus leiocarpus* obtained by Soxhlet extraction made it possible to identify twenty-five (25) chemical compounds. Among these compounds, myristoleic acid; Z-7-tetradecenal; 4,5-dimethyl-4-Hexene-3-un; 1,15-hexadecadiene and linoleoyl chloride were the most abundant. In this study, the observed antibacterial activities are explained by the results of the phytochemical analysis of the studied extracts. Indeed, the antimicrobial activity of most of these compounds, in particular flavonoids, tannins, alkaloids as well as terpenes, has already been demonstrated by several researchers [27,28,29]. The action mechanisms of such active substances are in forms of inhibiting the cell wall synthesis, the cell membrane function and the protein synthesis [30].

4. CONCLUSION

This work allowed us to highlight the antibacterial properties of *Anogeissus leiocarpus* on two strains of multiresistant *Salmonella typhimurium* isolated from broilers chickens faeces. The aqueous and 70% hydroethanolic extracts prepared showed bactericidal powers on the two strains of multiresistant *Salmonella typhimurium* which were the subject of study. Thus, the aqueous and 70% hydroethanolic extracts of *Anogeissus leiocarpus* recorded minimal bactericidal concentrations of 12.5 mg/ml and 25 mg/ml on the two bacterial germs. In view of the results obtained in the present work, this plant could be used as a phytomedicine to combat certain pathologies linked to *Salmonella typhimurium* in broilers breeding. Also, it would be interesting to undertake studies with a view to using these extracts of *Anogeissus leiocarpus* as antibacterial agents in food products that may harbor strains of *Salmonella typhimurium*.

ACKNOWLEDGEMENTS

The authors acknowledge the Microbiology Unit of the Biotechnology Laboratory (UFR Biosciences of the Félix Houphouët-Boigny University of Côte d'Ivoire) for providing *Salmonella typhimurium* isolates.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Campos M. Poultry is King of Proteins in IPC Outlook: International Poultry Council; 2019.
Available: <https://internationalpoultrycouncil.org/2019/03/08/poultry-is-king-of-proteins-in-ipc-outlook/>.
2. Organization for Economic Cooperation and Development/Food and Agriculture Organization (OECD/FAO). Agricultural Outlook for OECD and FAO; 2016.
3. OECD/FAO. Commodity Snapshots: Meat. Oecd-FAO Agricultural Outlook for 2017-2026. Paris: OECD Publishing; 2017.
4. Bichet H, Sanaa M, Dorchie PH, Reperant JM. Sanitary and zootechnical impact of clinical coccidiosis upon laying hen in Senegal. Rev. of Veterinary Medicine. 2003;154(6):431-438.
5. White RL, Friedrich LV, Mihm LB, Bosso JA. Assessment of the relationship between antimicrobial usage and susceptibility: differences between the hospital and specific patient-care areas. Clinical Infectious Diseases. 2000; 31: 16-23.
6. Beytut E, Ozcan K, Erginsoy S. Immuno histochemical detection of fungal elements in the tissues of goslings with pulmonary and systemic aspergillosis. Acta Veterinaria Hungarica. 2004; 52(1): 71-84.
7. Humbert F. Salmonellosis and poultry industry: Epidemiological aspects and impact on public health. The Veterenary Point. 1992;24(145):201-206.
8. Bonny AC, Assandi KR, Karou ATG. Inhibition of the growth of multidrugs resistant avian *Salmonella* strains by aqueous and ethanolic extracts of *Mallotus oppositifolius* (Geisel.) Müll.-Arg (Euphorbiaceae). Int J Agri Biosci. 2021; 10(3): 128-133.
9. Agaie BM, Onyeyili PA, Muhammad BY, Landan MJ. Some toxic effects of aqueous leaf extract of *Anogeissus leiocarpus* in rats. Journal of Pharmacology and Toxicology. 2007; 2(4):396-401.
10. Adejumbi JA, Ogundiya MO, Kolapo A, and kunade MB. Phytochemical composition and *in vitro* antimicrobial activity of *Anogeissus leiocarpus* on some common oral pathogens. Journal of Medicinal Plants Research. 2008; 2(8):193-196.
11. Ndukwe IG, Amupitan JO, Isah YS, Adegoke K S. Phytochemical and

- antimicrobial screening of the crude extracts from the root, stem bark and leaves of *Vitellariaparadoxa*. Afr. J. Biotech. 2007; 6 (16): 1905-1909.
12. Stephen MC, Victoria IJ, Martins U, Yahaya S, Stephen AP, Joy AI, et al. Phytochemical screening and antibacterial activities of *Anogeissus leiocarpa* (DC.) Gull. and Perr. leaf extracts against *Escherichia coli* (T. Escherichia) and *Salmonella typhimurium* (Loeffler). Australian Journal of Science and Technology. 2020; 5: 527-532.
 13. Usman AH, Ali T, Danjani AG. Phytochemical and antimicrobial studies of stem-bark extracts of *Anogeissus leiocarpus* found in Dutsin-ma, Katsina-Nigeria. Fudma Journal of Sciences. 2020; 4: 156-167.
 14. Sale M, Pukuma MS, Hamza FA. (2020). *In vitro* Antibacterial activity of the bark extracts of *Anogeissus leiocarpus* on some bacterial pathogens. Nigerian Journal of Microbiology. 2020; 34(1): 4911 – 4919.
 15. Ouattara A, Golly KJ, Touré A. Investigation on traditional use of *Pericopsis* (afroormosia) *laxiflora* (Benth.) stem bark in treatment of infectious diseases caused by *Staphylococcus aureus* and *Shigella sp.*, two multi-resistants bacteria. Res J Pharm Biol Chem Sci. 2016; 7: 377-83.
 16. Zirihi GN, Kra AKM. Evaluation of the antifungal activity of *Microglossa pyrifolia* (Lamarck) O. Kuntze (Asteraceae) « PYMI » on the *in vitro* growth of *Candida albicans*. Rev Med Pharmacol Afr. 2003; 17: 11-18.
 17. Kuete V, Penlap-BV, Etoa F-x, Modjo SL, Bogne P, Assob JC, Lontsi D. Antimicrobial activities of total extract and fractions fruit juice of *Citrus medica* Lin. (Rutaceae). Pharm. Med. Trad. Afr. 2004;13:91-101.
 18. Bssaibis F, Gmira N, Meziane M. Antibacterial activity of *Dittrichia viscosa* (L.) W. Greuter. Rev. Microbiol. Ind. San. and Environn. 2009;3:44-55.
 19. Toure A, Bahi C, Ouattara K, Djaman AJ, Coulibaly A. Phytochemical screening and *in vitro* antifungal activities of extracts of leaves of *Morinda morindoides* (*Morinda*, *Rubiaceae*). Journal of Medicinal Plants Research. 2011; 5 (31): 6780-86.
 20. Biyiti LF, Meko'o DJL, Tamzic V, Amvam ZPH. Research of antibacterial activity of four Cameroonian medicinal plants. Pharm. Méd. Trad. Afr. 2004;(13):11-20.
 21. Prabhjot SJ, Wazbir HB, Avneet K, Ankit B, Loveleen K. Antibacterial and phytochemical analysis of ethnomedicinal plants. Research Journal of Pharmaceutical, Biological and Chemical Sciences. 2014;5:1203-1208.
 22. Bahman F-N, Hassan A, Mehrangiz G (2022). Effects of different solvents to extract phytochemical materials of *Rhazya stricta* against *Salmonella typhimurium* activity. Gene Cell Tissue. 2022;9(1):1-8.
 23. Naphtali E, Tahir F, Agbo EB. Evaluation of activities of some plant leaf extract on typhoidal and non-typhoidal *Salmonella* isolate from selected hospitals in Bauchi. Nigeria GSC Biological and Pharmaceutical Sciences. 2020;11(02): 020–030.
 24. Fauchere LL, Avril JL. General and Medical Bacteriology. Editions Ellipses. Paris; 2002
 25. Razieh S, Jalal S, Somayyeh H. Anti-quorum sensing and antibacterial activities of *Satureja sahendica* hydroalcoholic extract against avian isolate of *Salmonella Typhimurium*. Iranian Journal of Veterinary Science and Technology. 2020;1(22):71-87.
DOI: 10.22067/veterinary.v12i1.85433.
 26. Dahiru M, Adamu MT, Kolawole OS, Isyaka MS. *Anogeissus leiocarpus*: A potential inhibitor of some extended spectrum β -Lactamase bacteria. Acta Scientific Microbiology. 2021;4:71-81.
 27. Abbiw KD. (1990). Useful plants of Ghana, West Africa: Uses of Wild and Cultivated plants. Intermediate Technology Publication, UK. 1990;98-212.
 28. Amvam ZPH, Biviti L, Tchoumbovngang F, Menut C, Lamatv G, Bouchet PH. (1998). Aromatic plants of Tropical Central Africa, Part XXXI1 Chemical composition and antifungal activity of thirteen essential oils from aromatic plants of Cameroon. Flavour and Fragrance Journal. 1998; 13: 107-114.
 29. Jayasinghe L, Malika KB, Nishantha JK, Gayathri UN, Ratnayake B, Noriyuki H, Yoshinory F. Antifungal constituents of the stem bark of *Bridelia retusta*, Phytochemistry. 2003; 62: 637-641.

30. Bhuiyan MNI, Chowdhury JU, Begum J. Chemical investigation of the leaf and rhizome essential oils of *Zingiber zerumbet* (L.) Smith from Bangladesh. *Bangladesh J. Pharmacol.* 2009;4:9-12.

© 2023 Ouattara et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106063>