



# Impact of Different Doses of Herbicides on Weed Density and Weed Control Efficiency in Maize

Shivendra Singh <sup>a++\*</sup>, Ram Ashish Yadav <sup>a#</sup>,  
Sunil Kumar Prajapati <sup>b++</sup>, Pradeep Kumar <sup>a++</sup>,  
Praveen Kumar Yadav <sup>a++</sup>, Naushad Khan <sup>a#</sup>  
and Prashun Sachan <sup>a++</sup>

<sup>a</sup> Department of Agronomy, C.S.A. University of Agriculture and Technology, Kanpur-208002, India.

<sup>b</sup> Division of Agronomy, ICAR-IARI, New Delhi-110012, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2024/v36i54539

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114591>

Original Research Article

Received: 13/01/2024

Accepted: 22/03/2024

Published: 29/03/2024

## ABSTRACT

A field experiment was conducted at Chandra Shekhar Azad University of Agriculture and Technology, Kanpur during Kharif season of 2021. To evaluate the effect of different doses of Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC on growth and yield of maize (*Zea mays* L.). The experiment was laid down in the Randomized block design with ten treatments which was replicated three times. Experimental Field was infested with *Echinochloa colona*, *Digitaria sanguinalis*, *Dactyloctenium aegyptium*, *Commelina benghalensis*, *Trianthema* and *Cyperus rotundus*. Ready-mix herbicide, Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC (*Calaris Xtra*) manufactured by *Syngenta* was used in the experiment. Among the herbicidal treatments, the

<sup>++</sup> Research Scholar;

<sup>#</sup> Professor;

\*Corresponding author: E-mail: shivendrduat@gmail.com;

application of Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC @ 1.750 kg a.i. ha<sup>-1</sup> recorded the lowest weed density and highest weed control efficiency (approx. 90%) at every growth stage of maize. However, among herbicidal treatments, grain yield of maize was maximum with the application of Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC @ 1.0 kg a.i. ha<sup>-1</sup>(T4) which might be due to toxicity effect of herbicide.

**Keywords:** Doses; ready-mix; herbicide.

## 1. INTRODUCTION

Maize (*Zea mays* L.) is a key cereal and adaptable crop in the *Poaceae* family. Its relevance covers many uses, which include human dietary needs, animal and poultry feed, and industrial utilisation for the production of maize starch, dextrose, maize syrup, and maize flakes [1]. Being a C4, plant is capable of utilizing solar radiation more efficiently than several other cereal crops [2]. It is grown in many agro-ecological zones worldwide, with the United States, China, Brazil, and Mexico being the top producers [3].

Maize, after rice and wheat, is India's most important cereal crop in terms of food security and agricultural revenue. The majority of maize is grown during the kharif season, and weed infestation is one of the most critical yield-limiting factors. However, the first six weeks following crop planting are the most essential period for crop weed competition, since initial sluggish growth in wider spacing of maize, along with favourable weather conditions, allow luxuriant weed development, which may cut production by 28-100% (Dass et. al., 2012).

Weeds alone represent for one-third of the entire projected productivity losses caused by pests, insects, diseases, and weeds globally. Weeds are the most major factor limiting output in maize farming in India. Weed infestation in maize is fairly prevalent, and it typically leads into a large

reduction in production, especially in dry land conditions [4]. Chemicals are more efficient in maize fields because they can control weeds before they develop, kill both vegetative and deep-rooted weeds, and are also more cost-effective and produce results rapidly [5].

## 2. MATERIALS AND METHODS

### 2.1 Experimental Site

The research was conducted at 'Student's Instructional Farm, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur. The experimental field had an even topography and good drainage facility. Geographically, experimental site falls under the sub-tropical and semi- arid tract of North India of Indo- Gangetic plains and lies on the right bank of holy river Ganga. It is located on 26° 28' 36" N latitude, 80° 18' 26" E longitude and at an altitude of 126 meters above mean sea level. The experimental plot was homogenous in fertility with assured irrigation and other required facilities.

### 2.2 Experimental Design and Treatment Details

The experiment was designed as Randomized block design (RBD) with 10 Treatments replicated thrice. The treatment was allocated randomly in each block. The treatment details are given in Table 1.

**Table 1. Treatment details**

T1	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.375 kg a.i. ha <sup>-1</sup>
T2	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.75 kg a.i. ha <sup>-1</sup>
T3	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.875 kg a.i. ha <sup>-1</sup>
T4	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.0 kg a.i. ha <sup>-1</sup>
T5	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.125 kg a.i. ha <sup>-1</sup>
T6	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.750 kg a.i. ha <sup>-1</sup>
T7	Atrazine 50% WP @ 1.0 kg a.i. ha <sup>-1</sup>
T8	Hand weeding at 15 and 30 DAS
T9	Weed Free
T10	Control (Weedy check)

(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) is ready mix or premix herbicide bearing trade name Calaris Xtra manufactured by Syngenta.

### 2.3 Seed Sowing and Spacing

The field was ploughed with a tractor drawn cultivator and after with the rotavator to obtain a fine tilth. The seed was sown at the spacing of 45 cm between rows and 15 cm between plant. The Maize Variety *Azad Uttam* was sown and the applied seed rate was 25 kg ha<sup>-1</sup>. Thinning and gap filling was after 20 days after sowing wherever it was required for maintaining optimum plant population.

### 2.3 Application of Herbicide

Herbicide application was made at 25 Days after sowing (DAS) of maize crop with the knapsack sprayer.

### 2.4 Weed Density (No. /m<sup>2</sup>)

The weed density of different species was recorded at 15, 30, 45 and 60 days after application (DAA) of herbicides.

## 3. RESULTS AND DISCUSSION

### 3.1 Weed Studies

Weed flora of the experimental plot were collected, identified and different species of weeds were observed during the course of investigation, which includes six species of weeds. *Echinochloa colona*, *Digitaria sanguinalis*, *Dactyloctenium aegyptium*, *Commelina benghalensis*, among monocot weeds, *Trianthema* spp. is among dicot weeds, and *Cyperus rotundus* among sedges were the weed flora predominantly observed in the experimental site.

### 3.2 Effect of Treatments on Density (per m<sup>2</sup>) of *Cyperus rotundus*

The data in Table-2 revealed that the effect of herbicides was found significantly in reducing the population of *Cyperus rotundus* at 15, 30, 45 and 60 days after application of herbicides. Among the herbicide applications it was observed that the significantly minimum density of *Cyperus rotundus* was recorded under the application of (Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.750 kg a.i. ha<sup>-1</sup>. This result was in the line with Samant et al., (2015) and Dey et al., [6].

### 3.3 Effect of Treatments on Density (per m<sup>2</sup>) of *Echinochloa colona*

Data pertaining to density of *Echinochloa* is given in Table-3. Among the herbicide applications it was observed that the significantly minimum density of *Echinochloa colona* was recorded under the treatment (Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.750 kg a.i. ha<sup>-1</sup> (T6). Maximum density of *Echinochloa* was observed in the weedy check(T10) at each stage of crop growth period. This result was supported by Ispita Dey [7].

### 3.4 Effect of Treatments on Density (per m<sup>2</sup>) of *Digitaria sanguinalis* and *Trianthema* sp.

Data pertaining to density of *Digitaria sanguinalis* and *Trianthema* sp. is given in Table-4. Among the herbicide applications it was observed that the significantly minimum density of *Digitaria sanguinalis* and *trianthema* was recorded under the treatment T6 at each stage of crop growth while maximum density was recorded at Weedy check(T10).

Table 2. Density (per m<sup>2</sup>) of *Cyperus rotundus*

Treatment	Treatment details	15 DAA	30 DAA	45 DAA	60 DAA
T1	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.375 kg a.i. ha <sup>-1</sup>	4.40 (18.90)	5.49 (29.67)	5.88 (34.10)	7.09 (49.7)
T2	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.75 kg a.i. ha <sup>-1</sup>	3.45 (11.40)	4.77 (22.30)	5.62 (31.10)	6.47 (41.40)
T3	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.875 kg a.i. ha <sup>-1</sup>	3.05 (8.80)	4.17 (16.90)	4.89 (23.40)	5.68 (31.80)
T4	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.0 kg a.i. ha <sup>-1</sup>	2.83 (7.50)	3.83 (14.20)	4.52 (19.90)	5.22 (26.70)
T5	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.125 kg a.i. ha <sup>-1</sup>	2.61 (6.30)	3.56 (12.20)	4.37 (18.60)	5.11 (25.60)
T6	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.750 kg a.i. ha <sup>-1</sup>	2.27 (4.67)	3.24 (10.00)	4.12 (16.50)	5.01 (24.60)

Treatment	Treatment details	15 DAA	30 DAA	45 DAA	60 DAA
T7	Atrazine 50% WP @ 1.0 kg a.i. ha <sup>-1</sup>	3.55 (12.10)	4.67 (21.30)	5.59 (30.70)	6.43 (40.80)
T8	Hand Weeding @ 15 and 30 DAS	2.61 (6.30)	3.70 (13.20)	5.47 (29.40)	6.12 (37.00)
T9	Weed Free	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)
T10	Control	4.91 (23.60)	6.80 (45.80)	8.23 (67.20)	9.74 (94.40)
	SE(m)±	0.34	0.36	0.45	0.83
	C.D at 5%	1.03	1.07	1.34	1.21

# Data in the parenthesis are actual values.

**Table 3. Density (per m<sup>2</sup>) of *Echinochloa colona***

Treatment	Treatment details	15 DAA	30 DAA	45 DAA	60 DAA
T1	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.375 kg a.i. ha <sup>-1</sup>	3.94 (15.10)	4.34 (18.40)	4.63 (20.90)	5.15 (26.00)
T2	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.75 kg a.i. ha <sup>-1</sup>	3.24 (10.00)	3.75 (13.60)	4.12 (16.50)	4.58 (20.50)
T3	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 0.875 kg a.i. ha <sup>-1</sup>	3.16 (9.50)	3.58 (12.30)	3.86 (14.40)	4.31 (18.10)
T4	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.0 kg a.i. ha <sup>-1</sup>	3.16 (9.50)	3.56 (12.20)	3.74 (13.50)	4.12 (16.50)
T5	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.125 kg a.i. ha <sup>-1</sup>	3.03 (8.70)	3.33 (10.60)	3.59 (12.40)	3.97 (15.30)
T6	(Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.750 kg a.i. ha <sup>-1</sup>	2.33 (4.95)	2.94 (8.20)	3.32 (10.50)	3.87 (14.5)
T7	Atrazine 50% WP @ 1.0 kg a.i. ha <sup>-1</sup>	3.24 (10.00)	3.69 (13.10)	3.97 (15.30)	4.53 (20.00)
T8	Hand Weeding @ 15 and 30 DAS	2.79 (7.30)	3.30 (10.40)	4.32 (18.20)	4.70 (21.60)
T9	Weed Free	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)
T10	Control	4.17 (16.90)	5.33 (27.90)	6.08 (36.50)	6.75 (45.10)
	SE(m)±	0.31	0.32	0.51	0.54
	C.D at 5%	0.94	0.97	1.51	1.62

\*DAA (Days after application of herbicide)

# Data in the parenthesis are actual values.

**Table 4. Density (per m<sup>2</sup>) of *Digitaria sanguinalis* and *Trianthema* sp.**

Treatment	<i>Digitaria sanguinalis</i>				<i>Trianthema</i>			
	15 DAA	30 DAA	45 DAA	60 DAA	15 DAA	30 DAA	45 DAA	60 DAA
T1	3.14 (9.40)	4.14 (16.67)	4.36 (18.50)	5.04 (24.90)	3.0 (8.50)	3.72 (13.40)	4.05 (15.90)	4.68 (21.40)
T2	2.79 (7.30)	3.49 (11.70)	4.20 (15.70)	4.40 (18.90)	2.02 (3.60)	2.81 (7.40)	3.45 (11.40)	3.91 (14.80)
T3	2.70 (6.80)	3.24 (10.00)	3.73 (13.40)	4.05 (15.90)	1.73 (2.50)	2.39 (5.20)	2.77 (7.20)	3.29 (10.30)

Treatment	<i>Digitaria sanguinalis</i>				<i>Trianthema</i>			
	15 DAA	30 DAA	45 DAA	60 DAA	15 DAA	30 DAA	45 DAA	60 DAA
T4	2.65 (6.50)	3.18 (9.60)	3.59 (12.40)	3.86 (14.40)	1.64 (2.20)	2.26 (4.60)	2.66 (6.60)	2.98 (8.40)
T5	2.59 (6.20)	3.07 (8.90)	3.36 (10.80)	3.42 (11.20)	1.58 (2.00)	2.02 (3.60)	2.35 (5.00)	2.74 (7.00)
T6	2.09 (3.87)	2.82 (7.50)	3.15 (9.40)	3.66 (12.9)	1.47 (1.67)	1.81 (2.79)	2.19 (4.31)	2.79 (7.3)
T7	2.79 (7.30)	3.38 (10.90)	3.90 (14.70)	4.23 (17.40)	2.10 (3.90)	2.70 (6.80)	3.36 (10.80)	3.87 (14.50)
T8	2.65 (6.50)	3.19 (9.70)	3.74 (13.50)	4.28 (17.80)	1.52 (1.80)	2.07 (3.80)	2.95 (8.20)	3.63 (12.70)
T9	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)
T10	3.78 (13.8)	4.80 (22.50)	5.96 (35.00)	6.72 (44.70)	3.55 (12.10)	5.50 (29.80)	7.64 (57.90)	8.08 (64.80)
SE(m)±	0.27	0.34	0.35	0.52	0.23	0.39	0.47	0.59
C.D at 5%	0.81	1.02	1.05	1.56	0.68	1.18	1.42	1.77

# Data in the parenthesis are actual values.

Table-5. Density (per m<sup>2</sup>) of *Dactyloctenium aegyptium* and *Commelina benghalensis*

Treatment	<i>Dactyloctenium aegyptium</i>				<i>Commelina benghalensis</i>			
	15 DAA	30 DAA	45 DAA	60 DAA	15 DAA	30 DAA	45 DAA	60 DAA
T1	3.34 (10.67)	4.47 (19.56)	4.86 (23.10)	5.47 (29.40)	3.13 (9.30)	4.00 (15.50)	4.82 (22.70)	5.31 (27.70)
T2	2.63 (6.40)	3.62 (12.60)	4.31 (18.10)	4.96 (24.10)	2.43 (5.40)	3.29 (10.30)	4.06 (16.00)	4.60 (20.70)
T3	2.32 (4.90)	3.08 (9.00)	3.61 (12.50)	4.23 (17.40)	2.17 (4.20)	2.79 (7.00)	3.41 (11.10)	3.86 (14.40)
T4	2.10 (3.90)	2.83 (7.50)	3.29 (10.30)	3.81 (14.00)	2.02 (3.60)	2.61 (6.30)	3.18 (9.60)	3.59 (12.40)
T5	2.01 (3.55)	2.77 (7.20)	3.18 (9.60)	3.62 (12.60)	1.92 (3.20)	2.39 (5.20)	2.95 (8.20)	3.35 (10.70)
T6	2.00 (3.50)	2.70 (6.80)	2.98 (8.40)	3.24 (10)	1.81 (2.79)	1.98 (3.45)	2.87 (7.75)	2.98 (8.4)
T7	2.66 (6.60)	3.59 (12.40)	4.29 (17.90)	4.96 (24.10)	2.68 (6.70)	3.41 (11.10)	4.12 (16.50)	4.77 (22.30)
T8	1.95 (3.30)	2.74 (7.00)	3.95 (15.10)	4.47 (19.50)	1.90 (3.10)	2.49 (5.70)	3.56 (12.20)	4.27 (17.70)
T9	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)
T10	3.87 (14.50)	5.39 (28.50)	6.40 (40.50)	7.75 (59.60)	3.80 (14.10)	4.89 (23.40)	5.76 (32.70)	6.86 (46.60)
SE(m)±	0.25	0.38	0.43	0.58	0.29	0.34	0.40	0.51
C.D at 5%	0.75	1.13	1.29	1.75	0.86	1.02	1.21	1.54

# Data in the parenthesis are actual values.

### 3.5 Effect of Treatments on Density (per m<sup>2</sup>) of *Dactyloctenium aegyptium* and *Commelina benghalensis*

Data pertaining to density of *Dactyloctenium aegyptium* and *Commelina benghalensis* is

given in Table-5. Minimum density was observed in the T6 treatment in both the weeds at each stage of crop growth. The maximum density of *Commelina benghalensis* and *Dactyloctenium* was recorded under weedy check(T10).

Table 6. Weed control efficiency (%) at 15 DAA

Treatments	<i>Cyperus rotundus</i>	<i>Echinochloa colona</i>	<i>Digitaria sanguinalis</i>	<i>Trianthema sp.</i>	<i>Commelina benghalensis</i>	<i>Dactyloctenium aegyptium</i>
T1	81.20	75.39	73.71	81.97	64.42	82.67
T2	85.30	79.49	80.91	85.76	73.69	86.57
T3	88.98	81.08	82.73	86.56	79.80	90.01
T4	89.20	85.78	86.59	88.34	84.78	89.67
T5	89.68	88.63	89.32	89.38	86.43	91.00
T6	90.28	90.63	90.98	91.52	88.90	91.28
T7	82.61	77.15	78.73	84.57	67.36	84.56
T8	88.86	79.47	76.69	92.77	86.91	90.50
T9	100	100	100	100	100	100
T10	-	-	-	-	-	-
SE(m)±	0.94	0.98	1.01	1.36	1.34	1.09
C.D at 5%	2.80	2.91	3.02	4.04	3.98	3.25

### 3.6 Effects of Treatments on Weed Control Efficiency (%) in Maize

The data pertaining to weed control efficiency is given in Table-6. At 15 DAA highest weed control efficiency (100%) was obtained with Weed free plot followed by two hand weeding at 15 and 30 DAS. The result was in conformity with the findings of Sanodiya *et al.* [8]. Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) @ 1.750 kg a.i. ha<sup>-1</sup> attained highest WCE at 15 DAA. Nadiger *et al.* [9] Barla *et al.* [10] Malik [11] and Radheyshyam [12] also obtained similar results.

### 4. CONCLUSION

The application of Mesotrione 2.27% w/w + Atrazine 22.7% w/w SC) at the rate of 1.750 kg a.i. ha<sup>-1</sup> is most efficient in controlling weeds and recorded the lowest weed density and highest weed control efficiency of approximately 90% on an average at every growth stage of maize.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

- Sachan DS, Khan N, Maurya CL, Singh B. Influence of Different Herbicides on the Growth, Growth Attributes and Yield of Maize (*Zea mays* L.) under Central Plains Zone of Uttar Pradesh. Journal of Experimental Agriculture International. 2024;46(3):9-19.
- Deewan P, Mundra SL, Singh D, Meena M, Verma R, Sharma NK. Effect of weed and nutrient management on growth, productivity and protein content of quality protein maize (*Zea mays* L.). Journal of Pharmacognosy and Phytochemistry. 2017;6(1):271-274.
- Sachan DS, Reddy KJ, Saini Y, Rai AK, Singh O, Laxman T. Assessing Grain Yield and Achieving Enhanced Quality in Maize by Next Generation Fertilizer: A Review. International Journal of Environment and Climate Change. 2023;13(8):626-637.
- Sulewska HA, Koziara WI, Śmiatacz KA, Szymańska GR, Panasiewicz KA. Efficacy of selected herbicides in weed control of maize. *Fragm. Agron.* 2012;29(3): 144-151.
- Chikoye D, Udensi UE, Lum AF. Evaluation of a new formulation of atrazine and metolachlor mixture for weed control in maize in Nigeria. *Crop Protection.* 2005;24(11):1016-1020.
- Dey P, Pratap T, Mishra S, Sahoo SK, Ghosh S. Different Weed Management Practices and Their Effect on Soil Micro Flora in Spring Season Sweet Corn (*Zea mays saccharata* L.) in Tarai Region of Uttarakhand, India. *International Journal of Current Microbiology and Applied Science.* 2018;7(9):2319-7706
- Ispita dey Thesis, Krishikosh; 2020.

8. Sanodiya P, Jha AK, Shrivastava A. Effect of integrated weed management on seed yield of fodder maize; 2013.
9. Nadiger S, Babu R, Aravinda Kumar BN. Bioefficacy of pre-emergence herbicides on weed management in maize. Karnataka Journal of Agricultural Science. 2013; 26(1):17-19.
10. Barla S, Upasani RR, Puran AN, Thakur R. Weed management in maize. Indian Journal of Weed Science. 2016;48(1): 67–69.
11. Malik N. Performance of maize (*Zea mays* L.) and associated weeds under different herbicides schedule (Doctoral dissertation, GB Pant University of Agriculture and Technology, Pantnagar-263145 (Uttarakhand); 2015.
12. Radheshyam Evaluation of post emergence herbicides in maize (*Zea mays* L.). M.Sc. Thesis submitted to the ICAR-Indian Agricultural research Institute, New Delhi; 2018.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/114591>