



Screening of Antibacterial Activity from Mangrove Plants (*Rhizophora mucronata*) Leaf Associated Actinobacterium of Streptomyces Species

John Francis^a, Lakshminarayanan Arivarasu^{a*}, Pitchiah Sivaperumal^b and Lakshmi Thangavelu^a

^a Department of Pharmacology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai-600077, India.

^b Department of Pharmacology, Marine Biomedical and Environmental Health Research Laboratory-Blue Laboratory, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai-600077, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i62A35698

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77805>

Original Research Article

Received 12 October 2021

Accepted 27 December 2021

Published 28 December 2021

ABSTRACT

Introduction: Mangrove is a shrub or small tree that grows in coastal saline water mostly. The term is also used for tropical coastal vegetation consisting of special species. Mangroves occur worldwide mostly in the tropics and subtropics, mainly between latitudes 25° N and 25° S. The total mangrove forest area of the world was 137,800 square kilometres spanning 118 countries and territories

Materials and Methods: The fresh leaves of were collected from Pichavaram area, Tamilnadu. The leaves were washed thoroughly with tap water then surface sterilized with 1% of hydrogen peroxide. Then the sample was macerated in mortar and pestle.

Results and Discussion: The antibacterial activity of the rhizophora extract was analysed using two types of assays, namely disc diffusion method and Minimum Inhibitory Concentration (MIC) assay. The disc diffusion test was done for different concentrations for the selected clinical isolates (*Klebsiella*, *streptococcus* and *vibrio*).

Conclusion: From this study ,we can conclude that has good antibacterial activity against these microorganisms (*Klebsiella*, *streptococcus* and *vibrio*) and further studies can be done in future.

Keywords: Mangrove; antibacterial; actinobacteria; xenobiotics; wetlands.

1. INTRODUCTION

Mangrove is a shrub or small tree that grows in coastal saline water mostly. The term is also used for tropical coastal vegetation consisting of special species. Mangroves occur worldwide mostly in the tropics and subtropics, mainly between latitudes 25° N and 25° S. The total mangrove forest area of the world was 137,800 square kilometres spanning 118 countries and territories. Mangrove swamps found in tidal areas as mangrove ecosystems play an important role as refuge, feeding, breast feeding areas and sustain an extensive food web based on detritus. Marine invertebrates contain a variety of therapeutically bioactive compounds that have shown promise in the treatment of a variety of ailments [1]. High toxic compounds were isolated from ascidians that have the ability to alter resting membrane potential but not the formation of toxin-formed channels [2]. Additionally, mangroves export nutrients to adjacent marine ecosystems such as seagrass communities and coral reefs [3]. Because of washing of land-derived wastes containing metals primarily from industrial and agricultural sites, domestic wastes, and weathering action triggered by monsoonal floods, the bioaccumulation of metals in mangroves and salt marshes showed the highest accumulation in the monsoon season, followed by other seasons [4].

In the mangrove environment, numerous bacterial classes are usually present [5]. Mangroves of sea areas are generally nitrogen deficient but are nonetheless highly productive. Endophytes are major microorganisms that live inside of plants without causing any harm to their hosts which have been isolated from root nodules and leaves, stem and fruits of a wide variety of plant species [6]. Isolated endophytic bacteria taken from the branches of the two major species of mangrove plants, *Avicennia nitida* (white mangrove) and *Rhizophora mangle* (Red mangrove), are mostly found in two different mangrove areas of Brazil. Actinobacteria particularly, the sort Streptomyces, are significant makers of bioactive auxiliary metabolites. Following quite a while of screening, it has gotten progressively hard to find new antimicrobials from actinobacteria separated from normal soil conditions. These days, an ever increasing number of investigates are centered around extraordinary territories and outrageous conditions [7]., for example, desert, marine, and mangrove since microorganisms in uncommon

conditions need to create special guard component against the pressure from their living spaces and can develop versatile biosynthetic pathways for orchestrating novel natural mixtures [8] be told, countless new bioactive mixtures delivered by actinobacterial strains living in exceptional conditions have been found as of late [9]. Actinobacteria in the sea are one of the most productive classes of secondary metabolite producers, and they're essential for industry [10-13]. Thousands of biologically active secondary metabolites are produced by many members of the Actinomycetales group [14]. Since the 1950s, several essential antibiotics, anticancer, antitumor, and immunosuppressive agents have been studied and screened in Actinobacteria from terrestrial sources. However, in recent years, the repeated rediscovery of the same compounds from terrestrial actinobacteria has made them less appealing for screening programmes. At the same time, due to the structural diversity and unusual biological activities of their secondary metabolites, actinobacteria isolated from the marine environment are currently attracting a lot of attention. They are effective sources of secondary metabolites of antibacterial, antifungal, anticancer, antitumor, cytotoxic, cytostatic, anti-inflammatory, anti-parasitic, anti-malaria, antiviral, antioxidant, anti-angiogenesis, and other biological activities. In this review, an evaluation is made on the current status of research on marine actinobacteria yielding pharmaceutically active secondary metabolites [15]. Bioactive compounds derived from marine actinobacteria have distinct chemical structures that could be used to develop new medicines to fight resistant pathogens. Our team has extensive knowledge and research experience that has translate into high quality publications [16–22],[12],[23],24], [18,25–33]. With the advancement of science and technology, new bioactive compounds synthesised by actinobacteria from different marine sources will be in higher demand in the future.

2. MATERIALS AND METHODS

2.1 Collection of Plant Material and Preparation

The fresh leaves of were collected from Pichavaram area, Tamilnadu. The leaves were washed thoroughly with tap water then surface sterilized with 1% of hydrogen peroxide. Then the sample was macerated in mortar and pestle.

2.2 Isolation of Actinobacteria

Isolation and enumeration of actinobacteria were carried out in Kuster's agar medium (KUA) supplemented with 0.5% (W/v) NaCl. To minimize the fungal and bacterial contamination, KUA medium was supplemented with cycloheximide (10 µg/ml) and nalidixic acid (10 µg/ml) respectively (Kathiresan *et al.*, 2005). Freshly macerated samples were serially diluted and inoculated on KUA medium and incubated at 36°C for 7 days. The colonies were counted and the population density has been expressed as colony forming units per gram (CFU/g) of sediments. Morphologically distinct colonies were selected and pure cultures were obtained.

2.3 Identification of Marine Actinobacteria

2.3.1 Aerial mass colour

The colour of the mature sporulating aerial mycelium was recorded in naked eye. When the aerial mass colour fell between two colours series, both the colours were recorded. If the aerial mass colour of a strain to be studied showed intermediate tints, then also, both the colour series were noted. The media used were Yeast Extract-Malt Extract Agar and Inorganic-Salt Starch Agar.

2.3.2 Melanoid pigments

The grouping was made on the production of melanoid pigments (*i.e.* greenish brown, brownish black or distinct brown, pigment modified by other colours) on the medium. The strains were grouped as melanoid pigment produced (+) and not produced (-). In a few cases, the production of melanoid pigments was delayed or weak, and therefore, it was not distinguishable. This is indicated as variable (V). This test was carried out on the media ISP-1 and ISP-7, as recommended by the International *Streptomyces* Project (Shirling and Gottlieb, 1966).

2.3.3 Reverse side pigments

Reverse side pigment production of the isolate was determined on ISP7 medium. The pigment production was noted as distinctive (+) and not distinctive or none (-). In case, a colour with low chroma such as pale yellow, olive or yellowish brown occurred, it was included in the latter group (-).

2.3.4 Soluble pigments

Soluble pigment production of isolate was observed on ISP7 medium. The diffusible pigment production other than melanin was considered positive (+) and not produced (-). The colour was recorded (red, orange, green, yellow, blue and violet).

2.3.5 Spore chain morphology

Spore morphological characters of the strains were studied by inoculating a loopful of one week old cultures into solidified agar medium containing sterile glass slides. The cultures were incubated at 28±2°C and examined periodically for the formation of aerial mycelium, sporophore structure and spore morphology.

2.4 Chemotaxonomical Characteristics

2.4.1 Hydrolysis

Hydrolysis was done for releasing amino acids. Harvested cells of each strain weighing 20 mg (fresh) were placed in an ampo bottle and 1 ml of 6 N HCl was added and sealed with an alcohol blast burner. The samples were kept at 121°C for 20 h in a sand bath. The bottles were cooled by keeping them at a room temperature of 28±2°C. Hydrolysis was also done for releasing sugars. Harvested cells of each strain weighing 50 mg (fresh) were placed in an ampo bottle and 1 ml of 0.5N HCl was added and sealed with an alcohol blast burner. The samples were kept at 110°C for 2 h. The bottles were then cooled by keeping them at a room temperature of 28±2°C.

2.4.2 Thin layer chromatography (TLC)

Spotting of the whole cell hydrolysates was made carefully on TLC plate using a microliter pipette. Spots were 5-10 mm in diameter. This was done by multiple applications on the same spot of very small portions of the sample, which were dried by a hand dryer.

2.4.3 Amino acids

Each sample (3 µl) was applied on the baselines of the TLC plate (20 cm x 20 cm). Adjacent to this, 1 µl of DL-diaminopimelic acid (an authentic material mixture of DAP isomers) and 1 µl of amino acetic acid (glycine) were spotted as standards. The TLC plate was developed with the solvent system containing methanol: pyridine: glacial acetic acid: H₂O (5: 0.5: 0.125: 2.5 v/v). It

took approximately more than 4 h for development. The spots were visualized by spraying with 0.4% ninhydrin solution in water-saturated n-butanol, followed by heating at 100⁰ C for 5 min. Spots of amino acids ran faster than DAP. The sample spots were immediately compared with the spots of the standards since spots gradually disappeared in a few hours.

2.4.4 Whole-cell sugars

On a cellulose TLC plate (20 cm x 20 cm), 5µl of samples was spotted along with 3 µl of sugar solutions as standards on the same plates. Galactose, arabinose, xylose and madurose were the sugars, which were used as standards. The TLC plate was developed with the solvent mixture containing ethyl acetate: pyridine: acetic acid: distilled water (8: 5: 1: 1.5 v/v). The development time was more than 4 h. Spots were visualized by spraying with aniline phthalate reagent (3.25 g of phthalic acid dissolved in 2 ml of aniline and made upto 100 ml with water saturated n-butanol). The sprayed plate was heated at 100⁰ C for 4 min. Hexoses appeared as yellowish brown spots and pentoses, as maroon coloured spots.

2.4.5 Assimilation of carbon source

The ability of the actinobacterial strain in utilizing various carbon compounds as source of energy was studied, following the method recommended by International Streptomyces Project (Shirling and Gottlieb, 1966). Chemically pure carbon source certified to be free of admixture with other carbohydrates and contaminating materials were used for this purpose. Carbon sources for this test were Arabinose, Xylose, Inositol, Mannitol, Fructose, Rhamnose, Sucrose and Raffinose. These carbon sources were sterilized by ether sterilization without heating. The media and plates were prepared and inoculated according to the convention of ISP project (Shirling and Gottlieb, 1966). For each of the carbon sources, utilization is expressed as positive (+), negative (-), or doubtful (±). In the 'doubtful' strains, only a trace of growth slightly greater than that of the control was noticed.

2.4.6 Preparation of actinobacterial extraction

The isolated actinobacteria were inoculated on the nutrient broth and incubated at room temperature for 5-7 days. The culture was centrifuged (10000 rpm for 10 min) and the supernatant was extracted with equal volume of

ethyl acetate (Naik et al., 2001). After the evaporation of ethyl acetate, it was re-extracted with the equal amount of chloroform and concentrated under vacuum at room temperature.

2.5 Bacterial Suspension

The pathogenic bacteria *Klebsiella pneumonia*, *Streptococcus* sp. and *Vibrio cholerae* was collected from the Department of Microbiology, Saveetha medical college and hospital, Tamilnadu. The bacterial pathogens were cultured in Muller –Hinton Broth for 24 hr at room temperature. From this bacterial suspension was prepared with saline and the optical density was measured at 600 nm. The concentration of microbial suspension was fixed as 10⁸ CFU/ml. 1ml of suspension was spread over on Muller Hinton agar plate and incubated for 24hrs at ambient temperature.

2.6 Antibacterial Activity

The antibacterial activity of actinobacterial extract was performed with disc diffusion method. Whatman filter paper discs (5mm) were impregnated with various concentrations (0.5, 1, 1.5, 2, 2.5 and 3mg/ml) of leaf extract using ethanol and methanol solvent. The inoculated plates were incubated for 24 hr at room temperature and the inhibition zones around the discs were measured. All the results were expressed from an average of three with standard deviation.

2.7 Minimum Inhibitory Concentration

Minimal Inhibition Concentration of actinobacterial extract on ethanol and methanol was determined in 5 concentrations (0 -50 µg/ml / 0.001 to 0.1 mg /ml) with blank (extract in Muller Hinton broth). The inoculated bacteria in test tubes are incubated for 24hr in ambient temperature then the optical density was observed.

3. RESULTS

The antibacterial activity of the rhizophora extract was analysed using two types of assays, namely disc diffusion method and Minimum Inhibitory Concentration (MIC) assay. The Table 1 shows a disc diffusion test was done. For different concentrations for the selected clinical isolates (*Klebsiella, streptococcus and vibrio sp.*). For one gram of klebsiella for 0 µg/ml concentration of

the crude extract the zone of inhibition is 0, for 50 µg/ml concentration the zone of inhibition was 5 with standard error of ± 1.04 mm for 100 µg/ml concentration the zone of inhibition is 11 ± 2.6 mm, 150 µg/ml the zone of inhibition is 14 ± 1.4 mm for 200 µg/ml the zone of inhibition is found to be 16 ± 1.3 mm and for 250 µg/ml the zone of inhibition seen is 20 ± 2.4 mm for 300 µg/ml the inhibition was 28 ± 1.8 mm.

Table 1. Morphology of *streptomyces species* species isolated from

Colour of aerial mycelium	gray
Melanoid pigment	-
Reverse side pigment	-
Soluble pigment	+
Spore chain	Spirale
Assimilation of Carbon source	
Arabinose	+
Xylose	-
Inositol	+
Mannitol	-
Fructose	+
Rhamnose	-
Sucrose	+
Raffinose	+

Similarly was done for streptococcus, for 50 µg/ml the zone of inhibition is 3 ± 2.1 mm, for 100 µg/ml the inhibition seen was 8 ± 1.9 mm and for 150 µg/ml the zone of inhibition seen is 12 ± 2.4 mm for 200 µg/ml of the crude extract the zone of inhibition seen is 16 ± 1.3 for 250 µg/ml the zone of inhibition is 20 ± 1.7 mm and for 300 µg/ml the inhibition is 24 ± 2.1 mm For one gram of vibrio, for 50 µg/ml of the crude extract, the zone of inhibition seen is 4 ± 1.8 mm for 100 µg/ml the zone of inhibition seen is 9 ± 2.7 mm, for 150 µg/ml the zone of inhibition seen is around 13 ± 2.4 mm and for 200 µg/ml of the crude extract it shows the zone of inhibition of 17 ± 2.1 mm, for 250 µg/ml the zone of inhibition seen is 22 ± 2.4 mm and for 300 µg/ml the inhibition seen is 26 ± 2.6 mm. From the disc diffusion susceptibility test done for the highest concentration of 300 µg/ml of the crude extract, *Klebsiella* showed the highest zone of inhibition of the selected clinical isolates and *vibrio sp.* showed the second highest zone of inhibition and followed streptococcus..

Table 3 shows the Minimum Inhibitory Concentration (MIC) test was also done on the selected extracts with positive control as tetracycline. For *klebsiella* the MIC was found to be 30 µg/ml of the crude extract of the seaweed

and for tetracycline the MIC is seen to be 10 µg/ml. For *vibrio* and streptococcus species the MIC was found to be 20 µg/ml of the crude extract of the seaweed and for tetracycline the MIC for *E.coli* is seen to be 20 µg/ml. For the MIC was found to be 40 µg/ml of the crude extract of the seaweed and for tetracycline the MIC for *E.coli* is seen to be 20 µg/ml.

4. DISCUSSION

Overall gram-positive microorganisms are viewed as more delicate than gram-negative microscopic organisms towards various antimicrobial mixtures on account of the distinction in the design of their cell dividers [34] yet our outcome showed that the concentrates are successful against both gram-positive and gram-negative microbes. Previous research suggests that the unsaturated fat methyl ester concentrations of *E. agallocha* leaves possess important antibacterial properties. Antimicrobial properties of substances are alluring apparatuses in the control of undesired bacteria, especially in the treatment of contaminations and in the treatment of food waste [35]. The dynamic constituents of plants generally meddle with development and digestion of microorganisms in a negative [36]. In the previous study, ethanol removal showed the presence of a few phytochemicals, making it more dynamic against bacterial strains in contrast with the fluid concentrate. The antibacterial activity of the mixtures distinguished are now depicted by Patra et al. A study by G Sahoo showed that it had maximum zone of inhibition against streptococcus aureus [37-51].

Staphylococcus aureus, another MDR, was likewise hindered by the ethanol leaf concentrates of the relative multitude of three mangroves though in the event of ampicillin, the restraint zone was similarly less. The current study discovered more inhibition of *Staphylococcus aureus* by ethanol concentrate of *R. mucronata* and *E. agallocha* leaves [52]. Sivaperumal et al. screened the antibacterial movement of chloroform unrefined concentrate and segment fractionated compounds from leaves of *E. agallocha* and uncovered that contrasted with the unrefined concentrate, segment fractionated compounds were more dynamic against *S. aureus* and *S. pneumonia*. Albeit in his investigation, no endeavor was made with unadulterated concentrates, very encouraging outcomes were gotten against the MDR strains of *Staphylococcus aureus* and

Table 2. This table shows the zone of inhibition for different concentrations for the clinical isolated species (*Klebsiella, streptococcus, vibrio*)

$\mu\text{g/ml}$	Zone of inhibition		
	<i>Klebsiella</i> (mm)	<i>Streptococcus</i> (mm)	<i>Vibrio</i> (mm)
0	0	0	0
50	5 \pm 1.04	3 \pm 2.1	4 \pm 1.8
100	11 \pm 2.6	8 \pm 1.9	9 \pm 2.7
150	14 \pm 1.4	12 \pm 2.4	13 \pm 2.4
200	20 \pm 1.3	16 \pm 1.3	17 \pm 2.1
250	24 \pm 2.4	20 \pm 1.7	22 \pm 2.4
300	28 \pm 1.8	24 \pm 2.1	26 \pm 2.6

Table 3. This table shows the Minimum Inhibitory Concentration (MIC) for the selected clinical isolates (*klebsiella, streptococcus and vibrio*) with positive control as tetracycline

	Minimum Inhibitory Concentration						MIC ($\mu\text{g/ml}$)
	0	10	20	30	40	50	
<i>Klebsiella</i>	+	+	+	-	-	-	30 $\mu\text{g/ml}$
Tetracycline	+	+	-	-	-	-	20 $\mu\text{g/ml}$
<i>streptococcus sp</i>	+	+	-	-	-	-	20 $\mu\text{g/ml}$
Tetracycline	+	-	-	-	-	-	10 $\mu\text{g/ml}$
<i>Vibrio sp</i>	+	+	-	-	-	-	20 $\mu\text{g/ml}$
Tetracycline	+	-	-	-	-	-	10 $\mu\text{g/ml}$

Salmonella typhi. The potential for using marine-resistant actinobacteria to treat radionuclides was discovered in previous research. The morphing of big functional categories of organic pollutants that have been handled. Interactions between radionuclides and marine actinobacteria were also confirmed using actinobacterial cells(4). Other studies like antibacterial activity, antifungal activity, anti-inflammatory activity on both marine as well as using nanoparticles from plant extracts are done [53–61].

The previous examination additionally features the anti-toxin capability of *S. alba* and *E. agallocha* against *P. vulgaris* and *P. mirabilis* which are astute microbes showing opposition and moderate action, individually, to ampicillin. In general, the mangroves of Goa have enormous potential for the planning of novel anti-infection agents against the pathogenic microbes and particularly on multidrug-safe strains. Additionally, their antimicrobial likely should be reached out against a board of microorganisms answerable for most basic microbial infections. Further examination is important to recognize those bioactive mixtures, which will be a stage for clinical applications. actinobacteria are very notable to deliver a wide assortment of auxiliary metabolites like colors and antimicrobials, particularly *streptomyces* species produces numerous metabolites, for example, polyketides, peptides and polyketide-peptides half and halves

was checked for antifungal and safe concealment

5. CONCLUSION

In this study, associated actinobacteria *Streptomyces* species was checked and concluded that is showed potent antibacterial activity and further studies can be done by isolating its individual component and can be checked for various properties like antifungal, antioxidant and insecticidal activity [62,63]. These results can be compared with other actinobacterial species to enhance the property and can be used for pharmaceutical purpose

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENT

The authors would like to thank Saveetha Dental College and Saveetha Institute of Medical and Technical Sciences for their kind support to utilize the facilities for the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Farhana S, Shoichiro I, Yuji N. Molecular identification, micronutrient content, antifungal and hemolytic activity of starfish *Asterias amurensis* collected from Kobe coast, Japan [Internet]. African Journal of Biotechnology. 2017; 16:163–70. Available: <http://dx.doi.org/10.5897/ajb2016.15776>
2. Syed MH, Gnanakkan A, Pitchiah S. Exploration of acute toxicity, analgesic, anti-inflammatory, and anti-pyretic activities of the black tunicate, *Phallusia nigra* (Savigny, 1816) using mice model. Environ Sci Pollut Res Int. 2021;28(5):5809–21.
3. Ando Y, Mitsugi N, Yano K, Karube I. Isolation of a bacterium from mangrove soil for degradation of sea sludge. Appl Biochem Biotechnol. 2001;95(3):175–82.
4. Sivaperumal P, Kamala K, Rajaram R. Adsorption of cesium ion by marine actinobacterium *Nocardiopsis* sp. 13H and their extracellular polymeric substances (EPS) role in bioremediation. Environ Sci Pollut Res Int. 2018;25(5):4254–67.
5. Holguin G, Gonzalez-Zamorano P, de-Bashan LE, Mendoza R, Amador E, Bashan Y. Mangrove health in an arid environment encroached by urban development—a case study [Internet]. Science of The Total Environment. 2006;363:260–74. Available: <http://dx.doi.org/10.1016/j.scitotenv.2005.05.026>
6. Azevedo JL. Endophytic Fungi from Brazilian Tropical Hosts and Their Biotechnological Applications [Internet]. Microbial Diversity and Biotechnology in Food Security. 2014;17–22. Available: http://dx.doi.org/10.1007/978-81-322-1801-2_2
7. Stackebrandt E, Schumann P. Introduction to the Taxonomy of Actinobacteria [Internet]. The Prokaryotes. 2006;297–321. Available: http://dx.doi.org/10.1007/0-387-30743-5_16
8. Manivasagan P, Venkatesan J, Kim S-K. Introduction to Marine Actinobacteria [Internet]. Marine Microbiology. 2013;1–19. Available: <http://dx.doi.org/10.1002/9783527665259.ch01>
9. Anandan R. An Introduction to Actinobacteria. 2016.
10. Devi VS, Subathra Devi V, Gnanavel BK. Properties of Concrete Manufactured Using Steel Slag [Internet]. Procedia Engineering. 2014;97:95–104. Available: <http://dx.doi.org/10.1016/j.proeng.2014.12.229>
11. Gupta P, Ariga P, Deogade SC. Effect of Monopoly-coating Agent on the Surface Roughness of a Tissue Conditioner Subjected to Cleansing and Disinfection: A Contact Profilometric Study. Contemp Clin Dent. 2018;9(Suppl 1):S122–6.
12. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. Microb Pathog. 2018;117:68–72.
13. Needhidasan S, Samuel M, Chidambaram R. Electronic waste - an emerging threat to the environment of urban India. J Environ Health Sci Eng. 2014;12(1):36.
14. Baltz R. Renaissance in antibacterial discovery from actinomycetes [Internet]. Current Opinion in Pharmacology. 2008;8:557–63. Available: <http://dx.doi.org/10.1016/j.coph.2008.04.008>
15. Manivasagan P, Kang K-H, Sivakumar K, Li-Chan ECY, Oh H-M, Kim S-K. Marine actinobacteria: an important source of bioactive natural products. Environ Toxicol Pharmacol. 2014;38(1):172–88.
16. Rajeshkumar S, Kumar SV, Ramaiah A, Agarwal H, Lakshmi T, Roopan SM. Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. Enzyme Microb Technol. 2018;117:91–5.
17. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly

- synthesized nanoparticles on hazardous dyes degradation. *Biocatal Agric Biotechnol.* 2019;19:101138.
18. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.-mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res.* 2020;27(8):8166–75.
 19. Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using *Gymnema sylvestre* leaf extract and evaluation of its antibacterial activity [Internet]. *South African Journal of Chemical Engineering.* 2020;32:1–4. Available:<http://dx.doi.org/10.1016/j.sajce.2019.11.005>
 20. Rajasekaran S, Damodharan D, Gopal K, Rajesh Kumar B, De Poures MV. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel.* 2020;277:118166.
 21. Santhoshkumar J, Sowmya B, Venkat Kumar S, Rajeshkumar S. Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*. *S Afr J Chem Eng.* 2019;29:17–23.
 22. Raj R K, D E, S R. β -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line. *J Biomed Mater Res A.* 2020;108(9):1899–908.
 23. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol.* 2019;38(6):694–702.
 24. Ezhilarasan D, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int.* 2018;17(3):192–7.
 25. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, Rajeshkumar S. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of *Tamarindus indica* on MCF-7 human breast cancer cell line. *J Drug Deliv Sci Technol.* 2020;55:101376.
 26. Dua K, Wadhwa R, Singhvi G, Rapalli V, Shukla SD, Shastri MD, et al. The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress. *Drug Dev Res.* 2019;80(6):714–30.
 27. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol.* 2018;89(10):1241–8.
 28. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol.* 2021;122:105030.
 29. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry? *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2021;132(1):118–9.
 30. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. *Syzygium cumini* extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med.* 2019;48(2):115–21.
 31. Duraisamy R, Krishnan CS, Ramasubramanian H, Sampathkumar J, Mariappan S, Navarasampatti Sivaprakasam A. Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments. *Implant Dent.* 2019;28(3):289–95.
 32. Gnanavel V, Roopan SM, Rajeshkumar S. Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products. *Aquaculture.* 2019;507:1–6.
 33. Markov A, Thangavelu L, Aravindhan S, Zekiy AO, Jarahian M, Chartrand MS, et al. Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders. *Stem Cell Res Ther.* 2021;12(1):192.
 34. Scherrer R, Gerhardt P. Molecular Sieving by the *Bacillus megaterium* Cell Wall and Protoplast [Internet]. *Journal of Bacteriology.* 1971;107:718–35. Available:<http://dx.doi.org/10.1128/jb.107.3.718-735.1971>
 35. Agoramoorthy G, Chandrasekaran M, Venkatesalu V, Hsu MJ. Antibacterial and antifungal activities of fatty acid methyl esters of the blind-your-eye mangrove from India [Internet]. *Brazilian Journal of Microbiology.* 2007;38:739–42.

- Available: <http://dx.doi.org/10.1590/s1517-83822007000400028>
36. Aboaba OOA, Smith OO, Olude SI, FO. Antibacterial Effect of Edible Plant Extract on *Escherichia coli* 0157:H7 [Internet]. *Pakistan Journal of Nutrition*. 2006;5:325–7. Available:<http://dx.doi.org/10.3923/pjn.2006.325.327>
 37. Sahoo G, Mulla NSS, Ansari ZA, Mohandass C. Antibacterial Activity of Mangrove Leaf Extracts against Human Pathogens. *Indian J Pharm Sci*. 2012; 74(4):348–51.
 38. Pushpaanjali G, Geetha RV, Lakshmi T. Knowledge and Awareness about Antibiotic Usage and Emerging Drug Resistance Bacteria among Dental Students. *Journal of Pharmaceutical Research International*. 2020;34–42.
 39. Aathira CM, Geetha RV, Lakshmi T. Knowledge and Awareness about the Mode of Transmission of Vector Borne Diseases among General Public. *Journal of Pharmaceutical Research International*. 2020;87–96.
 40. Baskar K, Lakshmi T. Knowledge, Attitude and Practices Regarding HPV Vaccination among Undergraduate and Postgraduate Dental Students in Chennai. *Journal of Pharmaceutical Research International*. 2020;95–100.
 41. Manya Suresh LT. Wound Healing Properties of *Aloe Barbadensis* Miller-In Vitro Assay. *Journal of Complementary Medicine Research*. 2020;11(5):30–4.
 42. First Report on Marine Actinobacterial Diversity around Madras Atomic Power Station (MAPS), India [Internet]. [cited 2021 Aug 31]. Available:<http://alinteridergisi.com/article/first-report-on-marine-actinobacterial-diversity-around-madras-atomic-power-station-maps-india/>
 43. Physicochemical Profile of *Acacia Catechu* Bark Extract – An in Vitro Study - *International Journal of Pharmaceutical and Phytopharmacological Research* [Internet]. [cited 2021 Aug 31]. Available:<https://eijppr.com/article/physicochemical-profile-of-acacia-catechu-bark-extract-an-in-vitro-stud>
 44. Lakshmi T. Antifungal Activity of *Ficus racemosa* Ethanolic Extract against Dermatophytes-An in vitro Study. *Journal of Research in Medical and Dental Science*. 2021;9(2):191–3.
 45. Awareness of Drug Abuse among Teenagers - *International Journal of Pharmaceutical and Phytopharmacological Research* [Internet]. [cited 2021 Aug 31]. Available:<https://eijppr.com/article/awareness-of-drug-abuse-among-teenagers>
 46. Mangal CSK, Anitha R, Lakshmi T. Inhibition of Nitric oxide Production and Nitric oxide Synthase Gene Expression in LPS Activated RAW 264.7 Macrophages by Thyme oleoresin from *Thymus vulgaris*. *J Young Pharm*. 2018;10(4):481.
 47. COX2 Inhibitory Activity of *Abutilon Indicum* - *Pharmaceutical Research and Allied Sciences* [Internet]. [cited 2021 Aug 31]. Available: <https://ijpras.com/article/cox2-inhibitory-activity-of-abutilon-indicum>
 48. Jibu RM, Geetha RV, Lakshmi T. Isolation, Detection and Molecular Characterization of *Staphylococcus aureus* from Postoperative Infections. *Journal of Pharmaceutical Research International*. 2020;63–7.
 49. Sindhu PK, Thangavelu L, Geetha RV, Rajeshkumar S, Raghunandhakumar S, Roy A. Anorectic drugs: an experimental and clinical perspective -A Review. *Journal of Complementary Medicine Research*. 2020;11(5):106–12.
 50. Nivethitha R, Thangavelu L, Geetha RV, Anitha R, Rajeshkumar S, Raghunandhakumar S. In Vitro Anticancer Effect of *Sesamum Indicum* Extract -. *Journal of Complementary Medicine Research*. 2020;11(5):99–105.
 51. Mariona P, Roy A, Lakshmi T. Survey on lifestyle and food habits of patients with PCOS and obesity. *Journal of Complementary Medicine Research*. 2020;11(5):93–8.
 52. Abeysinghe PD. Antibacterial Activity of some Medicinal Mangroves against Antibiotic Resistant Pathogenic Bacteria. *Indian J Pharm Sci*. 2010;72(2):167–72.
 53. Kamala K, Sivaperumal P, Gopalakrishnan R, Swarnakumar NS, Rajaram R. Isolation and characterization of biologically active alkaloids from marine actinobacteria *Nocardiosis* sp. NCS1 [Internet]. *Biocatalysis and Agricultural Biotechnology*. 2015;4:63–9. Available:<http://dx.doi.org/10.1016/j.bcab.2014.10.005>
 54. Sivaperumal P, Kamala K, Rajaram R. Bioactive DOPA melanin isolated and characterised from a marine

- actinobacterium *Streptomyces* sp. MVCS6 from Versova coast [Internet]. *Natural Product Research*. 2015;29:2117–21. Available: <http://dx.doi.org/10.1080/14786419.2014.988712>
55. Sivaperumal P, Kamala K, Rajaram R, Mishra SS. Melanin from marine *Streptomyces* sp. (MVCS13) with potential effect against ornamental fish pathogens of *Carassius auratus* (Linnaeus, 1758) [Internet]. *Biocatalysis and Agricultural Biotechnology*. 2014;3:134–41. Available: <http://dx.doi.org/10.1016/j.bcab.2014.09.007>
56. Desai C, Patel P, Markande AR, Kamala K, Sivaperumal P. Exploration of haloarchaea for their potential applications in food industry [Internet]. *International Journal of Environmental Science and Technology*. 2020;17:4455–64. Available: <http://dx.doi.org/10.1007/s13762-020-02773-2>
57. Kumar P, Sivaperumal P, Manigandan V, Rajaram R, Hussain M. Assessment of potential human health risk due to heavy metal contamination in edible finfish and shellfish collected around Ennore coast, India [Internet]. *Environmental Science and Pollution Research*. 2021;28:8151–67. Available: <http://dx.doi.org/10.1007/s11356-020-10764-6>
58. Sivaperumal P, Ramasamy P, Jacob Inbaneson S, Ravikumar S. Screening of antibacterial activity of mangrove leaf bioactive compounds against antibiotic resistant clinical isolates. 2010;2(5):348–53.
59. Ananthan G, Sivaperuma P, Mohamed Hussain S. Antibacterial Potential of Marine Ascidian *Phallusia arabica* Against Isolated Urinary Tract Infections Bacterial Pathogens [Internet]. *Asian Journal of Animal Sciences*. 2011;5:208–12. Available: <http://dx.doi.org/10.3923/ajas.2011.208.212>
60. Kamala K, Sivaperumal P. Biomedical Applications of Enzymes From Marine Actinobacteria [Internet]. *Marine Enzymes Biotechnology: Production and Industrial Applications, Part III - Application of Marine Enzymes*. 2017:107–23. Available: <http://dx.doi.org/10.1016/bs.afnr.2016.11.002>
61. Sivaperumal P, Kamala K, Rajaram R. Bioremediation of Industrial Waste Through Enzyme Producing Marine Microorganisms [Internet]. *Marine Enzymes Biotechnology: Production and Industrial Applications, Part III - Application of Marine Enzymes*. 2017:165–79. Available: <http://dx.doi.org/10.1016/bs.afnr.2016.10.006>
62. Rajendran R, Kunjusankaran RN, Sandhya R, Anilkumar A, Santhosh R, Patil SR. Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An in Vitro Study. *Pesqui Bras Odontopediatria Clin Integr*. 2019;19(0):4668.
63. Kumar MS, Vamsi G, Sripriya R, Sehgal PK. Expression of matrix metalloproteinases (MMP-8 and -9) in chronic periodontitis patients with and without diabetes mellitus. *J Periodontol*. 2006;77(11):1803–8.

© 2021 Francis et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77805>