



Assessment of Groundnut (*Arachis hypogaea* L.) Varieties under Various Sowing Windows for Crop Growth, Development and Yield in Western Maharashtra, India

Wankhede Sachin Yadavrao^{a*},
Kharbade Sattappa Bhambhar^{b++}, Misal Suraj Satish^{c,d#},
Jadhav Jaywant Dadaji^{e†}, Abdul Aziz Shaikh^{f‡}
and Sthool Vijay Atmaram^{g^}

^a Department of Agricultural Meteorology, College of Agriculture, Pune-411005, Mahatma Phule Krishi Vidyapeeth, Rahuri-413722, Maharashtra, India.

^b College of Agriculture, Karad-415 110, Mahatma Phule Krishi Vidyapeeth, Rahuri-413722, Maharashtra, India.

^c Krishi Vigyan Kendra, Mohol, Solapur, 413 213, Mahatma Phule Krishi Vidyapeeth, Rahuri, 413722, Maharashtra, India.

^d Mahatma Phule Krishi Vidyapeeth, Rahuri-413722, Maharashtra, India.

^e Agricultural Technical School, Solapur-413 002, Mahatma Phule Krishi Vidyapeeth, Rahuri-413722, Maharashtra, India.

^f Oilseeds Research Station, Jalgaon, 425 001, Mahatma Phule Krishi Vidyapeeth, Rahuri-413722, Maharashtra, India.

^g Department of Agricultural Meteorology, College of Agriculture, Pune-411005, Mahatma Phule Krishi Vidyapeeth, Rahuri-413722, Maharashtra, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i81948

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99837>

⁺⁺ Associate Dean;

[#] SMS Agrometeorology;

[†] Principal;

[‡] Groundnut Agronomist;

[^] Head;

*Corresponding author: E-mail: wankhedesachin04@gmail.com;

ABSTRACT

The present experiment was carried out at Department of Agricultural Meteorology Farm, Centre of Advanced Faculty Training, College of Agriculture, Pune, MPKV, Rahuri for assessment of different groundnut (*Arachis hypogaea* L.) varieties for crop growth, development and yields under various sowing windows in Western Maharashtra Plain Zone during the Kharif season of 2017 and 2018. The experiment was laid out in split plot design with three replications. The treatment comprised of four varieties viz., V₁: JL-501, V₂: RHRG-6083 (*Phule Unnati*), V₃: TAG-24 and V₄: JL-776 (*Phule Bharati*) as main plot and four sowing windows viz., S₁: 25th MW, S₂: 26th MW, S₃: 27th MW and S₄: 28th MW as sub plot treatments. From the result of the study, all the growth attributes were increased with the advancement in age of the groundnut crop. Plant height 35.27 and 33.34 cm, total number branches 11.34 and 10.55, total dry matter accumulation plant⁻¹ 34.36 and 32.11 g during both the years of experiment, which were found significantly higher in variety JL-776 over RHRG-6083, JL-501 and TAG-24. Among all the sowing windows 26th MW (S₂) recorded the highest growth attributes viz., plant height (33.67 and 31.94 cm), total number of branches plant⁻¹ (11.34 and 10.58 g) and total dry matter accumulation plant⁻¹ (34.75 and 32.78 g) during both year of experiment. 26th MW sowing window was at par with the 27th MW sowing window with all growth attributes. Pod yield (26.59 and 28.14 q ha⁻¹) and haulm yield (39.61 and 36.7 q ha⁻¹) were significantly higher in variety in JL-776 followed by RHRG-6083, JL-501 and TAG-24. Pod yield (27.25 and 28.84 q ha⁻¹) and haulm yield (40.60 and 37.61 q ha⁻¹) was higher in 26th MW sowing window, which were at par with 27th MW sowing window during both the year.

Keywords: Crop weather relationship; growth; phenology; post-harvest character; sowing date; yield.

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L., Fabaceae) is essentially a tropical plant and requires a long and warm growing season. The favourable climate for groundnut is a well distributed rainfall of at least 500 mm during the crop growing season with abundance of sunshine and relatively warm temperature, which is essential for maximum yield and quality of groundnut. Groundnut crop grows under wide range of temperatures but requires a long and warm growing season. Sowing, flowering and reproductive phases require an average rainfall distribution of 100, 150 and 400-500 mm, respectively. Moisture stress during vegetative period delays flowering, pod setting and results in low yields. High atmospheric humidity stimulates more flowering which results in increased peg setting. Pod development stage is most sensitive to moisture deficit. Pod yield is significantly influenced by day length. Long days promote vegetative growth at the expense of reproductive growth and increased crop growth rate resulting in decreased partitioning of photosynthesis to pods and decreased duration of effective pod filling phase [1]. Sowing of

rained and irrigated crop early in the season provide favourable weather conditions for proper growth and yield of groundnut. Delay in sowing by one-week results in linear decrease in pod yield of groundnut. In normal-sown crop, the pattern of flowering is regular with two distinct peaks of flowering, whereas in late-sown crop erratic pattern of flowering occurs [2]. The choice of a groundnut variety for any particular area depends on matching the variety with the length of the growing season. Groundnut varieties whose growth cycle is longer than the duration of growing season at a particular location either fail to mature or mature at a time when soil is too hard to dig the pods, therefore the investigation is carried out to assess the different groundnut varieties for crop growth, development and yields under various sowing dates in Western Maharashtra.

2. MATERIALS AND METHODS

2.1 Location of the Experimental Site, Soil and Climatic Condition

The field experiment was conducted for two consecutive years at Department of Agricultural

Meteorology farm, College of Agriculture, Pune during *kharif*, 2017 and 2018, India. The geographical location of the site (Pune) was 18° 32'N, latitude; 73°51'E, longitude and 557.7 MASL. The experimental site is situated in the sub-tropical region (Plain Zone) in the Pune. The soil is medium black having depth of about 1m. The average annual rainfall of Pune is 675 mm, which is distributed from second fortnight of June to second fortnight of October. Out of total rainfall, about 75 per cent is received from June to September from south-west monsoon, while remaining is received from north-east monsoon during October and November.

2.2 Nature of Season during Experimental Period

Daily and weekly mean meteorological data during the crop growth period (25th to 45th MW) of *kharif* 2017 and 2018 recorded in class 'A' observatory situated in the adjoining field. The daily maximum and minimum temperature during the crop growth period ranged from 34.4 and 12.7 °C during *kharif* 2017 and 34.7 and 13.3 °C during *kharif* 2018. During crop period, the weekly maximum and minimum temperatures varied from 27.1 to 33.4 °C and 14.1 to 23.9 °C, respectively, during 2017. It was varied from 24.2 to 37.7 °C and 13.3 to 24.6 °C respectively, during *kharif* 2018. Weekly relative humidity during morning (07.20 h LMT) and afternoon (14.20 h LMT) was 93.7 and 31.1 % in 2017, whereas it was 97 and 16 % in *kharif* 2018, respectively. The daily range of relative humidity during morning was 75-97 % and 72-97 % during the respective years while during afternoon was in the range of 26-98 %, while, it was between 16-92 % during two years of experimentation, respectively. The weekly wind velocity during the period ranged from 1.6 to 10.3 and 1.1 to 11.6 kmph during 2017 and 2018, respectively. The bright sunshine hour's day⁻¹ during crop growing period were 9.3 and 10.5 during 2017 and 2018, respectively. The weekly evaporation ranged from 2.2 to 6.7 and 2.2 to 5.3 mm per day in 2017 and 2018, respectively. The weekly photoperiod *i.e.* maximum possible sunshine hours which were fixed for the particular day in a year ranged from 10.38 to 13.87.

2.3 Experimental Details

The experiment was laid out in split plot design with three replications. The treatment comprised of four varieties *viz.*, V₁: JL-501, V₂: RHRG-6083

(*Phule Unnati*), V₃: TAG-24 and V₄: JL-776 (*Phule Bharati*) as main plot and four sowing windows *viz.*, S₁: 25rd MW (18th to 24th June), S₂: 26th MW (25th June to 01st July), S₃: 27th MW (2nd to 8th July) and S₄: 28th MW (09th to 15th July) as sub plot treatments. The gross and net plot size was 4.5 x 4.5 m² and 3.6 x 3.6 m², respectively. The allocation of treatments was done with random method. The certified seed of all the groundnut varieties was procured from the Groundnut Breeder, Oilseed Research Station, Jalgaon, MPKV, Rahuri. Sowing was done as per the treatments by dibbling one kernel at each hill with 30 cm inter-row and 7.5 cm intra-row distance keeping a seed rate of 100 kg ha⁻¹. The requisite plant population was maintained by thinning and gap filling where ever necessary. Urea and single super phosphate were used as source of N and P and applied as per recommended dose *i.e.* 25 kg N and 50 kg P₂O₅. Seed of groundnut was inoculated with *Rhizobium* culture @ 250 g 10 kg⁻¹ seed.

2.4 Growth Studies

Plant height (cm) was measured from the base of main stem to the base of apical bud of the plant at the end of various physiological growth stages, Number of branches Plant⁻¹ was counted from each observation plant from branching stage to physiological maturity, Dry matter accumulation plant⁻¹ (g) plant parts were sundried and then allow drying in thermostatically controlled hot air oven at 60 ± 2°C till constant weight was recorded, the dry matter study was carried out at the end of each growth stage during the period experiment.

2.5 Phenological Studies

The number of days from sowing to 50 per cent of the plants in the plot showed flowering was recorded by observing plants from net plot. The number of days required from sowing to physiological maturity of pods was recorded by observing plants from net plot.

2.6 Yield Studies

Pods from net plot were stripped after uprooting and dried in sunlight. The pod yield obtained was recorded as per treatments and pod yield⁻¹ was calculated. After stripping pods, haulms from net plot area of each treatment were sun dried and expressed quintal⁻¹.

3. RESULTS AND DISCUSSION

The biometric observations of groundnut were recorded on various growth characters *viz.*, plant height (cm) (Table 1), number of branches plant⁻¹ (Table 2), dry matter (g) (Table 3), days to 50 per cent flowering and days to maturity (Table 4) at regular interval of 28, 42, 56, 70, 84 days after sowing (DAS) and at harvest.

3.1 Plant Height (cm) (Table 1)

The mean plant height increased as the crop advanced in age. The mean plant height increased from 28 DAS (4.35 cm) to harvest (32.13 cm) during 2017 and from 28 DAS (3.91 cm) to harvest (30.23 cm) during 2018

3.1.1 Effect of varieties (Table 1A)

The mean plant height differences due to varieties were significant at all the stages of crop growth. It could be observed that at all the stages of crop growth, variety JL-776 (V₄) recorded significantly higher plant height (35.27 and 33.34 cm). This was followed by RHRG-6083 (V₂) (34.19 and 32.34) and JL-501 (V₁) (29.90 and 27.99 cm) during the year, 2017 and 2018, respectively. Plant height in variety TAG-24 (V₃) (29.15 and 27.26 cm) was found significantly the lowest among all varieties at all the growth stages of crop. This suggested that the variety JL-776 suitably acclimatize in soil and agro climatic conditions of Pune region thus, exploiting its potential in term of growth and development. Further, increase in plant height in groundnut is genetically governed phenomena affecting hormonal balance, nutrient absorbing capacity and photosynthetic ability. All these processes were reflected in height and other yield contributing characters. Increase in plant height due to favourable weather situations like higher GDD, photo thermal and heliothermal units which facilitates better growth (Table 1A).

3.1.2 Effect of sowing windows (Table 1B)

The mean plant height was significantly influenced throughout the crop growth period by different growing environments created through different sowing windows. The maximum plant height (34.15 and 32.27 cm) was recorded in 26th MW (S₂) sowing window and it was at par with 25th MW sowing window (32.90 and 30.94 cm) i.e. (S₁) this was followed by 27th MW sowing window (31.52 and 29.64 cm) and 28th MW sowing window (29.94 and 28.07 cm) during the

year 2017 and 2018, respectively. The plant height was increased very fast in between 28 to 70 DAS which might be due to active vegetative growth phase of the plant and at slow rate beyond 70 DAS as the plant enter from initial vegetative phase to reproductive phase. It can be concluded that plant height in the second sowing window (S₂) was significantly superior as compared to rest of the sowing windows during the crop growth period. It might be due to suitable weather conditions of different weather parameters during crop growing period. Plant height decreased significantly with delay in sowing. Reduced plant height with delay in sowing may be due to quick changes in photoperiod, which accelerated development towards reproductive stage and hence less time was available for vegetative growth. These results are similar to those reported by [3] (Table 1B).

3.1.3 Effect of interaction (Table 1C)

The plant height (cm) at all the stages of crop growth was significantly influenced by interaction between varieties and sowing windows during both the years. 26th MW sowing window (S₂) recorded higher plant height (36.55 and 38.48 cm) in variety JL-776 (V₄) which was followed by variety RHRG-6083 (V₂) (33.65 and 35.48 cm), JL-501 (V₁) (29.98 and 31.81 cm) and TAG-24 (V₃) (28.91 and 30.84 cm) during both the years 2017 and 2018, respectively (Table 1C).

3.2 Number of Branches Plant⁻¹ (Table 2)

The mean number of branches plant⁻¹ was progressively increased with advancement of the age of the crop, 10.27 and 9.44 at harvest during the year, 2017 and 2018, respectively.

3.2.1 Effect of varieties (Table 2A)

The mean number of branches plant⁻¹ as influenced due to varieties was significant at all the stages of crop growth. It could be observed that significantly higher number of branches plant⁻¹ was observed in var. JL-776 (V₄) (11.34 and 10.55) followed by var. RHRG-6083 (V₂) (10.73 and 9.89) and JL=501 (V₁) (9.62 and 8.78). Lower number of branches plant⁻¹ was observed in var. TAG-24 (V₃) (9.39 and 8.55 cm) at harvest during the year 2017 and 2018, respectively. This difference in varietal performance may be attributed to genotypic variation affecting branching pattern and duration (Table 2A).

3.2.2 Effect of sowing windows (Table 2B)

The maximum number of branches plant⁻¹ of groundnut (11.34 and 10.58 cm) was recorded in 26th MW (S₂) sowing window and it was at par with 25th MW sowing window (10.61 and 9.77 cm) *i.e.* (S₁) which was followed by 27th MW sowing window (10.01 and 9.17 cm) and 28th MW sowing window (9.12 and 8.25 cm) during 2017 and 2018, respectively. The number of branches plant⁻¹ of groundnut was increased very fast in between 28 to 56 DAS which might be due to active vegetative growth phase of the plant and at slow rate beyond 56 DAS as the plant enter from initial vegetative phase to reproductive phase. It was observed that number of branches plant⁻¹ in the sowing window (S₂) was significantly superior as compared to rest of the sowing windows during the crop growth period. It might be due to suitable weather conditions of different weather parameters during crop growing period. Number of branches plant⁻¹ decreased significantly with delay in sowing. Reduced number of branches plant⁻¹ with delay in sowing may be due to quick changes in photoperiod, which accelerated development towards reproductive stage and hence less time was available for vegetative growth [4] and [5] (Table 2B).

3.2.3 Effect of interaction (Table 2C)

The number of branches plant⁻¹ of groundnut at all the stages of crop growth was significantly influenced by interaction between varieties and sowing windows during both the years. 26th MW sowing window (S₂) recorded higher number of branches plant⁻¹ of groundnut (12.45 and 12.98 cm) in var. JL-776 (V₄) which was followed by var. RHRG-6083 (V₂) (10.80 and 11.64 cm), JL-501 (V₁) (9.75 and 10.59 cm) and TAG-24 (V₃) (9.31 and 10.15 cm) during the year 2017 and 2018, respectively (Table 2C).

3.3 Total Dry Matter Accumulation Plant⁻¹ (g) (Table 3)

During the crop growing period, increase in dry matter weight was continuous with the advancement in the crop age up to harvest of crop. The rate of increase was rapid during flowering and reproductive period. The mean dry matter of groundnut plant⁻¹ at harvest during both the years of 2017 and 2018, was 30.88 and 29.02 (g), respectively.

3.3.1 Effect of varieties (Table 3A)

The mean dry matter plant⁻¹ of groundnut as influenced due to varieties were significant at all the stages of crop growth. It could be observed that at harvest, variety JL-776 (V₄) recorded significantly higher dry matter plant⁻¹ (g) of groundnut (34.36 and 32.81 g) followed by RHRG-6083 (V₂) (31.08 and 29.11 g) and JL-501 (V₁) (29.75 and 27.78 g), whereas it was lowest in var. TAG-24 (V₃) (28.33 and 26.36 g) at harvest of crop during 2017 and 2018, respectively. This was consequent to significant growth and yield parameters of variety JL-776 which exploited its full potential under present agro-climatic condition since, dry matter accumulation is the result of all growth and yield attributes (Table 3A).

3.3.2 Effect of sowing windows (Table 3B)

The maximum dry matter accumulation plant⁻¹ (34.75 and 32.78 g) was recorded in 26th MW (S₂) sowing window and it was at par with 25th MW sowing window (S₁) (31.67 and 29.7 g) *i.e.* (S₁) this was followed by 27th MW sowing window (29.58 and 27.61 g) and 28th MW sowing window (27.53 and 25.97 g) during the year 2017 and 2018, respectively. The dry matter accumulation plant⁻¹ of groundnut was increased very fast in between 28 to 56 DAS which might be due to active vegetative growth phase of the plant and at slow rate beyond 56 DAS as the plant enter from initial vegetative phase to reproductive phase. The dry matter accumulation plant⁻¹ of groundnut in the sowing window (S₂) was significantly superior as compared to rest of the sowing windows during the crop growth period. It might be due to suitable weather conditions of different weather parameters during crop growing period. Dry matter accumulation plant⁻¹ of groundnut decreased significantly with delay in sowing. Reduced dry matter accumulation plant⁻¹ of groundnut with delay in sowing may be due to quick changes in photoperiod, which accelerated development towards reproductive stage and hence less time was available for vegetative growth. At all the stages of growth, the dry matter weight plant⁻¹ (g) showed decreasing trend with late sowings (S₂ to S₄). Besides, growth period of the crop also decreased with each successive delay in sowing which was also caused reduction in dry matter accumulation in late sowings. From all the observations it is observed that the sowing window 26th (MW) *i.e.* treatment S₂ was significantly superior over all the other sowing window treatments. The vegetative growth and

dry matter production were less in extended sowing windows. This decreased growth period might have reduced dry matter production with late sowing. They indicated July 7 as the optimum date of sowing and observed the highest net assimilation ratio and photosynthesis ratio at this sowing date. Similar results were reported by [5] and [6] (Table 3B).

3.3.3 Effect of interaction (Table 3C)

The dry matter accumulation plant⁻¹ of groundnut at all the stages of crop growth was significantly influenced by interaction between varieties and sowing windows during both the years. 26th MW sowing window (S₂) recorded higher dry matter accumulation plant⁻¹ of groundnut (41.03 and 43 g) in variety JL-776 (V₄) which was followed by variety RHRG-6083 (V₂) (31.36 and 33.33 g), JL-501 (V₁) (31.03 and 33.0 g) and TAG-24 (V₃) (29.03 and 31.0 g) during the year 2017 and 2018 respectively (Table 3C).

3.4 Days to 50 Per Cent Flowering and Maturity (Table 4)

Data regarding mean days to 50 per cent flowering and maturity of groundnut as influenced significantly by the different treatments are presented. The mean days to 50 per cent flowering were 32.98 and 32.48 during the year 2017 and 2018, respectively. The mean days to maturity were 108.46 and 107.68 for 2017 and 2018, respectively.

3.4.1 Effect of varieties (Table 4A)

Days to 50 percent flowering and days to maturity were influenced significantly due to different groundnut varieties. Maximum number of days to 50 per cent flowering (36 and 35.63) was observed in variety JL-776 (V₄) which was significantly higher. This was followed by var. RHRG-6083 (V₂) (33.92 and 33.38) and JL-501 (V₁) (31.17 and 30.63), while var. TAG-24 (V₃) was recorded significantly lower (30.83 and 30.29) during the year 2017 and 2018, respectively. Maximum number of days to maturity (113.08 and 111.75) was observed in variety JL-776 (V₄) which was significantly higher. This was followed by RHRG-6083 (V₂) (111.42 and 110.66) and JL-501 (V₁) (106.58 and 103.91), while var. TAG-24 (V₃) was recorded significantly lower (102.75 and 102.41) during the year 2017 and 2018, respectively.

3.4.2 Effect of sowing windows (Table 4B)

The days to 50 per cent flowering (35.08 and 34.54) was observed significantly higher in 26th MW sowing window. This was followed by 25th MW sowing window at (33.67 and 33.13), 27th MW sowing window (32.67 and 32.13) and 28th MW sowing window (30.5 and 30.13) during the year 2017 and 2018, respectively. The days to maturity (111.83 and 110.91) was observed significantly higher in 26th MW sowing window. This was followed by 25th MW sowing window at (109.83 and 109.16), 27th MW sowing window (107.58 and 106.64) and 28th MW sowing window (104.58 and 104.3) during the year 2017 and 2018, respectively. The similar results were observed by [6] at the time of sowing groundnut in *Kharif* season and indicated that the sowing groundnut in the second fortnight of June and first fortnight of July gave highest flower production. Later sowings reduced flower production. In any given locality, early sowings in first fortnight of July resulted in early flowering and early maturity [7]. Similar result was recorded by [8] in the field trials conducted on pea (*Pisum* sp.) cultivars of early and late maturing varieties sown at seven days intervals and concluded that the July sowings emerged faster and required less time to reach flowering and early harvesting.

3.4.3 Effects of interaction (Table 4C)

The days to 50 per cent flowering and days to maturity was significantly influenced by interaction between varieties and sowing windows during the year 2017 and 2018. Sowing at 26th MW sowing window (S₂) recorded maximum days to 50 per cent flowering (37.79 and 38.33) in variety JL-776 (V₄). This was followed by variety RHRG-6083 (V₂) (35.46 and 36.0), JL-501 (V₁) (32.46 and 33.0), and TAG-24 (V₃) (32.24 and 33.0) during the year 2017 and 2018 respectively. Sowing at 26th MW sowing window (S₂) recorded maximum days to maturity (115.66 and 117) in variety JL-776 (V₄). This was followed by variety RHRG-6083 (V₂) (113.66 and 114.67), JL-501 (V₁) (108.66 and 109.33), and TAG-24 (V₃) (105.66 and 106.33) during 2017 and 2018 respectively.

3.5 Yield Studies (Table 5)

3.5.1 Pod yield (kg ha⁻¹)

The mean pod yield of groundnut was 24.32 and 25.71 q ha⁻¹ was recorded during the year 2017 and 2018, respectively (Table 5).

3.5.1.1 Effect of varieties (Table 5A)

The pod yield was significantly higher in JL-776 (V_4) (26.59 and 28.14 $q\ ha^{-1}$) which was significantly superior rest of the groundnut varieties. This was followed by RHRG-6083 (V_2) (25.75 and 27.13 $q\ ha^{-1}$), JL-501 (V_1) (22.75 and 24.08 $q\ ha^{-1}$). The variety TAG-24 recorded significantly lower pod yield (22.19 and 23.49 $q\ ha^{-1}$) during the year 2017 and 2018, respectively. The differences in pod yield of groundnut varieties might be due to inherent genetical potential of variety.

3.5.1.2 Effect of sowing windows (Table 5B)

The pod yield of groundnut was influenced significantly due to extended sowing windows. The pod yield was maximum at 26th MW (S_2) sowing window (27.25 and 28.84 $q\ ha^{-1}$) and was at with 27th MW (25.89 and 27.27 $q\ ha^{-1}$). This was followed by 25th MW sowing window (23.35 and 24.72 $q\ ha^{-1}$) and 28th MW sowing window (20.79 and 22.01 $q\ ha^{-1}$) during the year 2017 and 2018, respectively. A sowing window of 26th MW was favourable to maximum pod production because of favourable weather condition. The longer growing season in early sowing time resulted in a better pod yield performance. These results confirm the finding reported by [5], [9] and [10].

3.5.1.3 Effects of interaction (Table 5C)

The pod yield ($q\ ha^{-1}$) was significantly influenced by interaction between varieties and sowing windows during the year 2017 and 2018. Sowing at 26th MW sowing window (S_2) recorded maximum pod yield (32.49 and 30.70 $q\ ha^{-1}$) in variety JL-776 (V_4). This was followed by variety RHRG-6083 (V_2) (29.96 and 28.30 $q\ ha^{-1}$), JL-501 (V_1) (26.86 and 25.38 $q\ ha^{-1}$), and TAG-24 (V_3) (26.04 and 24.60 $q\ ha^{-1}$) during the year 2017 and 2018, respectively. These results showed that delay in sowing of groundnut varieties could not able to assimilate the more biomass resulted in reduced pod yield of groundnut. (Table 5C)

3.5.2 Haulm yield ($kg\ ha^{-1}$)

The mean haulm yield of groundnut was 36.24 and 33.49 $q\ ha^{-1}$ during the year 2017 and 2018, respectively (Table 5).

3.5.2.1 Effect of varieties (Table 5A)

The haulm yield of groundnut was influenced significantly due to groundnut varieties. The haulm yield was significantly higher in JL-776 (V_4) (39.61 and 36.7 $q\ ha^{-1}$) and significantly superior rest of the groundnut varieties. This was followed by RHRG-6083 (V_2) (38.37 and 35.23 $q\ ha^{-1}$), JL-501 (V_1) (33.9 and 31.41 $q\ ha^{-1}$). The variety TAG-24 recorded significantly lower haulm yield (33.07 and 30.64 $q\ ha^{-1}$) during the year 2017 and 2018, respectively. The differences in haulm yield of groundnut varieties might be due to inherent genetical potential of groundnut variety (Table 5A).

3.5.2.2 Effect of sowing windows (Table 5B)

The haulm yield of groundnut was influenced significantly due to extended sowing windows. The haulm yield was maximum at 26th MW sowing window (40.60 and 37.61 $q\ ha^{-1}$), this was followed by 27th MW (38.57 and 35.41 $q\ ha^{-1}$) were at par with 25th MW sowing window. This was followed by 25th MW sowing window (34.80 and 32.24 $q\ ha^{-1}$) and 28th MW sowing window (30.98 and 28.71 $q\ ha^{-1}$) during the year 2017 and 2018, respectively. A sowing window of 26th MW was favourable to high haulm production because of favourable weather condition. The results are similar to those reported by [11]. (Table 5B)

3.5.2.3 Effects of interaction (Table 5C)

The haulm yield ($q\ ha^{-1}$) was significantly influenced by interaction between varieties and sowing windows during the year 2017 and 2018. Sowing at 26th MW sowing window (S_2) recorded maximum haulm yield (42.38 and 45.74 $q\ ha^{-1}$) in variety JL-776 (V_4). This was followed by variety RHRG-6083 (V_2) (39.08 and 42.17 $q\ ha^{-1}$), JL-501 (V_1) (35.04 and 37.82 $q\ ha^{-1}$) and TAG-24 (V_3) (33.97 and 36.66 $q\ ha^{-1}$) during the year 2017 and 2018, respectively. These results showed that delay in sowing of groundnut varieties could not able to assimilate the more biomass resulted in reduced haulm yield of groundnut (Table 5C).

Table 1. Plant height (cm) of groundnut (*Arachis hypogaea*) as influenced periodically by different treatments

| Treatment | 28 DAS | | | 42 DAS | | | 56 DAS | | | 70 DAS | | | 84 DAS | | | At harvest | | |
|--------------------------------------|-----------------------|----------------------|----------------------|-----------------------|------------------------|------------------------|---------------------|----------------------|---------------------|----------------------|---------------------|---------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled |
| A) Main plot: Varieties | | | | | | | | | | | | | | | | | | |
| V ₁ : JL-501 | 4.31 ^{bc} | 3.88 ^{bc} | 4.09 ^{bc} | 10.14 ^b | 9.25 ^{bc} | 9.70 ^{bc} | 20.34 ^c | 18.83 ^c | 19.59 ^c | 24.31 ^c | 23.69 ^c | 24.00 ^c | 29.42 ^b | 27.69 ^b | 28.55 ^b | 29.90 ^b | 27.99 ^b | 28.94 ^b |
| V ₂ : RHRG-6083 | 4.37 ^b | 3.94 ^{ab} | 4.16 ^b | 10.36 ^b | 9.47 ^{ab} | 9.92 ^{ab} | 21.50 ^b | 20.05 ^b | 20.78 ^b | 25.47 ^b | 24.85 ^b | 25.16 ^b | 33.75 ^a | 32.02 ^a | 32.89 ^a | 34.19 ^a | 32.34 ^a | 33.26 ^a |
| V ₃ : TAG-24 | 4.25 ^c | 3.82 ^c | 4.03 ^c | 10.03 ^b | 9.14 ^c | 9.59 ^c | 19.76 ^d | 18.34 ^d | 19.05 ^d | 23.78 ^d | 23.16 ^d | 23.47 ^d | 28.67 ^b | 26.94 ^b | 27.80 ^b | 29.15 ^b | 27.26 ^b | 28.21 ^b |
| V ₄ : JL-776 | 4.46 ^a | 4.01 ^a | 4.24 ^a | 10.83 ^a | 10.11 ^a | 10.47 ^a | 22.05 ^a | 20.59 ^a | 21.32 ^a | 26.12 ^a | 25.44 ^a | 25.78 ^a | 34.83 ^a | 33.00 ^a | 33.92 ^a | 35.27 ^a | 33.34 ^a | 34.30 ^a |
| S. E.± | 0.02 | 0.02 | 0.02 | 0.11 | 0.17 | 0.11 | 0.12 | 0.09 | 0.07 | 0.14 | 0.13 | 0.09 | 0.61 | 0.59 | 0.42 | 0.63 | 0.65 | 0.45 |
| C. D. at 5 % | 0.08 | 0.08 | 0.05 | 0.38 | 0.6 | 0.32 | 0.40 | 0.31 | 0.23 | 0.47 | 0.44 | 0.29 | 2.12 | 2.03 | 1.31 | 2.18 | 2.24 | 1.39 |
| B) Sub plot: Sowing windows | | | | | | | | | | | | | | | | | | |
| S ₁ : 25 th MW | 4.50 ^a | 4.08 ^a | 4.29 ^a | 11.08 ^a | 10.21 ^a | 10.65 ^a | 21.14 ^b | 19.67 ^b | 20.40 ^b | 25.60 ^b | 24.98 ^b | 25.29 ^b | 32.42 ^b | 30.69 ^b | 31.55 ^b | 32.90 ^b | 30.94 ^b | 31.92 ^b |
| S ₂ : 26 th MW | 4.39 ^b | 3.95 ^b | 4.17 ^b | 10.68 ^a | 9.81 ^a | 10.25 ^a | 21.86 ^a | 20.39 ^a | 21.12 ^a | 26.39 ^a | 25.83 ^a | 26.11 ^a | 33.67 ^a | 31.94 ^a | 32.80 ^a | 34.15 ^a | 32.27 ^a | 33.21 ^a |
| S ₃ : 27 th MW | 4.29 ^c | 3.86 ^c | 4.08 ^c | 10.04 ^b | 9.17 ^b | 9.60 ^b | 20.54 ^c | 19.07 ^c | 19.81 ^c | 24.54 ^c | 23.92 ^c | 24.23 ^c | 31.08 ^c | 29.35 ^c | 30.22 ^c | 31.52 ^c | 29.64 ^c | 30.58 ^c |
| S ₄ : 28 th MW | 4.20 ^d | 3.75 ^d | 3.98 ^d | 9.56 ^c | 8.79 ^c | 9.18 ^c | 20.12 ^d | 18.69 ^d | 19.40 ^d | 23.15 ^d | 22.42 ^d | 22.79 ^d | 29.50 ^d | 27.66 ^d | 28.58 ^d | 29.94 ^d | 28.07 ^d | 29.00 ^d |
| S. E.± | 0.02 | 0.02 | 0.01 | 0.14 | 0.09 | 0.08 | 0.07 | 0.06 | 0.05 | 0.08 | 0.09 | 0.06 | 0.23 | 0.24 | 0.17 | 0.22 | 0.22 | 0.16 |
| C. D. at 5 % | 0.05 | 0.06 | 0.04 | 0.40 | 0.27 | 0.23 | 0.2 | 0.18 | 0.13 | 0.25 | 0.26 | 0.17 | 0.68 | 0.7 | 0.47 | 0.65 | 0.64 | 0.44 |
| C) Interaction (AxB) | | | | | | | | | | | | | | | | | | |
| S ₁ V ₁ | 3.93 ^{cde} | 4.36 ^{def} | 4.14 ^{cdef} | 9.52 ^{cde} | 10.39 ^{cde} | 9.96 ^{cdefg} | 18.92 ^e | 20.39 ^f | 19.65 ^f | 24.52 ^{ef} | 25.14 ^{ef} | 24.83 ^{ef} | 27.94 ^{hij} | 29.67 ^{hi} | 28.80 ^{hi} | 28.22 ^{gh} | 30.15 ^{hi} | 29.18 ^{hi} |
| S ₂ V ₁ | 3.98 ^{bcd} | 4.41 ^{cde} | 4.20 ^{cde} | 9.69 ^{bcdde} | 10.56 ^{bcdde} | 10.13 ^{cde} | 20.19 ^c | 21.66 ^{cd} | 20.93 ^c | 25.19 ^{cd} | 25.81 ^{cd} | 25.50 ^{cd} | 29.60 ^{fg} | 31.33 ^g | 30.47 ^g | 29.98 ^{ef} | 31.81 ^{fg} | 30.90 ^g |
| S ₃ V ₁ | 3.87 ^{defg} | 4.30 ^{efgh} | 4.09 ^{efgh} | 9.04 ^{def} | 9.91 ^{def} | 9.48 ^{efghi} | 18.32 ^g | 19.79 ^{gh} | 19.06 ^{gh} | 23.38 ^h | 24.00 ^{hi} | 23.69 ^{ij} | 26.94 ^{jk} | 28.67 ^{jl} | 27.80 ^{jl} | 27.22 ^{hi} | 29.15 ^{ij} | 28.18 ^{kl} |
| S ₄ V ₁ | 3.72 ^f | 4.15 ^f | 3.94 ^f | 8.75 ^{ef} | 9.71 ^{ef} | 9.23 ^{efghi} | 17.91 ^h | 19.53 ^{ghi} | 18.72 ^{hi} | 21.66 ⁱ | 22.28 ^j | 21.97 ^k | 26.27 ^{kl} | 28.00 ^{kl} | 27.14 ^k | 26.55 ^{ij} | 28.48 ^{kl} | 27.52 ^{kl} |
| S ₁ V ₂ | 4.01 ^{bc} | 4.44 ^{bcd} | 4.22 ^{bcd} | 9.82 ^{bcd} | 10.69 ^{bcd} | 10.26 ^{bcd} | 20.33 ^c | 21.80 ^c | 21.07 ^c | 25.46 ^c | 26.08 ^c | 25.77 ^c | 32.94 ^{bcd} | 34.67 ^{bcd} | 33.80 ^{bcd} | 33.22 ^{bcd} | 35.15 ^{bc} | 34.18 ^{bc} |
| S ₂ V ₂ | 4.10 ^b | 4.53 ^b | 4.32 ^b | 10.00 ^{bc} | 10.87 ^{bc} | 10.44 ^{bc} | 20.85 ^b | 22.32 ^b | 21.58 ^b | 26.45 ^b | 27.07 ^b | 26.76 ^b | 33.27 ^{bc} | 35.00 ^{bc} | 34.14 ^{bc} | 33.65 ^{bc} | 35.48 ^{bc} | 34.56 ^{bc} |
| S ₃ V ₂ | 3.87 ^{defg} | 4.30 ^{efgh} | 4.09 ^{efgh} | 9.20 ^{cde} | 10.07 ^{cde} | 9.64 ^{defghi} | 19.79 ^d | 21.26 ^{de} | 20.53 ^{cd} | 24.44 ^{efg} | 25.06 ^{ef} | 24.75 ^{ef} | 31.60 ^{de} | 33.33 ^{de} | 32.47 ^{de} | 31.93 ^d | 33.64 ^{de} | 32.78 ^{de} |
| S ₄ V ₂ | 3.78 ^{fghi} | 4.21 ^{hi} | 4.00 ^{hi} | 8.87 ^{def} | 9.79 ^{def} | 9.33 ^{fghi} | 19.24 ^e | 20.61 ^f | 19.93 ^f | 23.06 ^f | 23.68 ^f | 23.37 ^f | 30.27 ^{ef} | 32.00 ^{ef} | 31.14 ^{ef} | 30.55 ^e | 32.48 ^{ef} | 31.52 ^{ef} |
| S ₁ V ₃ | 3.85 ^{efgh} | 4.28 ^{fgh} | 4.07 ^{fgh} | 9.43 ^{cde} | 10.30 ^{cde} | 9.86 ^{cdefgh} | 18.46 ^f | 19.93 ^g | 19.20 ^g | 23.91 ^g | 24.53 ^g | 24.22 ^{gh} | 27.60 ^{ijk} | 29.33 ^{hij} | 28.47 ^{hij} | 27.88 ^{ghi} | 29.81 ^{hij} | 28.85 ^{hij} |
| S ₂ V ₃ | 3.91 ^{cdefg} | 4.34 ^{defg} | 4.13 ^{defg} | 9.57 ^{cde} | 10.44 ^{cde} | 10.01 ^{cdef} | 18.95 ^e | 20.42 ^f | 19.68 ^f | 24.17 ^g | 24.79 ^g | 24.48 ^g | 28.60 ^{ghi} | 30.33 ^{gh} | 29.47 ^{gh} | 28.91 ^{fg} | 30.84 ^{gh} | 29.88 ^{gh} |
| S ₃ V ₃ | 3.80 ^{efghi} | 4.23 ^{ghi} | 4.01 ^{ghi} | 8.86 ^{ef} | 9.73 ^{ef} | 9.30 ^{fghi} | 18.03 ^{gh} | 19.50 ^{hi} | 18.76 ^{hi} | 23.01 ^h | 23.63 ^h | 23.32 ^h | 26.60 ^{jk} | 28.33 ^{ij} | 27.47 ^{ij} | 26.88 ^{hi} | 28.81 ^{ij} | 27.85 ^{ij} |
| S ₄ V ₃ | 3.72 ^{hi} | 4.15 ⁱ | 3.94 ⁱ | 8.71 ^{ef} | 9.65 ^{ef} | 9.18 ^{hi} | 17.94 ^h | 19.21 ⁱ | 18.58 ⁱ | 21.57 ⁱ | 22.19 ^j | 21.88 ^k | 24.94 ⁱ | 26.67 ^k | 25.80 ^k | 25.35 ^j | 27.15 ^k | 26.25 ^k |
| S ₁ V ₄ | 4.02 ^{bc} | 4.48 ^{bc} | 4.25 ^{bc} | 10.48 ^b | 11.35 ^b | 10.92 ^b | 20.96 ^b | 22.43 ^b | 21.70 ^b | 26.02 ^b | 26.64 ^b | 26.33 ^b | 34.27 ^b | 36.00 ^b | 35.14 ^b | 34.43 ^b | 36.48 ^b | 35.46 ^b |
| S ₂ V ₄ | 4.32 ^a | 4.73 ^a | 4.53 ^a | 11.57 ^a | 12.44 ^a | 12.01 ^a | 21.57 ^a | 23.04 ^a | 22.31 ^a | 27.51 ^a | 27.91 ^a | 27.71 ^a | 36.27 ^a | 38.00 ^a | 37.14 ^a | 36.55 ^a | 38.48 ^a | 37.52 ^a |
| S ₃ V ₄ | 3.91 ^{cdef} | 4.34 ^{defg} | 4.13 ^{defg} | 9.57 ^{cde} | 10.44 ^{cde} | 10.01 ^{cdef} | 20.15 ^c | 21.62 ^{cd} | 20.89 ^{cd} | 24.87 ^{de} | 25.49 ^{de} | 25.18 ^{de} | 32.27 ^{cd} | 34.00 ^{cd} | 33.14 ^{cd} | 32.55 ^{cd} | 34.48 ^{cd} | 33.52 ^{cd} |
| S ₄ V ₄ | 3.78 ^{ghi} | 4.29 ^{ghi} | 4.04 ^{ghi} | 8.82 ^f | 9.10 ^f | 8.96 ^f | 19.67 ^d | 21.11 ^e | 20.39 ^e | 23.38 ^h | 24.46 ^{gh} | 23.92 ^{hi} | 29.18 ^{gh} | 31.33 ^g | 30.26 ^g | 29.81 ^{ef} | 31.64 ^{fg} | 30.72 ^g |
| S. E.± | 0.04 | 0.04 | 0.03 | 0.28 | 0.19 | 0.17 | 0.14 | 0.12 | 0.09 | 0.17 | 0.18 | 0.12 | 0.47 | 0.48 | 0.33 | 0.44 | 0.44 | 0.31 |
| C. D. at 5 % | 0.11 | 0.12 | 0.08 | 0.81 | 0.55 | 0.47 | 0.4 | 0.36 | 0.26 | 0.49 | 0.51 | 0.34 | 1.36 | 1.4 | 0.95 | 1.3 | 1.28 | 0.89 |
| GM | 4.35 | 3.91 | 4.13 | 10.34 | 9.5 | 9.92 | 20.91 | 19.45 | 20.18 | 24.92 | 24.29 | 24.6 | 31.67 | 29.91 | 30.79 | 32.13 | 30.23 | 31.18 |

Note: Observations with same superscript are at par and with different superscript are significantly different.

Table 2. Mean number of branches plant⁻¹ of groundnut (*Arachis hypogaea*) as influenced periodically by different treatments

| Treatment | 28 DAS | | | 42 DAS | | | 56 DAS | | | 70 DAS | | | 84 DAS | | | At harvest | | |
|--------------------------------------|----------------------|----------------------|----------------------|-----------------------|-----------------------|-----------------------|----------------------|----------------------|----------------------|---------------------|----------------------|---------------------|---------------------|----------------------|---------------------|----------------------|----------------------|----------------------|
| | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled |
| A) Main plot: Varieties | | | | | | | | | | | | | | | | | | |
| V ₁ : JL-501 | 3.53 ^{bc} | 3.22 ^{bc} | 3.38 ^{bc} | 4.91 ^c | 4.44 ^c | 4.67 ^c | 8.31 ^b | 7.25 ^b | 7.78 ^b | 9.53 ^c | 8.39 ^c | 8.96 ^c | 9.54 ^c | 9.08 ^c | 9.31 ^c | 9.62 ^c | 8.79 ^c | 9.20 ^c |
| V ₂ : RHRG-6083 | 3.85 ^{ab} | 3.54 ^{ab} | 3.69 ^{ab} | 5.56 ^b | 5.09 ^b | 5.32 ^b | 8.72 ^{ab} | 7.66 ^{ab} | 8.19 ^{ab} | 10.80 ^b | 9.66 ^b | 10.23 ^b | 10.80 ^b | 10.34 ^b | 10.57 ^b | 10.73 ^b | 9.89 ^b | 10.31 ^b |
| V ₃ : TAG-24 | 3.42 ^c | 3.11 ^c | 3.27 ^c | 4.69 ^c | 4.22 ^c | 4.46 ^c | 7.77 ^c | 6.71 ^c | 7.24 ^c | 9.36 ^c | 8.22 ^c | 8.79 ^c | 9.39 ^c | 8.93 ^c | 9.16 ^c | 9.39 ^c | 8.55 ^c | 8.97 ^c |
| V ₄ : JL-776 | 3.89 ^a | 3.68 ^a | 3.78 ^a | 6.58 ^a | 6.11 ^a | 6.34 ^a | 9.19 ^a | 7.96 ^a | 8.57 ^a | 11.14 ^a | 9.93 ^a | 10.53 ^a | 11.23 ^a | 10.64 ^a | 10.94 ^a | 11.34 ^a | 10.55 ^a | 10.94 ^a |
| S. E.± | 0.09 | 0.11 | 0.07 | 0.14 | 0.11 | 0.09 | 0.19 | 0.15 | 0.12 | 0.08 | 0.11 | 0.07 | 0.07 | 0.11 | 0.07 | 0.09 | 0.07 | 0.06 |
| C. D. at 5 % | 0.33 | 0.38 | 0.23 | 0.48 | 0.39 | 0.28 | 0.66 | 0.53 | 0.38 | 0.29 | 0.38 | 0.22 | 0.25 | 0.38 | 0.21 | 0.32 | 0.24 | 0.18 |
| B) Sub plot: Sowing windows | | | | | | | | | | | | | | | | | | |
| S ₁ : 25 th MW | 3.67 ^{ab} | 3.36 ^{ab} | 3.51 ^{ab} | 5.89 ^b | 5.33 ^b | 5.61 ^b | 8.79 ^b | 7.73 ^b | 8.26 ^b | 10.61 ^b | 9.47 ^b | 10.04 ^b | 10.61 ^b | 10.15 ^b | 10.38 ^b | 10.61 ^b | 9.77 ^b | 10.19 ^b |
| S ₂ : 26 th MW | 4.04 ^a | 3.73 ^a | 3.89 ^a | 6.44 ^a | 6.06 ^a | 6.25 ^a | 9.48 ^a | 8.28 ^a | 8.88 ^a | 11.13 ^a | 9.99 ^a | 10.56 ^a | 11.19 ^a | 10.73 ^a | 10.96 ^a | 11.34 ^a | 10.58 ^a | 10.96 ^a |
| S ₃ : 27 th MW | 3.67 ^{ab} | 3.36 ^{ab} | 3.51 ^{ab} | 5.17 ^c | 4.70 ^c | 4.94 ^c | 8.31 ^c | 7.21 ^c | 7.76 ^c | 9.93 ^c | 8.79 ^c | 9.36 ^c | 10.01 ^c | 9.55 ^c | 9.78 ^c | 10.01 ^c | 9.17 ^c | 9.59 ^c |
| S ₄ : 28 th MW | 3.32 ^b | 3.10 ^b | 3.21 ^b | 4.23 ^c | 3.76 ^d | 3.99 ^d | 7.41 ^d | 6.35 ^d | 6.88 ^d | 9.16 ^d | 7.95 ^d | 8.56 ^d | 9.16 ^d | 8.58 ^d | 8.87 ^d | 9.12 ^d | 8.25 ^d | 8.69 ^d |
| S. E.± | 0.14 | 0.13 | 0.09 | 0.17 | 0.19 | 0.13 | 0.14 | 0.11 | 0.09 | 0.11 | 0.12 | 0.08 | 0.13 | 0.15 | 0.1 | 0.14 | 0.13 | 0.09 |
| C. D. at 5 % | 0.4 | 0.38 | 0.27 | 0.48 | 0.57 | 0.36 | 0.4 | 0.33 | 0.25 | 0.33 | 0.34 | 0.23 | 0.37 | 0.43 | 0.27 | 0.39 | 0.37 | 0.26 |
| C) Interaction (AxB) | | | | | | | | | | | | | | | | | | |
| S ₁ V ₁ | 2.69 ^d | 3.00 ^{cd} | 2.85 ^d | 4.70 ^{defgh} | 5.17 ^{defgh} | 4.94 ^{defgh} | 7.38 ^{defg} | 8.44 ^{defg} | 7.91 ^{defg} | 8.46 ^{fgn} | 9.6 ^{fgn} | 9.03 ^{gh} | 9.14 ^{ode} | 9.60 ^{ode} | 9.37 ^{ode} | 8.76 ^{efgh} | 9.60 ^{efgh} | 9.18 ^{efgh} |
| S ₂ V ₁ | 3.72 ^{ab} | 4.03 ^{ab} | 3.88 ^{ab} | 5.21 ^{def} | 5.68 ^{def} | 5.45 ^{def} | 7.67 ^{def} | 8.73 ^{def} | 8.20 ^{def} | 9.16 ^{def} | 10.30 ^{def} | 9.73 ^{def} | 9.88 ^c | 10.34 ^{def} | 10.11 ^c | 9.75 ^{cd} | 10.59 ^{cd} | 10.17 ^{cd} |
| S ₃ V ₁ | 3.44 ^{abcd} | 3.75 ^{abcd} | 3.60 ^{abcd} | 4.05 ^{fgh} | 4.52 ^{fgh} | 4.29 ^{fgh} | 7.17 ^{fgh} | 8.23 ^{fgh} | 7.70 ^{fgh} | 8.26 ^{ghi} | 9.40 ^{ghi} | 8.83 ^{ghi} | 8.94 ^{de} | 9.40 ^{de} | 9.17 ^{de} | 8.56 ^{fghi} | 9.40 ^{fghi} | 8.98 ^{fghi} |
| S ₄ V ₁ | 3.04 ^{bcd} | 3.35 ^{bcd} | 3.20 ^{bcd} | 3.78 ^{ghi} | 4.25 ^{ghi} | 4.02 ^{ghi} | 6.77 ^{ghi} | 7.83 ^{ghd} | 7.30 ^{ghi} | 7.69 ^{ij} | 8.83 ^l | 8.26 ^l | 8.37 ^{ef} | 8.83 ^{ef} | 8.60 ^{ef} | 8.04 ^{hi} | 8.88 ^{hi} | 8.46 ^{hi} |
| S ₁ V ₂ | 3.67 ^{abc} | 3.98 ^{abc} | 3.83 ^{abc} | 5.63 ^{cde} | 6.10 ^{cde} | 5.87 ^{cde} | 7.95 ^{cd} | 9.01 ^{cd} | 8.48 ^{cd} | 10.07 ^{bc} | 11.21 ^{bc} | 10.64 ^{bc} | 10.75 ^b | 11.21 ^b | 10.98 ^b | 10.37 ^{bc} | 11.21 ^{bc} | 10.79 ^{bc} |
| S ₂ V ₂ | 3.84 ^{ab} | 4.15 ^{ab} | 4.00 ^{ab} | 6.95 ^{ab} | 7.42 ^{ab} | 7.19 ^{ab} | 8.72 ^b | 9.78 ^b | 9.25 ^b | 10.48 ^b | 11.62 ^b | 11.05 ^b | 11.18 ^b | 11.64 ^b | 11.41 ^b | 10.80 ^b | 11.64 ^b | 11.22 ^b |
| S ₃ V ₂ | 3.42 ^{abcd} | 3.73 ^{abcd} | 3.58 ^{abcd} | 5.03 ^{defg} | 5.50 ^{defg} | 5.27 ^{defg} | 7.43 ^{defg} | 8.49 ^{defg} | 7.96 ^{defg} | 9.28 ^{de} | 10.42 ^{de} | 9.85 ^{de} | 9.96 ^c | 10.42 ^c | 10.19 ^c | 9.58 ^{cde} | 10.42 ^{cde} | 10.00 ^{cde} |
| S ₄ V ₂ | 3.22 ^{abcd} | 3.53 ^{abcd} | 3.37 ^{abcd} | 2.73 ⁱ | 3.20 ⁱ | 2.97 ⁱ | 6.53 ^{hi} | 7.59 ^{hi} | 7.06 ^{hi} | 8.80 ^{efg} | 9.94 ^{efg} | 9.37 ^{efg} | 9.48 ^{cd} | 9.94 ^{cd} | 9.71 ^{cd} | 8.81 ^{efgh} | 9.65 ^{efgh} | 9.23 ^{efgh} |
| S ₁ V ₃ | 3.27 ^{abcd} | 3.58 ^{abcd} | 3.43 ^{abcd} | 4.46 ^{efgh} | 4.93 ^{efgh} | 4.7 ^{efgh} | 7.20 ^{efgh} | 8.26 ^{efgh} | 7.73 ^{efgh} | 8.70 ^{efg} | 9.84 ^{efg} | 9.27 ^{efg} | 9.38 ^{cd} | 9.84 ^{cd} | 9.61 ^{cd} | 9.00 ^{defg} | 9.84 ^{defg} | 9.42 ^{defg} |
| S ₂ V ₃ | 3.36 ^{abcd} | 3.67 ^{abcd} | 3.51 ^{abcd} | 4.86 ^{defgh} | 5.33 ^{defgh} | 5.10 ^{defgh} | 7.35 ^{defg} | 8.41 ^{defg} | 7.88 ^{defg} | 8.91 ^{efg} | 10.05 ^{efg} | 9.48 ^{efg} | 9.69 ^{cd} | 10.15 ^{cd} | 9.92 ^{cd} | 9.31 ^{def} | 10.15 ^{def} | 9.73 ^{def} |
| S ₃ V ₃ | 3.02 ^{bcd} | 3.33 ^{bcd} | 3.18 ^{bcd} | 3.86 ^{ghi} | 4.33 ^{ghi} | 4.10 ^{ghi} | 6.46 ^{ij} | 7.52 ^{ij} | 6.99 ^{ij} | 7.87 ^{hij} | 9.01 ^{hij} | 8.44 ^{hij} | 8.55 ^{ef} | 9.01 ^{ef} | 8.78 ^{ef} | 8.17 ^{ghi} | 9.01 ^{ghi} | 8.59 ^{ghi} |
| S ₄ V ₃ | 2.81 ^{cd} | 3.12 ^{cd} | 2.96 ^{cd} | 3.69 ^{hi} | 4.16 ^{hi} | 3.93 ^{hi} | 5.82 ^j | 6.88 ^j | 6.35 ^j | 7.42 ^j | 8.56 ^k | 7.99 ^k | 8.10 ^k | 8.56 ^k | 8.33 ^k | 7.72 ^k | 8.56 ^k | 8.14 ^k |
| S ₁ V ₄ | 3.80 ^{ab} | 4.11 ^{ab} | 3.95 ^{ab} | 6.53 ^{abc} | 7.33 ^{abc} | 6.93 ^{abc} | 8.38 ^l | 9.44 ^{bc} | 8.91 ^{bc} | 10.64 ^b | 11.78 ^b | 11.21 ^b | 11.32 ^b | 11.78 ^b | 11.55 ^b | 10.94 ^b | 11.78 ^b | 11.36 ^b |
| S ₂ V ₄ | 4.02 ^a | 4.33 ^a | 4.17 ^a | 7.20 ^a | 7.33 ^a | 7.27 ^a | 9.38 ^a | 11.00 ^a | 10.19 ^a | 11.42 ^a | 12.56 ^a | 11.99 ^a | 12.15 ^a | 12.61 ^a | 12.38 ^a | 12.45 ^a | 12.98 ^a | 12.71 ^a |
| S ₃ V ₄ | 3.55 ^{abcd} | 3.86 ^{abcd} | 3.71 ^{abcd} | 5.86 ^{bcd} | 6.33 ^{bcd} | 6.10 ^{bcd} | 7.79 ^a | 8.99 ^{cde} | 8.39 ^{cde} | 9.75 ^{cd} | 10.89 ^{cd} | 10.32 ^{cd} | 10.74 ^b | 11.20 ^b | 10.97 ^b | 10.36 ^{bc} | 11.20 ^{bc} | 10.78 ^{bc} |
| S ₄ V ₄ | 3.34 ^{abcd} | 3.27 ^e | 3.31 ^{abcd} | 4.83 ^{defgh} | 5.30 ^{defgh} | 5.07 ^{defgh} | 6.27 ^{ij} | 7.33 ^{ij} | 6.8 ^{ij} | 7.91 ^{hij} | 9.33 ^{ghi} | 8.62 ^{ghi} | 8.36 ^{ef} | 9.33 ^{def} | 8.85 ^{def} | 8.44 ^{fghi} | 9.4 ^{fghi} | 8.92 ^{fghi} |
| S. E.± | 0.28 | 0.26 | 0.19 | 0.33 | 0.39 | 0.25 | 0.28 | 0.23 | 0.18 | 0.22 | 0.24 | 0.16 | 0.25 | 0.3 | 0.19 | 0.27 | 0.26 | 0.19 |
| C. D. at 5 % | NS | NS | 0.54 | 0.97 | 1.13 | 0.72 | 0.81 | 0.67 | 0.5 | 0.65 | 0.69 | 0.46 | 0.73 | 0.86 | 0.55 | 0.79 | 0.75 | 0.53 |
| GM | 3.67 | 3.39 | 3.53 | 5.43 | 4.96 | 5.2 | 8.50 | 7.39 | 7.94 | 10.21 | 9.05 | 9.63 | 10.24 | 9.75 | 10 | 10.27 | 9.44 | 9.86 |

Note: Observations with same superscript are at par and with different superscript are significantly different.

Table 3. Mean dry matter plant⁻¹ (g) of groundnut (*Arachis hypogaea*) as influenced periodically by different treatments

| Treatment | 28 DAS | | | 42 DAS | | | 56 DAS | | | 70 DAS | | | 84 DAS | | | At harvest | | |
|--------------------------------------|------------------------|------------------------|------------------------|---------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|-----------------------|-----------------------|-----------------------|----------------------|----------------------|----------------------|
| | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled |
| A) Main plot: Varieties | | | | | | | | | | | | | | | | | | |
| V ₁ : JL-501 | 2.14 ^c | 2.09 ^c | 2.12 ^c | 5.62 ^{bc} | 4.94 ^{bc} | 5.28 ^{bc} | 8.48 ^{bc} | 7.55 ^{bc} | 8.01 ^{bc} | 12.44 ^{bc} | 11.32 ^{bc} | 11.88 ^{bc} | 26.57 ^{bc} | 25.2 ^{bc} | 25.88 ^{bc} | 29.75 ^{bc} | 27.78 ^{bc} | 28.77 ^{bc} |
| V ₂ : RHRG-6083 | 2.34 ^b | 2.29 ^b | 2.31 ^b | 6.05 ^b | 5.37 ^b | 5.71 ^b | 9.07 ^{ab} | 8.14 ^{ab} | 8.60 ^{ab} | 12.83 ^{ab} | 11.71 ^{ab} | 12.27 ^{ab} | 28.08 ^b | 26.71 ^b | 27.39 ^b | 31.08 ^b | 29.11 ^b | 30.10 ^b |
| V ₃ : TAG-24 | 2.11 ^c | 2.07 ^d | 2.09 ^c | 5.40 ^c | 4.72 ^c | 5.06 ^c | 7.95 ^c | 7.02 ^c | 7.49 ^c | 11.95 ^c | 10.83 ^c | 11.39 ^c | 25.31 ^c | 23.94 ^c | 24.63 ^c | 28.33 ^c | 26.36 ^c | 27.35 ^c |
| V ₄ : JL-776 | 2.56 ^a | 2.54 ^a | 2.55 ^a | 6.88 ^a | 6.12 ^a | 6.50 ^a | 9.40 ^a | 8.31 ^a | 8.86 ^a | 13.24 ^a | 11.95 ^a | 12.60 ^a | 31.90 ^a | 30.23 ^a | 31.06 ^a | 34.36 ^a | 32.81 ^a | 33.58 ^a |
| S. E.± | 0.05 | 0.07 | 0.04 | 0.17 | 0.21 | 0.13 | 0.18 | 0.21 | 0.14 | 0.19 | 0.15 | 0.12 | 0.51 | 0.65 | 0.42 | 0.82 | 0.68 | 0.54 |
| C. D. at 5 % | 0.18 | 0.23 | 0.13 | 0.58 | 0.72 | 0.41 | 0.61 | 0.72 | 0.42 | 0.66 | 0.53 | 0.38 | 1.77 | 2.26 | 1.28 | 2.84 | 2.37 | 1.65 |
| B) Sub plot: Sowing windows | | | | | | | | | | | | | | | | | | |
| S ₁ : 25 th MW | 2.30 ^b | 2.25 ^b | 2.28 ^b | 6.05 ^b | 5.37 ^b | 5.71 ^b | 9.69 ^a | 8.76 ^a | 9.22 ^a | 13.84 ^a | 12.72 ^a | 13.28 ^a | 27.98 ^b | 26.61 ^b | 27.29 ^b | 31.67 ^b | 29.70 ^b | 30.68 ^b |
| S ₂ : 26 th MW | 2.83 ^a | 2.79 ^a | 2.81 ^a | 7.27 ^a | 6.59 ^a | 6.93 ^a | 8.92 ^b | 7.99 ^b | 8.45 ^b | 13.10 ^b | 11.98 ^b | 12.54 ^b | 31.10 ^a | 29.73 ^a | 30.42 ^a | 34.75 ^a | 32.78 ^a | 33.77 ^a |
| S ₃ : 27 th MW | 2.16 ^b | 2.12 ^b | 2.14 ^b | 5.60 ^c | 4.92 ^c | 5.26 ^c | 7.39 ^c | 7.86 ^c | 8.32 ^c | 12.33 ^c | 11.21 ^c | 11.77 ^c | 26.99 ^{bc} | 25.62 ^{bc} | 26.3b ^c | 29.58 ^c | 27.61 ^c | 28.60 ^c |
| S ₄ : 28 th MW | 1.85 ^c | 1.83 ^c | 1.84 ^c | 5.03 ^d | 4.27 ^d | 4.65 ^d | 7.98 ^d | 6.88 ^d | 7.43 ^d | 11.19 ^d | 9.90 ^d | 10.55 ^d | 25.79 ^c | 24.12 ^c | 24.96 ^c | 27.53 ^d | 25.97 ^c | 26.75 ^c |
| S. E.± | 0.07 | 0.06 | 0.05 | 0.11 | 0.13 | 0.08 | 0.14 | 0.16 | 0.11 | 0.15 | 0.17 | 0.11 | 0.56 | 0.52 | 0.38 | 0.69 | 0.68 | 0.48 |
| C. D. at 5 % | 0.21 | 0.19 | 0.14 | 0.31 | 0.37 | 0.24 | 0.42 | 0.46 | 0.3 | 0.42 | 0.49 | 0.31 | 1.65 | 1.53 | 1.09 | 2.03 | 1.98 | 1.38 |
| C) Interaction (AxB) | | | | | | | | | | | | | | | | | | |
| S ₁ V ₁ | 2.20 ^{cdefg} | 2.25 ^{cdefg} | 2.22 ^{cdefg} | 5.07 ^{def} | 5.75 ^{def} | 5.41 ^{def} | 7.67 ^{cde} | 8.60 ^{cde} | 8.14 ^{cde} | 11.70 ^{cde} | 12.82 ^{cde} | 12.26 ^{cde} | 25.43 ^{cdef} | 26.8 ^{cdef} | 26.12 ^{cdef} | 28.36 ^{bcd} | 30.33 ^{bcd} | 29.35 ^{bcd} |
| S ₂ V ₁ | 2.48 ^{bc} | 2.53 ^{bc} | 2.50 ^{bc} | 5.92 ^c | 6.60 ^c | 6.26 ^c | 8.19 ^{bcd} | 9.12 ^{bcd} | 8.66 ^{bcd} | 12.27 ^{bcd} | 13.39 ^{bcd} | 12.83 ^{bcd} | 27.20 ^{bcd} | 28.57 ^{bcd} | 27.88 ^{bcd} | 31.03 ^{bc} | 33.00 ^b | 32.02 ^{bc} |
| S ₃ V ₁ | 2.00 ^{defgh} | 2.05 ^{defgh} | 2.02 ^{defgh} | 4.71 ^{gh} | 5.39 ^{gh} | 5.05 ^{gh} | 7.19 ^{defg} | 8.12 ^{defg} | 7.66 ^{defg} | 10.96 ^{ef} | 12.08 ^{ef} | 11.52 ^{ef} | 24.48 ^{bcd} | 25.85 ^{def} | 25.17 ^{def} | 26.70 ^{cde} | 28.67 ^{cde} | 27.68 ^{cde} |
| S ₄ V ₁ | 1.70 ^h | 1.74 ^h | 1.72 ^h | 4.04 ^{hi} | 4.72 ^{hi} | 4.38 ^{hi} | 7.13 ^{efg} | 8.06 ^{efg} | 7.59 ^{efg} | 10.35 ^g | 11.47 ^{fg} | 10.91 ^{fg} | 23.68 ^{ef} | 25.05 ^{ef} | 24.37 ^{ef} | 25.03 ^{de} | 27.00 ^{de} | 26.02 ^{de} |
| S ₁ V ₂ | 2.28 ^{bcd} | 2.33 ^{bcd} | 2.31 ^{bcd} | 5.68 ^{cd} | 6.36 ^{cd} | 6.02 ^{cd} | 8.45 ^{bc} | 9.38 ^{bc} | 8.92 ^{bc} | 12.18 ^{bcd} | 13.30 ^{bcd} | 12.74 ^{bcd} | 27.44 ^{bcd} | 28.81 ^{bcd} | 28.12 ^{bcd} | 30.03 ^{bc} | 32.00 ^{bc} | 31.02 ^{bc} |
| S ₂ V ₂ | 2.71 ^b | 2.75 ^b | 2.73 ^b | 6.56 ^b | 7.24 ^b | 6.90 ^b | 8.79 ^b | 9.72 ^b | 9.26 ^b | 12.82 ^b | 13.94 ^b | 13.38 ^b | 28.52 ^{bc} | 29.89 ^{bc} | 29.20 ^{bc} | 31.36 ^b | 33.33 ^{bc} | 32.35 ^b |
| S ₃ V ₂ | 2.22 ^{cdef} | 2.27 ^{cdef} | 2.25 ^{cdef} | 5.02 ^{def} | 5.70 ^{def} | 5.36 ^{def} | 7.81 ^{bcd} | 8.74 ^{bcd} | 8.28 ^{bcd} | 11.36 ^{def} | 12.48 ^{def} | 11.92 ^{def} | 26.55 ^{bcd} | 27.92 ^{bcd} | 27.23 ^{bcd} | 28.36 ^{bcd} | 30.33 ^{bcd} | 29.35 ^{bcd} |
| S ₄ V ₂ | 1.94 ^{efgh} | 1.99 ^{efgh} | 1.96 ^{efgh} | 4.22 ^{ghi} | 4.90 ^{ghi} | 4.56 ^{ghi} | 7.49 ^{cdef} | 8.42 ^{cdef} | 7.96 ^{cdef} | 10.46 ^{fg} | 11.58 ^{fg} | 11.02 ^{fg} | 24.32 ^{def} | 25.69 ^{def} | 25.01 ^{def} | 26.70 ^{cde} | 28.67 ^{cde} | 27.68 ^{cde} |
| S ₁ V ₃ | 2.10 ^{cdefgh} | 2.15 ^{cdefgh} | 2.12 ^{cdefgh} | 4.87 ^{efg} | 5.55 ^{efg} | 5.21 ^{efg} | 7.34 ^{defg} | 8.27 ^{defg} | 7.81 ^{defg} | 11.23 ^{def} | 12.35 ^{def} | 11.79 ^{def} | 24.15 ^{def} | 25.52 ^{def} | 24.84 ^{def} | 27.36 ^{bcd} | 29.33 ^{bcd} | 28.35 ^{bcd} |
| S ₂ V ₃ | 2.45 ^{bcd} | 2.49 ^{bcd} | 2.47 ^{bcd} | 5.56 ^{cde} | 6.24 ^{cde} | 5.90 ^{cde} | 7.50 ^{cdef} | 8.43 ^{cdef} | 7.97 ^{cdef} | 11.83 ^{bcd} | 12.95 ^{bcd} | 12.39 ^{bcd} | 25.42 ^{cdef} | 26.79 ^{cdef} | 26.11 ^{cdef} | 29.03 ^{bcd} | 31.00 ^{bcd} | 30.02 ^{bcd} |
| S ₃ V ₃ | 1.90 ^{efgh} | 1.94 ^{efgh} | 1.92 ^{efgh} | 4.54 ^{ghi} | 5.22 ^{ghi} | 4.88 ^{ghi} | 6.78 ^{efg} | 7.71 ^{efg} | 7.25 ^{efg} | 10.47 ^{fg} | 11.59 ^{fg} | 11.03 ^{fg} | 23.14 ^f | 24.51 ^f | 23.83 ^f | 25.36 ^{de} | 27.33 ^{de} | 26.35 ^{de} |
| S ₄ V ₃ | 1.82 ^{gh} | 1.86 ^{gh} | 1.84 ^{gh} | 3.92 ⁱ | 4.60 ⁱ | 4.26 ⁱ | 6.47 ^{fg} | 7.40 ^{fg} | 6.94 ^{fg} | 9.81 ^{gh} | 10.93 ^{gh} | 10.37 ^{gh} | 23.06 ^f | 24.43 ^f | 23.74 ^f | 23.70 ^e | 25.67 ^e | 24.68 ^e |
| S ₁ V ₄ | 2.43 ^{bcd} | 2.48 ^{bcd} | 2.46 ^{bcd} | 5.87 ^c | 6.55 ^c | 6.21 ^c | 8.48 ^{bc} | 9.41 ^{bc} | 8.95 ^{bc} | 12.80 ^{bc} | 13.92 ^{bc} | 13.36 ^{bc} | 29.40 ^b | 30.77 ^b | 30.09 ^b | 31.70 ^b | 33.67 ^b | 32.68 ^b |
| S ₂ V ₄ | 3.52 ^a | 3.56 ^a | 3.54 ^a | 8.30 ^a | 8.98 ^a | 8.64 ^a | 10.54 ^a | 11.47 ^a | 11.01 ^a | 13.95 ^a | 15.07 ^a | 14.51 ^a | 37.80 ^a | 39.17 ^a | 38.48 ^a | 41.03 ^a | 43.00 ^a | 42.02 ^a |
| S ₃ V ₄ | 2.34 ^{bcd} | 2.39 ^{bcd} | 2.37 ^{bcd} | 5.43 ^{cde} | 6.11 ^{cde} | 5.77 ^{cde} | 7.77 ^{bcd} | 8.70 ^{bcd} | 8.24 ^{bcd} | 12.06 ^{bcd} | 13.18 ^{bcd} | 12.62 ^{bcd} | 28.29 ^{bc} | 29.66 ^{bc} | 28.98 ^{bc} | 30.03 ^{bc} | 32.00 ^{bc} | 31.02 ^{bc} |
| S ₄ V ₄ | 1.87 ^{gh} | 1.79 ^{gh} | 1.83 ^{gh} | 4.89 ^{efg} | 5.90 ^{cdef} | 5.39 ^{cdef} | 6.44 ^g | 8.03 ^{efg} | 7.24 ^{efg} | 9.00 ^h | 10.78 ^h | 9.89 ^h | 25.42 ^{cdef} | 28.00 ^{bcd} | 26.71 ^{bcd} | 28.46 ^{bcd} | 28.77 ^{bcd} | 28.62 ^{bcd} |
| S. E.± | 0.14 | 0.13 | 0.10 | 0.22 | 0.26 | 0.17 | 0.29 | 0.32 | 0.21 | 0.29 | 0.34 | 0.22 | 1.13 | 1.05 | 0.77 | 1.39 | 1.36 | 0.97 |
| C. D. at 5 % | 0.41 | 0.38 | 0.27 | 0.63 | 0.75 | 0.47 | 0.85 | 0.92 | 0.60 | 0.85 | 0.99 | 0.63 | 3.3 | 3.06 | 2.18 | 4.05 | 3.96 | 2.75 |
| GM | 2.29 | 2.25 | 2.27 | 5.99 | 5.29 | 5.64 | 8.73 | 7.75 | 8.24 | 12.61 | 11.45 | 12.03 | 27.96 | 26.52 | 27.24 | 30.88 | 29.02 | 29.95 |

Note: Observations with same superscript are at par and with different superscript are significantly different

Table 4. Mean number of days to 50 % flowering and maturity as influenced by different treatments in groundnut (*Arachis hypogaea*)

| Treatment | Days to 50 % flowering | | | Days to maturity | | |
|--------------------------------------|------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled |
| A) Main plot: Varieties | | | | | | |
| V ₁ : JL-501 | 31.17 ^c | 30.63 ^c | 30.90 ^c | 106.58 ^c | 105.91 ^c | 106.25 ^c |
| V ₂ : RHRG-6083 | 33.92 ^b | 33.38 ^b | 33.65 ^b | 111.42 ^b | 110.66 ^b | 111.04 ^b |
| V ₃ : TAG-24 | 30.83 ^c | 30.29 ^c | 30.56 ^c | 102.75 ^d | 102.41 ^d | 102.58 ^d |
| V ₄ : JL-776 | 36.00 ^a | 35.63 ^a | 35.81 ^a | 113.08 ^a | 111.75 ^a | 112.42 ^a |
| S. E.± | 0.13 | 0.19 | 0.12 | 0.25 | 0.28 | 0.21 |
| C. D. at 5 % | 0.45 | 0.66 | 0.36 | 0.85 | 0.96 | 0.65 |
| B) Sub plot: Sowing windows | | | | | | |
| S ₁ : 25 th MW | 33.67 ^b | 33.13 ^b | 33.40 ^b | 109.83 ^b | 109.16 ^b | 109.50 ^b |
| S ₂ : 26 th MW | 35.08 ^a | 34.54 ^a | 34.81 ^a | 111.83 ^a | 110.91 ^a | 111.37 ^a |
| S ₃ : 27 th MW | 32.67 ^c | 32.13 ^c | 32.40 ^c | 107.58 ^c | 106.64 ^c | 107.11 ^c |
| S ₄ : 28 th MW | 30.50 ^d | 30.13 ^d | 30.31 ^d | 104.58 ^d | 104.03 ^d | 104.30 ^d |
| S. E.± | 0.28 | 0.25 | 0.19 | 0.21 | 0.19 | 0.13 |
| C. D. at 5 % | 0.82 | 0.74 | 0.54 | 0.6 | 0.55 | 0.36 |
| C) Interaction (AxB) | | | | | | |
| S ₁ V ₁ | 31.13 ^g | 31.67 ^g | 31.40 ^g | 107.66 ^g | 108.33 ^g | 108.00 ^l |
| S ₂ V ₁ | 32.46 ^{ef} | 33.00 ^{ef} | 32.73 ^{ef} | 108.66 ^e | 109.33 ^e | 109.00 ^e |
| S ₃ V ₁ | 30.13 ^{gh} | 30.67 ^{gh} | 30.40 ^{gh} | 104.66 ^f | 105.33 ⁱ | 105.00 ^f |
| S ₄ V ₁ | 28.79 ^{hi} | 29.33 ^{hi} | 29.06 ^{hi} | 102.66 ^k | 103.33 ^k | 103.00 ^k |
| S ₁ V ₂ | 34.13 ^{cd} | 34.67 ^{cd} | 34.40 ^{cd} | 111.66 ^c | 112.33 ^c | 112.00 ^c |
| S ₂ V ₂ | 35.46 ^{bc} | 36.00 ^{bc} | 35.73 ^{bc} | 113.66 ^b | 114.67 ^b | 114.17 ^b |
| S ₃ V ₂ | 33.13 ^{de} | 33.67 ^{de} | 33.40 ^{de} | 109.66 ^d | 110.33 ^d | 110.00 ^d |
| S ₄ V ₂ | 30.79 ^g | 31.33 ^g | 31.06 ^g | 107.66 ^g | 108.33 ^g | 108.00 ^g |
| S ₁ V ₃ | 31.13 ^g | 31.67 ^g | 31.40 ^g | 103.66 ^j | 104.33 ^j | 104.00 ^l |
| S ₂ V ₃ | 32.24 ^{ef} | 33.00 ^{ef} | 32.73 ^{ef} | 105.66 ^h | 106.33 ^h | 106.00 ^h |
| S ₃ V ₃ | 30.13 ^{gh} | 30.67 ^{gh} | 30.40 ^{gh} | 101.66 ⁱ | 102.33 ⁱ | 102.00 ^l |
| S ₄ V ₃ | 27.46 ^j | 28.00 ^j | 27.73 ^j | 98.66 ^m | 98.00 ^m | 98.33 ^m |
| S ₁ V ₄ | 36.13 ^b | 36.67 ^b | 36.40 ^b | 113.66 ^b | 114.33 ^b | 114.00 ^b |
| S ₂ V ₄ | 37.79 ^a | 38.33 ^a | 38.06 ^a | 115.66 ^a | 117.00 ^a | 116.33 ^a |
| S ₃ V ₄ | 35.13 ^{bc} | 35.67 ^{bc} | 35.40 ^{bc} | 110.55 ^d | 112.33 ^c | 111.44 ^c |
| S ₄ V ₄ | 33.46 ^{de} | 33.33 ^{de} | 33.40 ^{de} | 107.11 ^f | 108.67 ^f | 107.89 ^f |
| S. E.± | 0.57 | 0.51 | 0.38 | 0.41 | 0.38 | 0.26 |
| C. D. at 5 % | NS | NS | 1.08 | 1.21 | 1.10 | 0.73 |
| General Mean | 32.98 | 32.48 | 32.73 | 108.46 | 107.68 | 108.07 |

Note: Observations with same superscript are at par and with different superscript are significantly different

Table 5. Mean pod and haulm yield (q ha⁻¹) of *kharif* groundnut (*Arachis hypogaea*) as influenced by different treatments

| Treatment | Pod yield (q ha ⁻¹) | | | Haulm yield (q ha ⁻¹) | | |
|--------------------------------------|---------------------------------|--------------------|--------------------|-----------------------------------|--------------------|--------------------|
| | 2017 | 2018 | Pooled | 2017 | 2018 | Pooled |
| A) Main plot: Varieties | | | | | | |
| V ₁ : JL-501 | 22.75 ^c | 24.08 ^c | 23.42 ^c | 33.90 ^c | 31.41 ^c | 32.65 ^c |
| V ₂ : RHRG-6083 | 25.75 ^d | 27.13 ^b | 26.44 ^b | 38.37 ^b | 35.23 ^b | 36.80 ^b |
| V ₃ : TAG-24 | 22.19 ^d | 23.49 ^d | 22.84 ^d | 33.07 ^d | 30.64 ^d | 31.85 ^d |
| V ₄ : JL-776 | 26.59 ^a | 28.14 ^a | 27.36 ^a | 39.61 ^a | 36.70 ^a | 38.16 ^a |
| S. E.± | 0.4 | 0.43 | 0.29 | 0.6 | 0.55 | 0.42 |
| C. D. at 5 % | 1.39 | 1.47 | 0.91 | 2.08 | 1.92 | 1.29 |
| B) Sub plot: Sowing windows | | | | | | |
| S ₁ : 25 th MW | 23.35 ^c | 24.72 ^c | 24.03 ^c | 34.80 ^c | 32.24 ^c | 33.52 ^c |
| S ₂ : 26 th MW | 27.25 ^a | 28.84 ^a | 28.04 ^a | 40.60 ^a | 37.61 ^a | 39.11 ^a |
| S ₃ : 27 th MW | 25.89 ^b | 27.27 ^b | 26.58 ^b | 38.57 ^b | 35.41 ^b | 36.99 ^b |
| S ₄ : 28 th MW | 20.79 ^d | 22.01 ^d | 21.40 ^d | 30.98 ^d | 28.71 ^d | 29.85 ^d |
| S. E.± | 0.22 | 0.23 | 0.16 | 0.33 | 0.3 | 0.21 |
| C. D. at 5 % | 0.64 | 0.68 | 0.45 | 0.95 | 0.88 | 0.61 |
| C) Interaction (AxB) | | | | | | |
| S ₁ V ₁ | 23.26 ⁱ | 21.98 ^l | 22.62 ⁱ | 30.34 ^l | 32.75 ^l | 31.55 ^l |
| S ₂ V ₁ | 26.86 ^e | 25.38 ^e | 26.12 ^e | 35.04 ^e | 37.82 ^e | 36.43 ^e |
| S ₃ V ₁ | 25.22 ^g | 23.83 ^g | 24.53 ^g | 32.90 ^g | 35.51 ^g | 34.20 ^g |
| S ₄ V ₁ | 20.97 ^k | 19.81 ^k | 20.39 ^k | 27.35 ^k | 29.52 ^k | 28.44 ^k |
| S ₁ V ₂ | 26.24 ^f | 24.79 ^f | 25.51 ^f | 34.22 ^f | 36.94 ^f | 35.58 ^f |
| S ₂ V ₂ | 29.96 ^c | 28.30 ^c | 29.13 ^c | 39.08 ^c | 42.17 ^c | 40.62 ^c |
| S ₃ V ₂ | 28.40 ^d | 27.31 ^d | 27.86 ^d | 36.41 ^d | 40.69 ^d | 38.55 ^d |
| S ₄ V ₂ | 23.91 ^h | 22.59 ^h | 23.25 ^h | 31.19 ^h | 33.67 ^h | 32.43 ^h |
| S ₁ V ₃ | 22.47 ^j | 21.23 ^j | 21.85 ^j | 29.31 ^j | 31.64 ^j | 30.48 ^j |
| S ₂ V ₃ | 26.04 ^f | 24.60 ^f | 25.32 ^f | 33.97 ^f | 36.66 ^f | 35.31 ^f |
| S ₃ V ₃ | 24.94 ^g | 23.57 ^g | 24.26 ^g | 32.54 ^g | 35.12 ^g | 33.83 ^g |
| S ₄ V ₃ | 20.51 ^k | 19.37 ^k | 19.94 ^k | 26.75 ^k | 28.87 ^k | 27.81 ^k |
| S ₁ V ₄ | 26.89 ^e | 25.41 ^e | 26.15 ^e | 35.08 ^e | 37.86 ^e | 36.47 ^e |
| S ₂ V ₄ | 32.49 ^a | 30.70 ^a | 31.59 ^a | 42.38 ^a | 45.74 ^a | 44.06 ^a |
| S ₃ V ₄ | 30.52 ^b | 28.84 ^b | 29.68 ^b | 39.81 ^b | 42.97 ^b | 41.39 ^b |
| S ₄ V ₄ | 22.65 ^j | 21.40 ^j | 22.02 ^j | 29.54 ^j | 31.88 ^j | 30.71 ^j |
| S. E.± | 0.44 | 0.46 | 0.32 | 0.65 | 0.61 | 0.43 |
| C. D. at 5% | 1.28 | 1.36 | 0.90 | 1.91 | 1.77 | 1.21 |
| General Mean | 24.32 | 25.71 | 25.01 | 36.24 | 33.49 | 34.87 |

Note: Observations with same superscript are at par and with different superscript are significantly different

4. CONCLUSIONS

Amongst all the groundnut (*Arachis hypogaea*) varieties, JL-776 (*Phule Bharati*) variety found significantly superior under extended sowing windows followed by varieties RHRG-6083, TAG-24 and JL-501. A sowing window of 26th MW was favourable to maximum pod production because of favourable weather condition. Sowing during 26th MW was observed to be most suitable and optimum for groundnut considering the growth and yield attributes. This sowing window was at par with 27th MW sowing window.

ACKNOWLEDGEMENT

Author thankful to Department of Agricultural Meteorology, Division of Plant Pathology, Division of Agril. Entomology, College of Agriculture, Pune and Mahatma Phule Krishi Vidyapeeth, Rahuri.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nigam SN, Rao RCN and Wynne JC. Effects of temperature and photoperiod on vegetative and reproductive growth of groundnut (*Arachis hypogaea* L.). *Journal of Agronomy and Crop Science*. 1998; 181(2):117-124.
2. Murthy, PSS and Rao RCN. Physiological basis of variations in pod yield of rainfed groundnut (*Arachis hypogaea* L.) under different dates of sowing. *Indian Journal of Agronomy*. 1986;31:106-108.
3. Gosh TK and Das Gupta DK. Effects of date of sowing on growth and yield of groundnut in lateritic upland. *Plant Science*. 1975;7:56-60.
4. Enikuomehin OA, Olowe VIO, Alao OS and Atayese MO. Assessment of Cercospora leaf spot disease of sesame in different planting dates in South Western Nigeria. *Moor Journal of Agricultural Research*. 2002;3(1):76-82.
5. Frimpong A. Characterization of groundnut (*Arachis hypogaea* L.) in Northern Ghana. *Pakistan Journal of Biological Sciences*. 2004;7:838-842.
6. Gopalkrishnan S, Varisai MS and Verma L. Studies on time of sowing on the growth and yield of *Arachis hypogaea* L. *Agricultural Journal*. 1967;54(6):283-287.
7. Smartt J. Factors influencing yield and quality of groundnuts in northern Rhodesia. *Emp. Journal of Expo Agriculture*. 1964;32: 343-351.
8. Gritton ET and Ebert RD. Interaction of planting date and powdery mildew on pea plant performance. *American Society for Horticultural Science*. 1975;100(2): 137-142.
9. Banik NC, Nath R and Chakraborty PK. Effect of dates of sowing on growth and yield of groundnut crop. *Journal of Crop and Weed*. 2009;5(2):59-62.
10. Bala HMB, Ogunlela VB, Tanimu B, Kuchinda NC. Response of two groundnut (*Arachis hypogaea* L.) varieties to sowing date and NPK fertilizer rate in a semiarid environment: Growth and growth attributes. *Asian Journal of Crop Science*. 2011;3:141-150.
11. Ntare BR, Williams JH and Nadunguru BJ. Effects of seasonal variation in temperature and cultivar on yield and yield determination of irrigated groundnut (*Arachis hypogaea* L.) during the dry season in the Sahel of West Africa. *The Journal of Agricultural Science*. 1998; 131(4):439-448.

© 2023 Wankhede et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/99837>