



First Record of the *Glugea* sp. on the *Alosa fallax* (Lacepède, 1803) Caught in the Marine Waters of Syria (Eastern Mediterranean)

**Reem Al-Sheikh Rasheed^{a++*}, Mohammad Galiya^{a#}
and Zouhair Almajid^{a#}**

^a Department of Zoology, College of Sciences, Tishreen University, Lattakia, Syria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v25i1653

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105360>

Original Research Article

Received: 18/06/2023

Accepted: 23/08/2023

Published: 01/09/2023

ABSTRACT

During the implementation of the current research entitled "Study of diet, growth, reproduction, and determination of the concentrations of some trace elements in *Alosa fallax* (Lacépède, 1803) in the marine waters of the Lattakian coast", we found the parasite *Glugea* sp. inside one of the *Alosa fallax* caught on 7/24/2023 AD from the Ras Al-Baseet area (35.8524° N, 35.8378° E) using Purse nets. Infection with the parasite *Glugea* sp. In colonies of yellowish-white masses over the digestive system and muscles. The anatomical and histological characteristics were consistent with scientific references, and this is the first record of the parasite *Glugea* sp. in the Syrian marine waters and the eastern Mediterranean.

⁺⁺ Ph.D. Student;

[#] Professor;

*Corresponding author: Email: mmrom92rm@gmail.com;

Keywords: *Glugea*; *Alosa fallax*; *microsporidea*; Syria.

1. INTRODUCTION

Clupeidae occupies the first place in economic fisheries in the world, as the annual catch rate of its species reached 11 million tons (more than 20% of the total catch of fish in the world), of which more than a thousand tons are caught annually in the Mediterranean Sea [1,2].

Alosa fallax is characterized by a moderately deep body (the maximum body height is medium), short notched jaw, colour: bright blue back, silver to white fins, with a dark spot behind the gill cover, and (5-6) other spots to follow [3].

Pelagic fish that migrate long distances. enter estuaries. It feeds on zooplankton and small fish. reproduction season in the spring. They spawn in tidal places and estuaries. Lay eggs on the bottom. It is spread in the Mediterranean from Finland and Iceland to Morocco and the countries of the eastern Mediterranean [4,5,6,7, 8].

Glugea sp. belongs to the class Microsporidea and they are protozoan parasites, it is believed to be closely related to the fungus Mecoza. These parasites are transmitted between hosts by monocytes cells bacteria and may be transmitted by ingestion by fish directly without the presence of an intermediary. Microsporidea can be found in almost any environment and can infect large numbers of vertebrate and invertebrate species [9]. They range in length from less than 1 μ m to more than 30 μ m [10].

Juvenile hosts are more likely to cause mortality than adults [7]. It causes the enlargement of host cells that look like tumors and are full of bacteria, and this is common in fish infected with the genus *Glugea* [5,6,7,8].

The immune response is often weak or non-existent, as the parasites are protected within host cells or encapsulated in xenomas. However,

parasites can attack the host's immune system during primary infection or if the host cell or tumor ruptures [11].

Microsporidea infection may result in altered host-sex ratios by killing or transforming males into females (sex inversion) of the host as well as changes in behavior and alterations in feeding [12].

2. MATERIALS AND METHODS

Alosa fallax was collected with an average of 1,100 individuals from the Ras Al-Baseet area on the coast of Lattakia Governorate, Syria, eastern Mediterranean. (35.8524° N, 35.8378° E) from 10/19/2021 to 7/24/2023 using a Purse net to carry out a biological study (nutrition, reproduction, growth, and determination of heavy metal concentrations in the muscles of the *Alosa fallax*) and during the anatomy of the sample collected on 24/7/2023 we noticed the presence of xenomas located on the internal organs of the fish (stomach and intestines). A microscopic swab was taken from one of the tumors (xenomas), and the presence of germs was noted inside the tumors.

The sample was kept in 4% formaldehyde for a day to stabilize it, then a series of routine procedures were performed to treat it with paraffin wax and to obtain a permanent slide containing the tissue sections of this parasitic infection according to the reference [13].

3. RESULTS AND DISCUSSIONS

Alosa fallax was collected from the Syrian marine waters using Purse nets during the period from 10/19/2021 to 7/24/2023, and its total number reached 1,100 individuals. On 7/24/2023, one of the individuals was found infected with a parasitic disease (Fig. 1), as it was found during the anatomy of the sample. The presence of fatty



Fig. 1. *Alosa fallax* caught from Ras al-Bassit on 24/7/2023, with a total length of 21 cm and a weight of 59.83 gr

tumors in the viscera (stomach and intestines) was recorded, as well as some tumors within the muscles (Fig. 2).

A smear was taken from one of the tumors and examined microscopically using a light microscope, and it was found that there were germs within the tumors (Fig. 3). Then a part of the digestive system was taken and a histological study was conducted and it was found that there are injuries and infections within the tissue (Fig. 4).

Upon research and reference comparison, it was found that the parasite belongs to the *Microsporidea* class specifically of the genus *Glugea* sp., There is no scientific article yet on the infection of *Alosa fallax* fishes with the parasite *Glugea* sp. in the eastern basin of the Mediterranean. We will mention some of the parasites of the genus *Glugea*, which have been recorded infecting different fish species in the following Table 1.



Fig. 2. Fatty tumors within the bowels of *Alosa fallax* Caught from Ras al-Baset on 7/24/2023




Fig. 3. Microscopic smears at 40 magnification of sebaceous tumors taken from parasite-infected fish showing the spores inside the tumors




Fig. 4. Histological section of fatty tumor on the guts of *Alosa fallax* fish infected with the parasite *Glugea* sp

Table 1. Some parasites of the genus *Glugea* that have been recorded to be infecting different fish species

Parasite	The host	References
<i>Glugea hertwigi</i>	Rainbow smelt <i>Osmerus mordax</i> ,	Canada [14].
<i>Glugea gasterostei</i>	Three-spined stickleback <i>Gasterosteus aculeatus</i> (Actinopterygii: Gasterosteiformes)	[15].
<i>Glugea nagelia</i>	<i>Cephalopholis hemistiktos</i> (Actinopterygii: Serranidae)	Red Sea [16].
<i>Glugea jazanensis</i>	<i>Lutjanus bohar</i>	Red Sea [17].
<i>Glugea arabica</i>	<i>Epinephelus polyphekadion</i>	Red Sea [18].
<i>Glugea serranus</i>	Blacktail comber, <i>Serranus atricauda</i> (Teleostei: Serranidae)	Madeira Archipelago (Portugal) [19].
<i>Glugea sardinellensis</i>	<i>Sardinella aurita</i>	Tunisian coasts. [20].
<i>Glugea eda</i>	Striated fusilier, <i>Caesio striata</i> ,	Red Sea off Saudi Arabia [21].
<i>Glugea sardinellensis</i>	<i>Sardinella aurita</i>	Algeria [22].
<i>Glugea thunni</i>	Blue fins tunna <i>Thunnus thynnus</i>	Atlantic Ocean [23].
<i>Glugea</i> sp.	<i>Alosa fallax</i>	Syrian marine waters Current research,2023

4. CONCLUSIONS

The results showed that one of the individuals of *Alosa fallax* analyzed was infected with the parasite *Glugea* sp.

In terms of fatty tumors full of bacteria, this is the first record of parasitic infection with this parasite among marine fish in the Syrian and eastern Mediterranean waters.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Syskan vi. Food-Fishes of Atlantic ocean, Dictionnry, Moscow, Agricultural Publishing (in Russian).1988;360.
- FAO. SOFI2018-the state of food security and Nutrition in the word; 2018.
- Whithead PIP; Bauchot NL, Hureau JC; Nilsoen J. Tortonese E. Fishes of the North-eastern Atlantic and Mediterranean.Unesco, Paris. 1986;II.
- Golani D. Öztürk B. Başusta N. Fishes of the Eastern Mediterranean–Turkish marine research foundation, Istanbul, Turkey. 2006;24.
- Sindermann CJ. 'Principal diseases of marine fish and shellfish'. (Academic Press: San Diego);1990.
- Lom J, Dyková I. Protozoan parasites of fishes. Elsevier; 1992.
- Shaw RW, Kent ML. Fish microsporidia. In 'The Microsporidia and Microsporidiosis'. (Eds M Wittner, LM Weiss.) (ASM Press: Washington DC). 1999;447–501
- Lom J, Nilsen F. Fish microsporidia: Fine Structural Diversity; 2003.
- Kent ML, Shaw RW, Sanders JL. Microsporidia in fish. In: Weiss LM, Becnel JJ (eds) Microsporidia: pathogens of opportunity. Wiley, Hoboken; 2014.
- Larsson JIR. Identification of microsporidia. Acta Protozool. 1999;38:161–197.
- Rohde K. (Ed.). *Marine parasitology*. Csiro Publishing. 2005;590.
- Dunn AM, Terry RS, Smith JE. Transovarial transmission in the Microsporidia. Advances in Parasitology. 2001;48:57–100.
- Carleton HM, Drury RA, Willington EA, Cameron H. cited from Carleton, Histological techniques, 4th Ed, Oxford Univ. Press, N. 4, Toronto; 1962.
- Lovy J, Kostka M, Dykova I, Arsenault G, Peckova H, Wright GM, Speare DJ. Phylogeny and morphology of *Glugea hertwigi* from rainbow smelt *Osmerus mordax* found in Prince Edward Island, Canada. Diseases of Aquatic Organisms. 2009;86(3):235-243.
- Tokarev YS, Voronin VN, Senderskiy IV, The Microsporidium *Glugea gasterostei* Voronin 1974 (Microsporidia: Marinosporidia) from the three-spined stickleback *Gasterosteus aculeatus* (Actinopterygii: Gasterosteiformes) as an independent species. Parazitologiya Issi IV .2015; 49(2):81–92
- Abdel-Baki AAS, Tamih AF, Al-Qahtani HA, Al-Quraishy S, Mansour L, *Glugea jazanensis* sp. Nov. infecting *Lutjanus bohar* in the Red Sea: Ultra- structure and phylogeny. Dis Aquat Organ. 2015b;116(3):185–190Amsterdam; New York.and phylogeny. Int. J. Parasitol. 33:107–127.
- Abdel-Baki AAS, Al-Quraishy S, Rocha S, Dkhil MA, Casal G, Azevedo C. Ultrastructure and phylogeny of *Glugea nagelia* n. sp. (Micro- sporidia: Glugeidae), infecting the intestinal wall of the yellowfin hind, *Cephalopholis hemistiktos* (Actinopterygii: Serranidae), from the Red Sea. Folia Parasitologica. 2015a;62(1):1–7
- Azevedo C, Abdel-Baki AAS, Rocha S, Al-Quraishy S, Casal G. Ultrastruc- ture and phylogeny of *Glugea arabica* n. sp. (Microsporidia), infecting the marine fish *Epinephelus polyphemadion* from the Red Sea. Eur J Protistol. 2016;52(228):11–21
- Casal G, Rocha S, Costa G, Al-Quraishy S, Azevedo C . Ultrastructural and molecular characterization of *Glugea serranus* n. sp, a microsporidian infecting the blacktail comber, *Serranus atricauda* (Teleostei: Serranidae), in the Madeira Archipelago (Portugal). Parasitol Res. 2016;115(10):3963–3972
- Mansour L, Thabet A, Harrath AH, Al Omar S, Mukhtar A, Sayed SR, Abdel-Baki AAS. New microsporidia, *Glugea sardinellensis* n. sp. (Microsporea, Glugeida) found in *Sardinella aurita* Valenciennes, 1847, collected of Tunisian coasts. Acta Protozool. 2016;55(4):281–290.
- Mansour, Lamjed JY, Zhang Heba M, Abdel-Haleem, Ahmed B. Darwish Saleh Al-Quraishy, and Abdel-Azeem S. Abdel-Baki. Ultrastructural description and phylogeny of a novel microsporidian,

- Glugea eda n. sp. from the striated fusilier, *Caesio striata*, in the Red Sea of Saudi Arabia. *Acta Tropica*. 2020;204:105331.
22. Ramdani S, Ramdane Z, Slamovits CH, Trilles JPG lugea sp. infecting *Sardinella aurita* in Algeria. *Journal of Parasitic Diseases*. 2022;46(3):672-685.
23. López-Verdejo A, Montero FE, de la Gándara, F., Gallego MA, Ortega A, Raga JA, Palacios-Abella JFA. Severe microsporidian disease in cultured Atlantic Bluefin Tuna (*Thunnus thynnus*). *IMA Fungus*, Chicago. 2022;13(1):1-14.

© 2023 Rasheed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105360>