



# Influence of Levels of Water Soluble Fertilizer on Yield, Quality and Economics of Sugarcane Through Fertigation

Gagan B. Shetty <sup>a+++\*</sup>, Ananthakumar M. A. <sup>b#</sup>,  
Bhagyalakshmi T. <sup>c#</sup>, Yogananda S. B. <sup>d†</sup>,  
Avte Shubhangi Basveshwar <sup>e++</sup>, Varalakshmi V. <sup>f++</sup>,  
Vanitha T. <sup>a‡</sup> and Harsha B. R. <sup>g^</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, College of Agriculture, GKVK, Bangalore- 560065, India.

<sup>b</sup> (SS & AC), Water Technology Centre, ZARS, V.C. Farm, Mandya- 571405, India.

<sup>c</sup> Department of SS & AC, COA, V.C. Farm, Mandya- 571405, India.

<sup>d</sup> Department of Agronomy, COA, V.C. Farm, Mandya- 571405, India.

<sup>e</sup> Department of Soil Science and Agricultural Chemistry, Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani, Maharashtra-431402, India.

<sup>f</sup> Department of Soil Science and Agricultural Chemistry, College of Agriculture, VC Farm, Mandya- 571405, India.

<sup>g</sup> ICAR-KVK Siwan, Dr. Rajendra Prasad Central Agricultural University, Pusa Samastipur, Bihar-841408, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJPSS/2023/v35i213974

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106558>

<sup>++</sup> Ph. D Scholar;

<sup>#</sup> Assistant Professor;

<sup>†</sup> Associate Professor;

<sup>‡</sup> Senior Research Fellow;

<sup>^</sup> Scientist;

\*Corresponding author: E-mail: gaganbshetty1997@gmail.com;

## ABSTRACT

A field experiment was conducted at Zonal Agricultural Research Station, V. C. Farm, Mandya during 2018-19 to study the effect of water-soluble fertilizer levels (urea, MAP, MOP) under drip fertigation on yield, quality and economics of sugarcane. The investigation was carried out in Randomized Complete Block Design with seven treatments and replicated thrice. The treatments comprised of four levels of fertigation viz., 150, 125, 100 and 75 per cent Recommended Dosage of Fertilizers (RDF) through water soluble fertilizers, 100 per cent RDF through conventional fertilizers without FYM, 100 per cent RDF through conventional fertilizers with FYM and control. The results revealed that application of 125 per cent RDF through water soluble fertilizers recorded significantly higher yield and yield parameters viz., number of millable canes clump<sup>-1</sup> (14.33), cane length (346.33 cm), number of internodes cane<sup>-1</sup>(21.00), cane girth (13.80 cm), single cane weight (2.97 Kg) and cane yield (212 t ha<sup>-1</sup>) and was on par with 150 per cent RDF and 100 per cent RDF through water soluble fertilizer application. Least number of millable canes clump<sup>-1</sup>(8.67), cane length (220.00 cm), number of internodes cane<sup>-1</sup> (17.87), cane girth (10.2 cm), single cane weight (1.72 Kg) and cane yield (112.60 t ha<sup>-1</sup>) was recorded in control. Quality parameters like pol (18.0%), CCS production (12.44%), sugar yield (26.38 t ha<sup>-1</sup>) and juice extraction per centage (69.65 %) were observed significantly higher with 125 per cent RDF through drip fertigation and was on par with 150 per cent RDF and 100 per cent RDF through water soluble fertilizer application. Economic perspectives such as higher gross returns (Rs. 487600 ha<sup>-1</sup>), net returns (Rs. 368532 ha<sup>-1</sup>) and B: C ratio (4.10) was noticed with application of 125 per cent RDF and hence concluded that fertigation at 125 per cent RDF found optimum for higher yield and net returns.

**Keywords:** Yield; quality; economics; drip fertigation; sugarcane; RDF.

## 1. INTRODUCTION

Sugarcane (*Saccharum officinarum* L.) is a versatile crop that provides sugar, bio-fuel, fiber and manure besides many by products. The crop is grown mainly to manufacture sugar and for making *gur* and *khandasari*. It is one of the important commercial crops of sugar in the world. Globally it is cultivated over an area of 24.5 m. ha with a production of 1850 m.t and productivity of 75.5 t ha<sup>-1</sup> [1]. In India, sugarcane is grown under diverse agro-climatic situations covering an area of 5.35 m. ha producing 366 m. t of sugarcane with the productivity of 69.02 t ha<sup>-1</sup> with the distinction of being the second largest producer of sugar after Brazil, and the world's biggest consumer of the sweetener (22.5 m. t) [2]. Karnataka ranks third in area (5.0 lakh ha), fourth in production (47 m t) and second in productivity of 94 t ha<sup>-1</sup> [3].

Sugarcane being a long duration crop, produces huge amount of biomass and requires large quantity of water compared to other crops. Water requirement of sugarcane under conventional

method of cultivation varies from 2000 to 2500 mm depending upon soil type and climate. Vagaries of monsoon and declining ground-water resource due to over exploitation have resulted in shortage of fresh water supply for agricultural use. Further, India's water demand will nearly double by 2030 from the present 740 billion m<sup>3</sup> to 1.3 trillion m<sup>3</sup>, thus necessitating efficient water management for improving agricultural productivity.

In conventional method of irrigation and fertilizer application, there is considerable loss of water and leaching of mobile nutrients, particularly nitrogen which in turn leads to pollution of water bodies and deterioration of soil health. Providing optimum soil condition throughout the growing period of sugarcane is of paramount importance to realize higher yields. Therefore, drip fertigation, one of the promising potential technologies offers the great scope to increase the cane productivity up to 200-220 t ha<sup>-1</sup> by saving 40-50 per cent irrigation water with increase in nutrient efficiency by 40 per cent [4].

The combined application of water and fertilizers is ideal for proper crop growth, since irrigation water acts as a carrier for the nutrients that the crops require. Through irrigation system, soluble fertilizers are thus transported directly to the feeding zone by frequent application in small quantities. This provides valuable alternative opportunities for growing crops under conditions close to those of nutrient solution, when properly handled. Though fertigation related research in sugarcane is not limited, identification of location specific optimum levels of fertilizers in sugarcane for drip fertigation to enhance productivity and studying the impact of application of MAP (Mono Ammonium Phosphate) which is an acidic fertilizer requires prime attention. Keeping these facts in mind, the present investigation was taken to know the effect of water-soluble fertilizer levels under drip fertigation on yield, quality and economics of sugarcane.

## 2. MATERIALS AND METHODS

The experiment was conducted at Zonal Agricultural Research Station, V. C. Farm, Mandya, during 2018-19. The Soil of the experimental site was red sandy loam with medium organic carbon (0.56 per cent), low available nitrogen (273 kg ha<sup>-1</sup>), medium available phosphorus (28 kg ha<sup>-1</sup>) and available potassium (268 kg ha<sup>-1</sup>). The investigation was carried out in Randomized Complete Block Design with seven treatments and replicated thrice. The treatments comprised of four levels of fertigation viz., 150, 125, 100 and 75 per cent RDF through water soluble fertilizers (WSF), 100 per cent RDF through conventional fertilizers without FYM, 100 per cent RDF through conventional fertilizers with FYM and control (Table 1). The land was prepared by ploughing with tractor drawn disc plough followed by disc harrowing and passing cultivator twice to bring the soil to fine tilth.

Layout was prepared with gross plot size of 15.0 m × 10.0 m. Drip irrigation system (pump, filter units, main line and sub line) was installed. The laterals were placed at 1.95 m apart. The drip line was passed in between 30 cm apart paired row at 20 cm. Inline emitters were placed 40 cm apart with discharge rate of 4 lph. The recommended farm yard manure (15 t ha<sup>-1</sup>) was applied uniformly to all the treatments two week before planting of sets except for control plot. The conventional fertilizers were applied as per recommended dose of fertilizer (250: 100: 125 kg of NPK ha<sup>-1</sup>), 10 percent N, 100 percent of P & K

were applied as basal dose and remaining N was applied as top dressing at 45, 75, 105 days after planting with 20, 30 & 40 percent N. While for drip fertigation plots N, P and K were applied as per nutrient scheduling twice in a week. No irrigation and fertilizers were applied to control plot.

The healthy and viable two eye budded sets were collected from a well grown nine-month-old plant crop of sugarcane and planted in a zig-zag manner in paired rows with spacing of 30 cm between rows and 180 cm between pairs of rows (180-30-180 cm) using VCF-0517 variety. Sugarcane variety VCF-0517 was developed at Zonal Agriculture Research Station, V.C Farm, Mandya and unveiled by UAS-B at Krishi Mela in the year 2017. A new high yielding midlate maturing genotype VCF- 0517 was developed from general cross collection of Co-8371. High yielding (80-90 t/acre), high tillering, better quality, suitable for wide row planting, high jaggery yield and quality and good ratooning ability are general characteristics of this variety. It occupied 80 % of the area in Southern Karnataka.

Weed management was done through application of Metribuzin 70 per cent @ 600 g ha<sup>-1</sup> at two days after planting. Optimum plant population was maintained by filling the gaps at 30 days after planting. Hand weeding was done at 45, 90 and 135 days after planting (DAP) to keep plots weed free. Earthing up was carried out by tractor drawn implement at 105 days after planting. Wrapping and propping were done when the crop attained 10 months age to prevent lodging. Healthy crop stand was maintained by adopting need-based plant protection and recommended package of practices. During harvest five canes from each plot were cut randomly and juice was extracted. Juice samples were analysed for brix, pol and purity per cent content as per the standard procedure. Data recorded on various observations viz., yield and quality and soil parameters generated from treatments imposed were subjected to analysis of variance as per the procedures outlined by Rangaswamy [5]. The level of significance used in 'F' and 't' tests were at  $p \leq 0.05$  and critical difference values were calculated wherever the 'F' test was found to be significant.

### 2.1 Juice Quality Parameters

The juice samples extracted by means of a power-driven sugarcane crusher from five canes

**Table 1. Treatment details**

Treatment	Details
T <sub>1</sub>	Control
T <sub>2</sub>	RDF through conventional fertilizers +No FYM
T <sub>3</sub>	RDF through conventional fertilizers + FYM
T <sub>4</sub>	150% RDF through water soluble fertilizers + FYM
T <sub>5</sub>	125% RDF through water soluble fertilizers + FYM
T <sub>6</sub>	100% RDF through water soluble fertilizers + FYM
T <sub>7</sub>	75% RDF through water soluble fertilizers + FYM

**Note:** RDF: Recommended Dosage of Fertilizer- 250:100:125 kg N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O ha<sup>-1</sup>

FYM: Farm Yard Manure

Conventional fertilizers: Urea, SSP and MOP

Water soluble fertilizers: Urea, MAP and MOP

selected at randomly from the net plot area at harvest were analysed for the following quality parameters.\

$$\text{CCS (\%)} = \{S - (B - S) \times 0.4\} \times 0.73$$

Where,

S= Sucrose per cent in juice

B= Brix per cent in juice

### 2.1.1 Brix (%)

The brix readings of the filtered juice samples were recorded with the help of brix hydrometer standardized for 27.5 °C. The juice temperatures were recorded for necessary temperature corrections and expressed in per cent Brix values.

### 2.1.2 Sucrose or pol (%)

The juice samples were clarified as per Horne's dry lead sub acetate clarification method [6] and filtered through Whatman number 1 filter paper. The pol per cent readings of the filtrates were recorded with the help of polarimeter. The pol readings so recorded were correlated with observed degrees brix with the help of Schmitz table so as to get the values of pol per cent of juice which is synonymously used for sucrose per cent of juice.

### 2.1.3 Purity (%)

It is the ratio of pol per cent of juice to the corrected degrees of brix expressed in percentage and the values were computed as per the following formula.

$$\text{Purity coefficient (\%)} = \frac{\text{Pol per cent in juice}}{\text{Corrected brix \%}} \times 100$$

### 2.1.4 Commercial Cane Sugar (CCS %)

It is the amount of white commercial sugar that can be obtained from total cane juice after removing total soluble solids. The values of commercial cane sugar on per cent cane basis were computed from the following formula.

### 2.1.5 Sugar yield (t ha<sup>-1</sup>)

Sugar yield was calculated by using the following formula as suggested by Venkatachari and Sastry [7].

$$\text{Sugar yield (t ha}^{-1}\text{)} = \text{CCS (\%)} \times \text{cane yield (t ha}^{-1}\text{)} / 100$$

Where,

CCS = Commercial cane sugar (%)

### 2.1.6 Juice extraction (%)

$$\text{Juice extraction per cent (\%)} = \frac{\text{Juice weight}}{\text{Cane weight}} \times 100$$

## 3. RESULTS AND DISCUSSION

### 3.1 Yield and Yield Attributes of Sugarcane

Drip fertigation of 125 per cent RDF through water soluble fertilizers noticed significantly higher single cane weight (2.97 Kg), cane length (346.33 cm), cane girth (13.80 cm), number of millable canes clump<sup>-1</sup> (14.33), number of internodes cane<sup>-1</sup> (21.00) and cane yield (212 t ha<sup>-1</sup>) and was observed to be on par with application of 150 per cent RDF and 100 per cent RDF through water soluble fertilizers, respectively.

Cane yield differed significantly with the application of varied fertigation levels of WSF.

**Table 2. Yield and yield parameters of sugarcane as influenced by levels of water-soluble fertilizers through fertigation**

Treatments	Single Cane weight (kg)	Cane length (cm)	Cane girth (cm)	No. of millable canes clump <sup>-1</sup>	No. of internodes Cane <sup>-1</sup>	Cane yield (t ha <sup>-1</sup> )	Single Cane weight (kg)
T <sub>1</sub>	1.72	220.00	10.2	8.67	17.87	112.60	1.72
T <sub>2</sub>	2.05	260.00	11.4	9.80	18.21	171.70	2.05
T <sub>3</sub>	2.10	265.00	11.8	10.50	18.40	175.00	2.10
T <sub>4</sub>	2.63	337.67	13.00	14.00	20.80	205.00	2.63
T <sub>5</sub>	2.97	346.33	13.80	14.33	21.00	212.00	2.97
T <sub>6</sub>	2.56	336.00	12.30	13.33	20.50	200.00	2.56
T <sub>7</sub>	2.44	305.00	12.40	13.00	19.53	195.00	2.44
S.Em ±	0.10	8.41	0.35	0.34	0.39	5.07	0.10
CD ( $p \leq 0.05$ )	0.31	25.90	1.08	1.05	1.19	15.61	0.31

Note: T<sub>1</sub>: Control

T<sub>2</sub>: RDF through conventional fertilizer +No FYM


T<sub>3</sub>: RDF through conventional fertilizers + FYM

T<sub>4</sub>: 150% RDF through water soluble fertilizers + FYM

T<sub>5</sub>: 125% RDF through water soluble fertilizers + FYM

T<sub>6</sub>: 100% RDF through water soluble fertilizers + FYM

T<sub>7</sub>: 75% RDF through water soluble fertilizers + FYM

**Fig. 1. Cane yield (t ha<sup>-1</sup>) as influenced by levels of water-soluble fertilizers through fertigation**

Cane yield with the application of 125 per cent RDF through WSF was significantly higher (212.0 t ha<sup>-1</sup>) than recorded in control (112.60 t ha<sup>-1</sup>), soil application of 100 per cent RDF through conventional fertilizer with FYM (175.00 t ha<sup>-1</sup>) and without FYM (171.70 t ha<sup>-1</sup>) and with 75 per cent RDF through WSF (195.00 t ha<sup>-1</sup>) treatments and was on par with 150 RDF per cent (205.00 t ha<sup>-1</sup>) and 100 per cent RDF

(200.00 t ha<sup>-1</sup>) through WSF application. The conventional method of cane cultivation recorded the lowest cane yield (171.7 and 175 t ha<sup>-1</sup>, respectively) when compared to treatments with varied water soluble fertilizer levels under drip fertigation. This might be due to considerable wastage of plant nutrients to alternate drying and wetting with loss of nutrients through deep percolation below root zone and volatilization of

nitrogen resulting in imbalance in soil water metabolism and nutrient environment [8]. In addition, drip fertigation of WSF at 75 per cent RDF resulted in significantly higher cane yield (195.00 t ha<sup>-1</sup>) when compared with application RDF through conventional fertilization without FYM (171.70 t ha<sup>-1</sup>) and RDF through conventional fertilization with FYM (175.00 t ha<sup>-1</sup>), it clearly indicates that application of fertilizers through drip fertigation system can improve sugarcane growth and yield parameters with 25 per cent lower dose of fertilizers (Table 2, Fig. 1, respectively).

The increased yield and yield parameters in case of drip fertigation treatments is mainly due to better accessibility of essential nutrients and soil moisture at rhizosphere to encounter the crop requirements with optimum sunlight and better aeration leading to improved yield attributes. The dual row planting system recorded higher individual cane weight and cane yield and was observed to be on par with the similar system of planting at 100 per cent RDF and lower cane yield was noticed in conventional irrigation method. Such findings are comparable to those of [9] under drip fertigation at 125 per cent RDF through WSF. The increased individual cane length, girth and number of internodes lead to increased single cane weight and this advantageous influence was due to greater uptake of macro nutrients thereby improving the

cell activities which eventually resulted in higher cane weight [10]. Comparable results were also reported by Mahadkar et al. [11] Deshmukh and Katake [12] Subramani [13].

### 3.2 Juice Quality

Drip fertigation of 125% RDF through water soluble fertilizers noticed significantly higher Juice extraction per cent (69.65%), pol (%) (18.00 per cent), commercial cane sugar per cent (12.44 %), sugar yield (26.38 t ha<sup>-1</sup>) and was observed to be on par with application of 150% RDF and 100% RDF through water soluble fertilizers, respectively (Table 3, Fig. 2, respectively). The significant difference due to varied.

WSF levels on quality parameters like brix (%) and purity (%) of cane was not noticed. The Juice extraction per cent, brix, pol, purity, CCS per cent and sugar yield were increased with increase in fertilizer levels up to 125 per cent RDF then declined by further increased in fertilizer level; the results are conformity with [14].

The results revealed that, juice quality declined beyond the application of fertilizer 125 per cent RDF. The possible reason for this might be, with increased dose of nitrogen and increased activity of enzymes, which is responsible for

**Table 3. Sugarcane juice quality as influenced by levels of water-soluble fertilizers through fertigation**

Treatments	Juice extraction (%)	Brix (%)	Pol (%)	Purity (%)	CCS (%)	Sugar yield (t ha <sup>-1</sup> )	Juice extraction (%)
T <sub>1</sub>	53.15	19.07	15.28	80.40	10.05	11.33	53.15
T <sub>2</sub>	58.03	20.00	16.20	81.13	10.72	18.40	58.03
T <sub>3</sub>	59.98	19.87	16.13	81.32	10.69	18.68	59.98
T <sub>4</sub>	69.15	19.93	17.50	87.85	12.06	24.74	69.15
T <sub>5</sub>	69.65	20.40	18.00	88.25	12.44	26.38	69.65
T <sub>6</sub>	67.46	20.20	17.50	86.81	11.99	24.64	67.46
T <sub>7</sub>	62.24	19.60	16.87	86.12	11.51	22.44	62.24
S.Em ±	1.20	0.59	0.13	2.24	0.16	0.62	1.20
CD ( $p \leq 0.05$ )	3.71	NS	0.39	NS	0.51	1.92	3.71

Note: T<sub>1</sub>: Control

T<sub>2</sub>: RDF through conventional fertilizer +No FYM


T<sub>3</sub>: RDF through conventional fertilizers + FYM

T<sub>4</sub>: 150% RDF through water soluble fertilizers + FYM

T<sub>5</sub>: 125% RDF through water soluble fertilizers + FYM

T<sub>6</sub>: 100% RDF through water soluble fertilizers + FYM

T<sub>7</sub>: 75% RDF through water soluble fertilizers + FYM



**Fig. 2. Sugar yield (t ha<sup>-1</sup>) as influenced by levels of water-soluble fertilizers through fertigation**

**Note:**

T<sub>1</sub>: Control

T<sub>2</sub>: RDF through conventional fertilizer +No FYM

T<sub>3</sub>: RDF through conventional fertilizers + FYM

T<sub>4</sub>: 150% RDF through water soluble fertilizers + FYM

T<sub>5</sub>: 125% RDF through water soluble fertilizers + FYM

T<sub>6</sub>: 100% RDF through water soluble fertilizers + FYM

T<sub>7</sub>: 75% RDF through water soluble fertilizers + FYM

RDF-250:100:125 NPK kg ha<sup>-1</sup>

**Table 4. Economics of sugarcane as influenced by levels of water-soluble fertilizers through fertigation**

Treatments	Cost of cultivation (Rs ha <sup>-1</sup> )	Gross returns (Rs ha <sup>-1</sup> )	Net returns (Rs ha <sup>-1</sup> )	B:C ratio
T <sub>1</sub>	85350	258980	173630	3.03
T <sub>2</sub>	97924	394910	296985	4.03
T <sub>3</sub>	109924	402500	292575	3.66
T <sub>4</sub>	121211	471500	350288	3.89
T <sub>5</sub>	119067	487600	368532	4.10
T <sub>6</sub>	116924	460000	343075	3.93
T <sub>7</sub>	114780	448500	333719	3.90

degradation of sucrose and changing into glucose and fructose. This is in accordance with [15] but they reported the poor quality of juice beyond 300 kg N ha<sup>-1</sup>. The higher brix, sucrose (pol %) and CCS per cent, sugar yield, juice extraction per cent of sugarcane was obtained in the treatment which received fertigation with 125 per cent RDF than the other fertigation levels. Similar findings were reported by Singandhupe et al. [16].

### 3.3 Economics

Application of WSF at 150 per cent RDF noticed higher cost of cultivation (Rs. 121211 ha<sup>-1</sup>) when compared with rest of the treatments. However, application of 125 per cent RDF through WSF recorded higher gross returns (Rs. 487600 ha<sup>-1</sup>),

net returns (Rs. 368532 ha<sup>-1</sup>) and B: C ratio (4.10) as compared to control (Rs. 258980 ha<sup>-1</sup>, Rs. 173630 ha<sup>-1</sup>, 3.03, respectively), soil application of 100 per cent RDF through conventional fertilizer with FYM (GR-Rs. 402500, NR-Rs. 292575 ha<sup>-1</sup>, B:C ratio-3.66, respectively) and without FYM (GR-Rs. 394910, NR- Rs. 296985 ha<sup>-1</sup>, B:C ratio-4.03) and with 75 per cent RDF through WSF (Rs. 448500 ha<sup>-1</sup>, Rs. 333719 ha<sup>-1</sup>, 3.90) and was on par with 150 RDF per cent (Rs. 471500 ha<sup>-1</sup>, Rs. 350288.6 ha<sup>-1</sup>, 3.89) and 100 per cent RDF (Rs. 460000 ha<sup>-1</sup>, Rs. 343075 ha<sup>-1</sup>, 3.93) through WSF application (Table 4).

The higher gross return, net return and B:C ratio were mainly due to greater cane yield and sugar yield as reported by Guruswamy and Mahendran

[17] Veeraputhiran et al. [18] Gururaj [19] Pawar et al. [20]. Soil application of 100 per cent RDF registered lower gross return, net return and B:C ratio and was mainly due lower cane and sugar yield. This result is analogous with the conclusions of [21].

#### 4. CONCLUSION

The brix values, pol and purity percentage determine the quality of cane. The cane quality is good if it contains 12- 13 per cent of sucrose and purity with minimum amount of reducing sugar. In the present study, the quality parameters were affected due to excess application of fertilizer, i.e more than 125 per cent of RDF. Furthermore, higher gross returns (Rs. 487600 ha<sup>-1</sup>), net returns (Rs. 368532 ha<sup>-1</sup>) and B: C ratio (4.10) was noticed with application of 125 per cent RDF and hence concluded that drip fertigation is an innovative technology for maximizing the yield. Though the cost of drip fertigation unit was quite high, considering its longer life period, the benefit accrued out of it will be for longer period. Drip fertigation of 125 % RDF through WSF recorded significantly greater cane yield of 212 t ha<sup>-1</sup> and least cane yield was registered in control and hence concluded that fertigation at 125 per cent RDF found optimum for higher yield and quality of sugarcane.

#### ACKNOWLEDGEMENT

I would like to thanks to my teacher, major advisor and Chairperson of the Advisory Committee, Dr. Ananthakumar M. A, Assistant Professor, Water technology centre, ZARS, V.C. Farm, Mandya, University of Agricultural Sciences, Bangalore, for his valuable suggestions and encouragement throughout my work and to complete my research successfully.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Anonymous Agri. Situation in India, agristat; 2018. Available:<http://www.des.kar.nic.in>.
2. Anonymous Agri. Situation in India, agristat; 2019a. Available:<http://www.des.kar.nic.in>.
3. Anonymous Agri. Situation in India, agristat; 2019b. Available:<http://www.des.kar.nic.in>.
4. Sonawane DA, Sabale RN. Effect of different sources of organic nitrogen on growth, yield and quality of Suru sugarcane. J. Maharashtra Agric. Univ. 2000;25(1):15-17.
5. Rangaswamy RA. Text Book of Agricultural Statistics. New Age International Publishers, New Delhi. 2010;537.
6. Meade C. Cane Sugar Handbook (X Ed.), John Wiley and Sons, New York, p.788. Methods. W. Junk, N.V.N.V. Publication. The Hague. 1977;343-381.
7. Venkatachari A, Sastry SK. Nutrient requirements of sugarcane at Rudrur, Andhra Pradesh. Proc. 4<sup>th</sup> All India Conf. Sugarcane Researchers and Dev. Workers. 1960;146.
8. Ridge DR, Hewson SA. Drip irrigation management strategies. Proceedings of the Australian Society of Sugarcane Technologies, held at Bundaberg, Queensland, Australia. 1995;17: 8-15.
9. Mahendran SJ, Stephen Arul AC, Prabagar P, Rajarathinam AND Yeyasrinivas, R, Effect of paired row planting on growth, yield and economics of sugarcane under drip fertigation system. Sugarcane J. 2005;14:44- 52.
10. Jayaram SK, Thanunathan A, Jeyabal AND Thiruppathi M. Influence of fertigation on sugarcane yield, economics and post-harvest soil status under sandy loam soil condition. Plant Arc. 2010;10(2):773-775.
11. Mahadkar UV, Raut RS, Shinde JB, Pawar DD, Gaikwad CB., Response of pre-seasonal sugarcane to planting material, inter row spacing and fertilizer levels under drip irrigation, Proc. Nation. Sem. Relevance of Micro- irrigation in Sugarcane, 9-10<sup>th</sup> February, VSI Pune, Maharashtra, India. 2005;121-130.
12. Deshmukh AS, Katake SS. Techno-economic evaluation of micro irrigation in sugarcane In: Proceedings of International Conference on Plasticulture and Precision Farming, 17-21 November, New Delhi. 2014;218.
13. Subramani T. Optimization of nutrient requirement for hybrid chillies under drip fertigation system in open field cultivation. Ph.D. Thesis. Tamil Nadu Agric. Univ., Coimbatore; 2008.
14. More A, Meyer JH, Moodleys P, Berg, M., Drip irrigated sugarcane response to nitrogen applied through fertigation in late



- and early season cycles. South African Sugar Tech. Assoc. 2005;81:333-342.
15. Singh PN, Mohan SC, Water use and yield response of sugarcane under different irrigation scheduling and nitrogen levels in sub-tropical region. Agric. Water Manag. 1994;26:253-264.
  16. Singandhupe RB, Bankar MC, Anand PSB, Patil NG. Management of drip irrigated sugarcane in Western India. Archives of Agron & Soil Sci. 2008;4(6): 629- 649.
  17. Guruswamy A, Mahendran PP, Study on the influence of irrigation regimes and fertigation levels on sugarcane under subsurface drip fertigation system. Res. J.Agric.Sci. 2010;12(7):545-550.
  18. Veeraputhiran R, Balasubramanian R, Pandian BJ, Chelladurai M, Tamilselvi, R, Renganathan VG. Effect of subsurface drip fertigation on cane yield, water use efficiency and economics of Sugarcane. Madras Agric. J. 2012;99(4-6):255-258.
  19. Gururaj K. Optimization of water and nutrient requirement through drip fertigation in aerobic rice. M.Sc. (Agri.) Thesis, Univ. of Agril. Sci., Bangalore; 2013.
  20. Pawar DD, Dingre SK, Durgude AG, Enhancing nutrient use and sugarcane (*Saccharum officinarum*) productivity with reduced cost through drip fertigation in western Maharashtra. Indian J. Agric. Sci. 2014;84(7):844–849.
  21. Bhoi PG, Pawar DD, Raskar BS, Bangar AR. and Shinde SH. Effects of water-soluble fertilizers through drip on growth, yield and quality of suru sugarcane. Microirrigation (Eds. Singh, H. P., Kaushish, S. P., Ashwani Kumar, Murthy, T. S. and Jose C. Samuel). 2001; 520-525.

© 2023 Shetty et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/106558>