



Effect of Organic Manures and Natural Farming on Soil Properties and Nutrient Uptake by Carrot

Sumit Kumar ^a, Vijay Pal Singh Panghal ^{a++*}, Narender ^{b++}
and Shiv Kumar ^a

^a Department of Vegetable Science, CCS Haryana Agricultural University, Hisar, India.

^b Department of Soil Science, CCS Haryana Agricultural University, Hisar, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i214089

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109005>

Original Research Article

Received: 04/09/2023

Accepted: 10/11/2023

Published: 11/11/2023

ABSTRACT

The present investigation carried out at the Research Farm Area of Deen Dayal Upadhyaya Center of Excellence for Organic Farming, Chaudhary Charan Singh Haryana Agricultural University, Hisar during *Rabi season* 2022-23 with the objectives to evaluate the effect of organic manures and natural farming on growth, yield and quality of carrot as well as on microbial count and nutrient uptake and economics of carrot. The experiment consisting of 8 treatments *i.e.*, T₁- RDN through Farm Yard Manure (FYM), T₂- RDN through Vermicompost (VC), T₃- RDN through Poultry Manure (PM), T₄- RDN through FYM + Biofertilizers, T₅- RDN through Vermicompost + Biofertilizers, T₆- RDN through Poultry Manure + Biofertilizers (*Azotobacter* + PSB), T₇- Cow based bio-formulations and T₈- Control was laid out in randomized block design with three replications. Total sugar, reducing sugar, non-reducing sugar %, TSS and N content and its uptake in root and shoot was also recorded the highest in treatment T₅ whereas, P, K contents and their uptake in root and shoot

⁺⁺ Assistant Scientist;

*Corresponding author: E-mail: sumitbidlan2@gmail.com;

were recorded highest in treatment T₄. Soil organic carbon (0.69 %), available N (165.5 kg ha⁻¹), P (16.82 kg ha⁻¹), K (369.60 kg ha⁻¹) were observed maximum in treatment T₅ and remained at par with the treatments receiving organic manures conjointly with biofertilizers. Microbial population of soil was recorded the maximum in treatment T₅ (6.94×10^6 cfu count g⁻¹ of soil) which was at par to treatments T₁, T₂, T₄ and T₇ but significantly higher over the control (6.52×10^6 cfu count g⁻¹ of soil).

Keywords: Biofertilizers; microbial population; organic carbon; quality.

1. INTRODUCTION

Carrot (*Daucus carota*) is an important vegetable crop grown almost all over the world in spring, summer and autumn in temperate regions and during winter in tropical and sub-tropical regions. It falls under the family Umbelliferae. Genus *Daucus* contains about 60 species of which very few are cultivated [1]. Carrots possess a unique taproot structure that has been modified into a conical shape. The root is hollow and the stem is short, erect, and not very long. The leaves are quadripinnate [2]. The main reason why carrot roots are highly regarded as a food component is due to their abundant content of beta carotene, which serves as a precursor to vitamin A [3]. Additionally, carrots are a good source of carbohydrates and essential minerals such as calcium, phosphorus, and iron [4].

Organic farming relies on utilizing available knowledge, techniques, and materials to work in harmony with nature and sustainably manage natural resources within a specific agro-ecological system [5]. In organic agriculture, there are various sources of nitrogen, phosphorus, and potassium, with Farm Yard Manure (FYM), vermicompost, and poultry manure, along with biofertilizers, being the most popular options [6]. Biofertilizers and microorganisms play a very significant role in improving soil fertility by fixing atmospheric nitrogen, both, in association with plant roots and without it, solubilise insoluble soil phosphates and produces plant growth substances in the soil [7]. They are in fact being promoted to harvest the naturally available, biological system of nutrient mobilization [8]. Carrots are heavy feeder of nutrients, uptake around 100 kg of nitrogen, 50 kg of P₂O₅, and 110 kg of K₂O per hectare. They are highly responsive to nutrient availability and soil moisture [9]. While inorganic fertilization is important for promoting growth and development in plants but organic sources of nutrients offer advantages such as consistent and slow nutrient release, maintaining an ideal CN ratio (carbon nitrogen ratio), enhancing water retention capacity, and improving microbial

activity in the soil without adverse residual effects [10]. Additionally, organic manures have shown positive effects on the quality attributes of different vegetables. Recognizing the significance of organic manures, this experiment aimed to investigate the impact of various organic manures, either alone or in combination with inorganic fertilizers, on the yield and quality attributes of carrots. Furthermore, the study also assessed their effects on experimental field's soil physical and chemical properties.

2. MATERIALS AND METHODS

Initial soil samples were collected randomly from several field locations in a zig-zag pattern at 0-15 cm of soil depth before crop sowing with a hand auger. A combined sample was ground and passed through a 2 mm sieve and thereafter put away in the polythene pack for further analysis. The chemical analysis of soil samples for texture, pH, EC, available N, P, K and organic carbon, was done by standard methods as outlined by Antil. However, organic manures (FYM, PM and VC) were collected from the DDUCE-OF CCS HAU, Hisar, and incorporated into the field before sowing of crop. Before analysis, the manure's samples were air-dried, ground and passed through a 2 mm sieve. Digestion (with the di-acid mixture) is the main step followed, thereafter; standard methods were adopted for analysis purpose.

2.1 Experimental Site, Design and Field Management

The experiment was carried out at Research farm of Deen Dayal Upadhyaya Centre of Excellence for Organic Farming of CCS Haryana Agricultural University, Hisar during the *rabi* season 2022-23 under organic system of cultivation without using any chemicals is located at 29°8'20" N latitude and 75°42'04" E longitude at an elevation of 213 metre above mean sea level (MSL) in the sub-tropics of the country and characterized by semi-arid climatic zone. The soil of experimental field was sandy loam in texture

containing 157, 15.70 and 364.00 kg ha⁻¹ available nitrogen, phosphorus and potassium, respectively in 0-15 cm soil depth with EC 0.78 dS m⁻¹, pH 7.80 and organic carbon content 0.67 per cent.

The treatment combinations were setup as per recommended dose of Nitrogen for carrot crop in that is 60 Kg ha⁻¹. The experiment was laid out in a randomized block design (RBD) with three replications and 8 treatment combinations. Seed of carrot cultivar Punjab carrot 161 was sown in net plot of 4.5m × 4.0 m on 16 October, 2023. FYM, vermicompost, poultry manures and biofertilizers applied @ 9210, 3157, 2143 kg ha⁻¹ and 50 ml each of *Azotobacter* and PSB from department of biotechnology CCS HAU, Hisar for seed treatment. Cow based bio-formulations (Ghanajeevamrit and Jeevamrit) was applied @ 250 kg Ghanajeevamrit + 250 FYM kg/ha and 125 litre Jeevamrit in three split doses at 20, 40 and 60 DAS @ 5 %, 10% and 10%, respectively was applied in the treatments were T₁: RDN (recommended dose of nitrogen), through FYM, T₂: RDN through vermicompost, T₃: RDN through poultry manure, T₄: RDN through FYM + biofertilizers, T₅: RDN through vermicompost + biofertilizers, T₆: RDN through poultry manure + biofertilizer, T₇: cow based bio-formulations (Ghanajeevamrit and Jeevamrit) and T₈: control.

2.2 Data Collected

Soil parameters viz., pH, electric conductivity, organic carbon %, available NPK in soil and microbial population in soil before and after harvest was calculated. Nutrient content and uptake by carrot roots and leaves were also analysed after the harvest.

3. RESULTS AND DISCUSSION

Application of organic manures alone and with biofertilizers significantly influence the soil parameters i.e., pH, EC, OC%, available NPK, soil microbial population and nutrient content and uptake by roots and leaves.

The significant maximum value of nitrogen content and uptake by roots and leaves (2.30, 2.80% and 71.04, 53.12 kg/ha, respectively) were recorded in T₅ with the application of RDN through vermicompost and seed was treated with the biofertilizers (*Azotobacter* + PSB) which was statistically higher than other treatments except T₄, T₂ and T₁ respectively (Table 1). The

significant higher value of phosphorus uptake by roots (25.62 kg/ha) was recorded in T₄ with the application of RDN through FYM and seed treated with the biofertilizers (*Azotobacter* + PSB) which was statistically at par with treatment T₁, and T₆ whereas, P uptake by leaves was recorded in T₅ with the application of RDN through vermicompost and seed treated with the biofertilizer (*Azotobacter* + PSB) which was statistically at par with treatment T₄, T₂ and T₁ (Table 1). K content and uptake were recorded maximum with treatment T₄ (2.40, 2.20 % and 68.33 and 37.59 kg/ha, respectively) which was closely followed by treatment T₁ (Table 1). However, the minimum nutrient content and uptake by roots and leaves was recorded in control in which no external manures or formulations were applied. The nutrient content of soil might have enhanced due to breakdown of organic sources of manures which become readily accessible to plants and increase the concentration of NPK in plants. Furthermore, applications of biofertilizers might have enhanced the root proliferation coupled with increased availability of nutrients in the rhizosphere by N₂ fixation, P and K solubilization which led to augmented nutrient uptake by the plants. These findings confirm the results of Mansuri *et al.* [11]. The solubilization and mobilization of native nutrients and micronutrients by soil micro fauna, chelation of complex intermediate organic compounds formed during the breakdown of organic manure, accumulation of various nutrients in different plant parts may be responsible for the increased nutrient uptake with organic manure and biofertilizers [12].

The data summarized in the Table 2 showed the result of pH, EC and OC% of soil which was recorded in the range of 7.60 to 7.87, 0.74 to 0.78 (dS/m) and 0.67 to 0.69%. Different treatments of organic manures alone or with biofertilizers and cow based bio-formulations have non-significant effect on the pH, EC and OC % of soil after harvest. Sur *et al.* (2010) who have also reported the beneficial effects of organic manures over inorganic fertilizers. The maximum available NPK in soil (164.9, 16.82 and 369.3 kg/ha) was observed with the T₅, application of RDN through vermicompost and seed treated with biofertilizers (*Azotobacter*+PSB) which was significantly higher than other treatments except T₂ and T₁, whereas, available N was also at par with T₄ and available K also with T₃, T₄ and T₆. Whereas, minimum available N (155.1 kg/ha) was found in control (Table 2). Significantly maximum soil microbial population

(6.94×10^6 cfu/g) was recorded with T5, the application of RDN through vermicompost along with biofertilizers (*Azotobacter* + PSB) used as seed treatment, which was closely followed by the T₁, T₂, T₄ and T₇, whereas, minimum soil microbial population recorded in control treatment. However, all treatments including control significantly improved soil microbial population than observed before sowing ($5.56 \times$

10^6). Application of solely organic manures or in combination with biofertilizers and cow based bio-formulations enhanced soil organic matter availability and provided substrate that act as an energy source for soil microbes growth and increased bacterial multiplication. The addition of organic matter into soil resulted in increasing soil microbial populations and soil biological activity Weon et al., [13] and Alvarez et al., [14].




Fig. 1. Effect of organic manures and natural farming on N content and uptake by carrot roots and leaves




Fig. 2. Effect of organic manures and natural farming on NPK in soil after harvest

Table 1. Effect of organic manures and natural farming on NPK content and uptake by roots and leaves after harvest

Treatments	N content (%)		N uptake (kg ha ⁻¹)		P content (%)		P uptake (kg ha ⁻¹)		K content (%)		K uptake (kg ha ⁻¹)	
	Roots	Leaves	Roots	Leaves	Roots	Leaves	Roots	Leaves	Roots	Leaves	Roots	Leaves
T ₁ -	2.15	2.60	60.64	42.33	0.88	0.84	24.82	13.69	2.31	2.10	65.15	34.24
T ₂ -	2.20	2.72	63.57	50.17	0.77	0.77	22.25	14.20	1.93	1.66	55.77	30.62
T ₃ -	1.83	2.39	48.10	34.07	0.83	0.80	21.82	11.39	2.10	1.82	55.20	25.91
T ₄ -	2.30	2.65	65.48	45.28	0.90	0.86	25.62	14.69	2.40	2.20	68.33	37.59
T ₅ -	2.30	2.80	71.04	53.12	0.79	0.78	24.40	14.80	2.00	1.63	61.78	30.92
T ₆ -	1.85	2.39	51.35	36.75	0.85	0.81	23.59	12.44	2.14	1.90	59.39	29.18
T ₇ -	1.53	2.20	39.64	34.12	0.75	0.74	19.43	11.49	1.90	1.54	49.23	23.92
T ₈ -	1.32	2.09	32.50	32.03	0.72	0.71	17.73	10.86	1.84	1.50	45.30	22.95
SEm (±)	0.9	0.24	0.90	0.63	0.03	0.02	0.37	0.20	0.05	0.03	1.20	1.40
CD at 5 %	0.22	0.35	9.20	7.92	0.06	0.05	1.85	1.63	0.21	0.14	5.70	4.23

Table 2. Effect of organic manures and natural farming on pH, EC, OC, available NPK and soil microbial population in soil after harvest

Treatmentss	pH	EC (dS/m)	OC (%)	Available NPK (kg ha ⁻¹)			Soil microbial population (10 ⁶ cfu count g ⁻¹ soil)
				N (kg ha ⁻¹)	P (kg ha ⁻¹)	K (kg ha ⁻¹)	
T ₁ -	7.63	0.77	0.68	163.6	16.08	367.2	6.75 × 10 ⁶
T ₂ -	7.60	0.75	0.68	164.0	16.51	368.8	6.87 × 10 ⁶
T ₃ -	7.77	0.77	0.67	159.0	15.85	367.0	6.65 × 10 ⁶
T ₄ -	7.63	0.75	0.68	164.9	16.19	367.7	6.80 × 10 ⁶
T ₅ -	7.60	0.74	0.69	165.3	16.82	369.3	6.94 × 10 ⁶
T ₆ -	7.73	0.76	0.67	159.6	15.91	367.2	6.67 × 10 ⁶
T ₇ -	7.70	0.77	0.67	157.3	15.81	364.9	6.74 × 10 ⁶
T ₈ -	7.87	0.78	0.67	155.1	14.75	361.3	6.52 × 10 ⁶
SEm (±)	0.12	0.01	0.01	1.586	0.31	1.42	0.08
CD at 5 %	N.S	N.S	N.S	4.9	0.9	4.3	0.21
Initial	7.80	0.78	0.67	157	15.70	364.0	5.56 × 10 ⁶

4. CONCLUSION

The study's findings indicate that applying the recommended dose of nutrients (RDN) using vermicompost and biofertilizers leads to notable improvements in the quality attributes of carrots. Moreover, when these nutrient sources are used together (FYM, vermicompost and poultry manure), they contribute to enhancing soil properties and increasing the population of beneficial soil microorganisms. As a result, this combined approach has the potential to sustain higher NPK content in leaves and improve the microbial population of soil over the long term.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Salunkhee PK, Kadam SS. Handbook of vegetable Science and Technology; 2005.
2. Bhattarai BP, Maharjan A. Effect of organic nutrient management on the growth and yield of carrot (*Daucus carota* L.) and soil fertility status. Nepalese Journal of Agricultural Sciences. 2013;16.
3. Ahmad T, Cawood M, Iqbal Q, Ariño A, Batool A, Tariq RMS, Akhtar S. Phytochemicals in *Daucus carota* and their health benefits. Foods. 2019;8(9):424.
4. Ganapathi M. Influence of organics, micronutrients and plant Growth regulators on productivity potential in Carrot (*Daucus carota* L.). Department of crop physiology. M.Sc. Thesis submitted to the University of Agricultural Sciences, Dharwad; 2006.
5. Rajbhandari BP, Gautam BR. Bio-intensive farming system: a manual. Kathmandu: WOREC Nepal Publications; 1998.
6. Khayat M. Evaluation Effect of Farmyard Manure (FYM) to improve cereal crop yield. Journal of Crop Nutrition Science. 2021;7(1):59-67.
7. Bhardwaj D, Ansari MW, Sahoo RK, Tuteja N. Biofertilizers function as key player in sustainable agriculture by improving soil fertility, plant tolerance and crop productivity. Microbial cell factories. 2014;13:1-10.
8. Venkateshwarlu B. Role of bio-fertilizers in organic farming: Organic farming in rain fed agriculture: Central institute for dry land agriculture, Hyderabad. 2008;85-95.
9. Rani SM, Reddy, Malla K. Effect of different organic manures and inorganic fertilizers on growth, yield and quality of carrot (*Daucus carota* L.). Karnataka Journal of Agricultural Scienc. 2007;20:686-88.
10. Yadav A, Jakhar RK, Kumari N, Yadav GN, Kant R, Sharma SL, Kumar A. Response of organic manures and fertilizers on yield and quality of carrot under sandy soil condition. The Pharma Innovation Journal. 2022;11(4):1822-27.
11. Mansuri SA, Shakhela RR, Patel NK, Jat JS. Effect of different zinc enriched organics on nutrient content and uptake by summer pearl millet [*Pennisetum glaucum* L.] on loamy sand. Journal of Pharmacognosy and Phytochemistry. 2019;8(1):724-730.
12. Kumar A, Garhwal RS, Dinesh, Ankush. Impact of various organic and inorganic sources of fertilizers on yield, yield attributes, and nutrients accumulation in direct seeded basmati rice. Indian Journal of Ecology. 2022;49(2):435-439.
13. Weon HY, Kwon JS, Suh JS, Choi WY. Soil microbial flora and chemical properties as influenced by the application of pig manure compost. Korean journal of Soil Science Fertilizer. 1999;32:76–83.
14. Alvarez MB, Gagnè S, Antoun H. Effect of compost on rhizosphere microflora of tomato and on the incidence of plant growth promoting rhizobacteria. Applied Environmental Microbiology. 1995;61:194–199.

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109005>