



## **Effect of Long Term Soil Test Crop Response Based Nutrient Management in Wheat (*Triticum aestivum* L.) on Physical Properties in a Vertisol**

Divya Bhayal <sup>a\*</sup>, P. S. Kulhare <sup>a</sup>, G. S. Tagore <sup>a</sup>, A. K. Upadhyay <sup>a</sup>, Lalita Bhayal <sup>b</sup> and Aakash <sup>c</sup>

<sup>a</sup> Department of Soil Science, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh (482 004), India.

<sup>b</sup> Department of Agronomy, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh (482 004), India.

<sup>c</sup> Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh (221 005), India.

### **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/IJPSS/2022/v34i2231484

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91300>

**Original Research Article**

**Received 24 July 2022**  
**Accepted 26 August 2022**  
**Published 27 August 2022**

### **ABSTRACT**

A field experiment was carried out during *Rabi* season of 2020–21 and 2021–22 at the Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur research field to find out the effect of long term soil test crop response based nutrient management in wheat (*Triticum aestivum* L.) on physical properties in a Vertisol. The treatments details were T<sub>1</sub>; Control, T<sub>2</sub>; 120 N: 80 P<sub>2</sub>O<sub>5</sub>: 60 K<sub>2</sub>O kg ha<sup>-1</sup>, T<sub>3</sub>; Target yield 4.5 t ha<sup>-1</sup>, T<sub>4</sub>; Target yield 6.0 t ha<sup>-1</sup>, T<sub>5</sub>; Target yield 4.5 t + FYM 5 t ha<sup>-1</sup>, T<sub>6</sub>; Target yield 6.0 t + FYM 5 t ha<sup>-1</sup> and carried out in Randomized Block Design (RBD) with four replications. The results indicated that the STCR based treatment with target yield 6.0 t ha<sup>-1</sup> along with application of FYM 5 t ha<sup>-1</sup> recorded lowest bulk density (1.31, 1.33 and 1.35 Mg m<sup>-3</sup>) and particle density (2.62, 2.62 and 2.63 Mg m<sup>-3</sup>) at 0-15, 15-30 and 30-45 cm soil depth. The treatment with target yield 6.0 t ha<sup>-1</sup> along with application of FYM 5 t ha<sup>-1</sup> recorded highest porosity (50.07, 49.28 and 48.52 %) as well as water holding capacity (44.42, 40.16 and 37.44 %) at 0-15, 15-30 and 30-45 cm soil depth.

\*Corresponding author: E-mail: divyabhayal123@gmail.com;

The long term study based on STCR clearly demonstrated that balanced fertilizer application not only improved the physical properties of soil but also save the fertilizer without impairing soil health.

**Keywords:** FYM; bulk density; particle density; porosity; STCR; water holding capacity.

## 1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is the world's most widely cultivated food grain crop. Besides staple food for human beings, wheat straw also serves as good source of feed for animals [1]. It is grown on approximately 32 million hectares with a total production of 107.8 million tonnes and a productivity of 3370 kg ha<sup>-1</sup> [2]. In Madhya Pradesh, it is grown in about 49 lakh ha with an annual production of 125.6 lakh tonnes and productivity of 2563 kg ha<sup>-1</sup> [2]. The major challenge confronting India during the 21<sup>st</sup> century is to produce enough food, fodder, fiber and fuel to meet the diversified need of ever-increasing human population as well as animal's population of the country. The application of inorganic fertilizers leads to the increasing crop yield due to the immediate supply of plant nutrients [3], but its use over a long run may decrease the stability of macro and micro aggregate, moisture retention and increase bulk and particle density thus decreasing the productivity [4]. Fertilization of crops based on generalized recommendation leads to under fertilization or over fertilization, results in deteriorating physical properties (bulk density, particle density, porosity etc.), lower productivity along with environmental pollution. Therefore, it is essential to protect the soil health by adopting balanced fertilization through soil testing and organic source as an integrated manner. Among the various methods of fertilizers recommendation, soil test based nutrient recommendation cares crop demand, response of nutrients and expected crop yield. Ramamoorthy et al. [5] developed the methodology of soil test based nutrient recommendation for targeted yield. It is a more quantitative, precise and meaningful approach because it involves combined use of soil and plant analysis, which provide information on real balance between applied nutrient and available nutrients of soil [6]. The inorganic fertilizer along with FYM plays an important role in increasing crop yields, since inorganic fertilizers fulfilled crop nutrients need quickly and FYM provides both macro and micronutrients gradually and continuously to crop. FYM improve physical properties of soil such as aggregation, permeability and water holding capacity. It contains all the essential elements which mitigate

the ill effect of imbalanced use of fertilizers [7]. Long term fertilizer experiment provides best possible means to study the changes in soil properties, dynamics of nutrient processes and future strategies for maintaining soil health [8]. Such experiments in India suggested that under continuous cropping, changes in soil fertility due to imbalanced fertilization may be recognized as an important factor limiting crop yields. Thus, it is essential to measure the changes in soil properties that occurred due to soil test based fertilizer application. Therefore, the present experiment was undertaken with the objectives to study the effect of long term STCR based nutrient management in wheat on soil bulk density, particle density, water holding capacity and porosity.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Details

The experiment was started in Rabi 2020-21 and 2021-22 at the Jawaharlal Nehru Krishi Vishwa Vidyalaya (JNKVV), Jabalpur research field. The experimental site was situated in the South-Eastern part of Madhya Pradesh at 23°13' North latitude, 79°57' East longitudes and at an elevation of 393 meter above mean sea level. The soil samples (0-15, 15-30 and 30-45 cm depth) were collected from the experimental site. The experiment was laid out in randomized block design (RBD) with six treatments each replicated four times. The treatment details were T<sub>1</sub>: Control, T<sub>2</sub>:120 N: 80 P<sub>2</sub>O<sub>5</sub>: 60 K<sub>2</sub>O kg ha<sup>-1</sup>, T<sub>3</sub>: target yield 4.5 t ha<sup>-1</sup>, T<sub>4</sub>: target yield 6.0 t ha<sup>-1</sup>, T<sub>5</sub>: target yield 4.5 t + FYM 5 t ha<sup>-1</sup> and T<sub>6</sub>: target yield 6.0 t + FYM 5 t ha<sup>-1</sup>. The STCR based fertilizer doses were calculated from the fertilizer adjustment equations for target yield 4.5 and 6.0 t ha<sup>-1</sup> of wheat. The fertilizer adjustment equations are given below:

$$\begin{aligned} FN &= 4.40 T - 0.40 SN \\ FP_2O_5 &= 4.00 T - 5.73 SP \\ FK_2O &= 2.53 T - 0.16 SK \end{aligned}$$

Whereas:

FN = Fertilizer nitrogen (kg ha<sup>-1</sup>); FP<sub>2</sub>O<sub>5</sub>= Fertilizer phosphorus (kg ha<sup>-1</sup>); FK<sub>2</sub>O = Fertilizer potassium (kg ha<sup>-1</sup>); T = Desired yield target (t ha<sup>-1</sup>); SN = Available soil nitrogen (kg ha<sup>-1</sup>); SP =

Available soil phosphorus ( $\text{kg ha}^{-1}$ ); SK= Available soil potassium ( $\text{kg ha}^{-1}$ ).

## 2.2 Soil Sampling and Analysis

The processed soil sample were analyzed for physical properties (bulk density, particle density, water holding capacity and porosity) from the following methods: the bulk density of soil at 0-15, 15-30 and 30-45 cm were determined using core sampler method [9], particle density of soil at 0-15, 15-30 and 30-45 cm were determined by Pycnometer [10], water holding capacity at 0-15, 15-30 and 30-45 cm were determined by Keen's Box [9] and porosity of soil at 0-15, 15-30 and 30-45 cm were determined using empirical method as described by Gupta and Dhakshinamoorthy, [11].

## 3. RESULTS AND DISCUSSION

### 3.1 Bulk Density

The application of STCR based NPK level as GRD, target yield of  $4.5 \text{ t ha}^{-1}$ , target yield of  $6.0 \text{ t ha}^{-1}$ , target yield of  $4.5 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  and target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  had significantly decreased the bulk density over control at all soil depth i.e. 0-15, 15-30 and 30-45 cm in the pooled data (Table 1 and Figure 1). However, the application of nutrient for target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$ , target yield of  $6.0 \text{ t ha}^{-1}$  and target yield of  $4.5 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  recorded significantly decreased bulk density over GRD at all soil depth except 30-45 cm in the pooled data. Target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  and target yield of  $4.5 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  had significantly less bulk density over target yield of  $6.0 \text{ t ha}^{-1}$  at all soil depth in the pooled data. The lowest bulk density was observed at 0-15 cm depth  $1.31 \text{ Mg m}^{-3}$ , at 15-30 cm depth  $1.33 \text{ Mg m}^{-3}$ , and at 30-45 cm depth  $1.35 \text{ Mg m}^{-3}$  with nutrient application for target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  in pooled data.

The significant reduction of bulk density in all soil depth with nutrient application for target yield of  $4.5 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  and target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  than NPK alone (as GRD, target yield of  $4.5 \text{ t ha}^{-1}$ , target yield of  $6.0 \text{ t ha}^{-1}$ ) might be due to the addition of FYM enhanced

the OC content which increased root biomass production that might have augmented organic matter content of the soil hence reduced the BD. The addition of FYM promotes the total porosity of soils as the microbial decomposition products of organic manures such as polysaccharides and bacterial gums are known to act as soil particle binding agents. These binding agents increase the porosity and decrease the bulk density of the soil by improving the aggregation. The lowest bulk density in target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  at all soil depths was attributed application of FYM as  $5 \text{ t ha}^{-1}$  in combination with nutrient application for target yield of  $6.0 \text{ t ha}^{-1}$  could provide a better environment for the crop development and health of the soil. It could retain for the crop development and health of the soil than NPK alone retaining more water due to increased porosity. These results of decrease in bulk density due to addition of FYM are line with results reported by Bandyopadhyay et al., [12], Thakur et al., [13], Gudadhe et al., [14], Moharana et al., [15].

### 3.2 Particle Density

The application of STCR based NPK level as GRD, target yield of  $4.5 \text{ t ha}^{-1}$ , target yield of  $6.0 \text{ t ha}^{-1}$ , target yield of  $4.5 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  and target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  reported slightly decreased the particle density over control at all soil depth in the pooled data but all the treatments were found non-significant to each other (Table 1). The lowest particle density was observed at 0-15 cm depth  $2.62 \text{ Mg m}^{-3}$ , at 15-30 cm depth  $2.62 \text{ Mg m}^{-3}$ , and at 30-45 cm depth  $2.63 \text{ Mg m}^{-3}$  with nutrient application for target yield of  $6.0 \text{ t ha}^{-1} + \text{FYM } 5 \text{ t ha}^{-1}$  in pooled data respectively (Table 1). The particle density is depends on texture and exclusively on the mineralogical composition of the soil material and content of organic matter. The continuous manuring in soil might result in slight decrease in particle density of soil at all soil depths. The particle density of soils slightly increased with depth possibly due to lower organic matter in sub-surface soil reported by Ram et al., [16]. The particle density did not show any significant change due to continuous application of fertilizer and manures. Similar findings are also reported by Nandapure et al., [17].




Fig. 1. Effect of STCR based NPK levels on Bulk density at different depth in soil

Table 1. Effect of STCR based NPK levels on post harvest soil bulk density and particle density (pooled data)

Treatment	Bulk density (Mg m <sup>-3</sup> )			Particle density (Mg m <sup>-3</sup> )		
	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
T <sub>1</sub> Control	1.38	1.44	1.46	2.66	2.67	2.67
T <sub>2</sub> GRD (120 N: 80 P <sub>2</sub> O <sub>5</sub> : 60 K <sub>2</sub> O kg ha <sup>-1</sup> )	1.36	1.41	1.43	2.64	2.64	2.65
T <sub>3</sub> Target Yield 4.5 t ha <sup>-1</sup>	1.36	1.40	1.42	2.63	2.64	2.64
T <sub>4</sub> Target Yield 6 t ha <sup>-1</sup>	1.34	1.39	1.42	2.63	2.64	2.63
T <sub>5</sub> Target Yield 4.5 t ha <sup>-1</sup> + FYM 5 t ha <sup>-1</sup>	1.32	1.36	1.38	2.62	2.63	2.64
T <sub>6</sub> Target Yield 6 t ha <sup>-1</sup> + FYM 5 t ha <sup>-1</sup>	1.31	1.33	1.35	2.62	2.62	2.63
SEm ±	0.005	0.008	0.008	0.01	0.01	0.01
CD (p=0.05)	0.01	0.02	0.02	NS	NS	NS

### 3.3 Water Holding Capacity

The water holding capacity at 0-15 cm depth at target yield of 4.5 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> and target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> was found significantly superior to control, GRD in pooled data. While WHC with target yield of 4.5 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> and target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> also significantly increased at 15-30 cm soil depth (Table 2 and Figure 2). However, the WHC at target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> was found significantly superior to target yield of 4.5 t ha<sup>-1</sup> and target yield of 6.0 t ha<sup>-1</sup> but it was found at par with target yield of 4.5 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> in pooled data. The water holding capacity at 30-45 cm with target yield of 6.0 t ha<sup>-1</sup>, target yield of 4.5 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> and target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> and GRD was found at par during both the years. However, the water holding capacity with target yield of 4.5 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> and target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> were found significantly superior to GRD

and target yield of 4.5 t ha<sup>-1</sup> but the two treatments were found at par in pooled data. The maximum water holding capacity was observed at 0-15 cm depth 44.42 %, 15-30 cm depth 40.16 % and at 30-45 cm depth 37.44 % with nutrient application for target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup>.

The continuous application of FYM decrease bulk density, increase porosity and reduce crust formation thus increase macro and micro pores in soil which help to increase water holding capacity. The higher WHC of the surface soil was attributed to the effect of increased proportion of aggregation. As the depth increase, there will be less aggregation and more compaction in the soil as compared to surface layers which decrease the water holding capacity. The decrease in water holding capacity with increase in depth was also reported by Hati et al., [18] and Pant and Ram [19].




Fig. 2. Effect of STCR based NPK levels on water holding capacity at different depth in soil

Table 2. Effect of STCR based NPK levels on post harvest water holding capacity (WHC) and porosity (pooled data)

Treatment	WHC %			Porosity %		
	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
T <sub>1</sub> Control	37.26	34.26	31.57	48.04	46.01	45.43
T <sub>2</sub> GRD (120 N: 80 P <sub>2</sub> O <sub>5</sub> : 60 K <sub>2</sub> O kg ha <sup>-1</sup> )	38.55	36.06	34.17	48.53	46.65	45.88
T <sub>3</sub> Target Yield 4.5 t ha <sup>-1</sup>	39.84	36.93	34.10	48.50	46.89	46.12
T <sub>4</sub> Target Yield 6 t ha <sup>-1</sup>	40.26	37.68	35.36	48.91	47.38	46.12
T <sub>5</sub> Target Yield 4.5 t ha <sup>-1</sup> + FYM 5 t ha <sup>-1</sup>	42.66	38.44	36.57	49.64	48.23	47.60
T <sub>6</sub> Target Yield 6 t ha <sup>-1</sup> + FYM 5 t ha <sup>-1</sup>	44.42	40.16	37.44	50.07	49.28	48.52
SEm ±	0.67	0.83	0.81	0.17	0.37	0.39
CD (p=0.05)	1.93	2.41	2.33	0.49	1.07	1.14

### 3.4 Porosity

The data presented in Table 2 indicated that the porosity with target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> was found significantly superior to target yield of 6.0 t ha<sup>-1</sup> in pooled data at 0-15 cm and 15-30 cm soil depth. The porosity at target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> also found significant over target yield of 4.5 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup>. While the porosity with target yield of 4.5 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> and target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup> were found significantly superior over control at 30-45 cm but the treatment were found at par at 30-45 cm soil depth in pooled data. The maximum soil porosity was observed at 0-15 cm 50.07%, 15-30 cm 49.28% and 30-45 cm 48.52% with nutrient application for target yield of 6.0 t ha<sup>-1</sup> + FYM 5 t ha<sup>-1</sup>.

The addition of FYM promotes the total porosity of soils as the microbial decomposition products

of organic manures such as polysaccharides and bacterial gums are known to act as soil particle binding agents. These binding agents increase the porosity and decrease the bulk density of the soil by improving the aggregation. The higher soil porosity of the soil of the surface layer than sub surface layer it might be due to ready exchange of O<sub>2</sub> and CO<sub>2</sub> between the soil and atmosphere thereby, promoting better root growth in soil. Similar findings were also reported by Hati et al., [18], Thakur et al., [13] and Moharana et al., [15].

### 4. CONCLUSION

In the present research, target yield 6.0 t ha<sup>-1</sup> along with the application of FYM 5 t ha<sup>-1</sup> recorded the lowest bulk density and particle density, highest porosity and water holding capacity. Based on the two years of research, it is concluded that prescription-based fertilizer

application i.e. target yield 6.0 t ha<sup>-1</sup> along with the application of FYM 5 t ha<sup>-1</sup> enhanced the soil physical properties and saved the fertilizer without negatively impacting the soil health.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Sarwar N, Maqsood M, Mubeen K, Shehzad M, Bhullar MS, Qamar R and Akbar N. Effect of different levels of irrigation on yield and yield components of wheat cultivars. *Pakistan Journal of Agriculture Science*. 2006;47:371-734.
2. Anonymous. Agriculture statistics at a glance. Directorate of economics and statistics, Department of Agriculture and Cooperation, ministry of Agriculture, government of India. 2021;4.
3. López Pérez A, Casanova E, Chacon LA, Paz P, Guerrero JR. Residual effect of three phosphate rocks from Tachina [Venezuela] in a greenhouse experiment with maize (*Zea mays* L.) as indicator plant. *Revista Científica UNET*. 1990; 4(1-2):29-48.
4. Sarkar S, Singh SR, Singh RP. The effect of organic and inorganic fertilizers on soil physical condition and the productivity of a rice-lentil cropping sequence in India. *The Journal of Agricultural Science*. 2003; 140(4):419-425.
5. Ramamoorthy B, Narasimham RL and Dinesh RS. Fertilizer application for specific yield targets on Sonora 64 (wheat). *Indian Farming*. 1967;17(5):43-45.
6. Sharma VK, Pandey RN, Kumar S, Chobhe KA and Chandra S. Soil test crop response based fertilizer recommendation under integrated nutrient management for higher productivity of pearl millet (*Pennisetum glaucum*) and (*Triticum aestivum*) wheat under long term experiment. *Indian Journal of Agricultural Sciences*. 2016;86:1076-1081.
7. Ray PK, Jana AK, Maitra DN, Saha MN, Chaudhury J, Saha S, Saha AR. Fertilizer prescriptions on soil test basis for jute, rice and wheat in a Typic Ustochrept. *Journal of the Indian Society of Soil Science*. 2000;48:79-84.
8. Swarup A. Role of soil organic matter and balance fertilization in combating land degradation in dry area and sustaining crop productivity. Food security and climate change in dry areas Proceedings of an International Conference 1-4 February 2010, Amman, Jordan (ICARDA). 2010;282-290.
9. Richards LA. Diagnosis and improvement of Saline and Alkaline soils USDA Agriculture Handbook 60, Washington D.C.; 1954.
10. Black CA. Methods of Soil Analysis, part II, Chemical and properties, No. 9 series, Agronomy, American Soc. Agron. Madison, Wisconsin, USA; 1965.
11. Gupta RP and Dhakshinamoorthy C. Procedures for Physical Analysis of Soils and Collection of Agrometeorological Data, Division of Agricultural Physics, Indian Agricultural Research Institute, New Delhi; 1980.
12. Bandyopadhyay KK, Mishra AK, Ghosh PK and Hati KM. Effect of integrated use of farmyard manure and chemical fertilizers on soil physical properties and productivity of soybean. *Soil & Tillage Research*. 2010;110(1):115-125.
13. Thakur Rishikesh, Sawarkar SD, Vaishya UK and Muneshwar Singh Impact of continuous use of inorganic fertilizers and organic manure on soil properties and productivity under soybean-wheat intensive cropping of a Vertisol. *Journal of the Indian Society of Soil Science*. 2011; 59(1):74-81.
14. Gudadhe N, Dhonde MB and Hirwe NA. Effect of integrated nutrient management on soil properties under cotton-chickpea cropping sequence in Vertisols of Deccan plateau of India. *Indian Journal of Agricultural Research*. 2015;49(3):207-14.
15. Moharana PC, Sharma BM and Biswas Dr. Changes in the soil properties and availability of micronutrients after six-year application of organic and chemical fertilizers using STCR-based targeted yield equations under pearl millet-wheat cropping system. *Journal of Plant Nutrition*. 2017;40(2):165-176.
16. Ram D, Ram T and Shubhash Chand. Characterization and Classification of flood-prone soils of eastern plains of Rajasthan for their corrective measures. *J. of Indian Society of Soil Science*. 2010; 58(2):228-232.

17. Nandapure SP, Sonune BA, Gabhane VV, Katkar RN and Patil RT. Long term effects of integrated nutrient management on soil physical properties and crop productivity in shorghum-wheat cropping sequence in a vertisol. Indian J. Agric. Res. 2011;45(4): 336-340.
18. Hati KM, Swarup A, Dwivedi AK, Misra AK. Changes in soil physical properties and organic carbon status at the topsoil horizon of a vertisol of central India after 28 years of continuous cropping, fertilizer manuring. Agriculture Ecosystem & Environment. 2007;119:127-134.
19. Pant PK and Ram S. Long-term Manuring and Fertilization Effects on Soil Physical Properties after Forty Two Cycles under Rice-wheat System in North Indian Mollisols. Int. J. Curr. Microbiol. App. Sci. 2018;7(7):232-240.

© 2022 Bhayal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/91300>*