



# Simple Tests for $n$ -th Roots of Natural Numbers being Natural Numbers and Elementary Methods to Determine Their Values

Bernd E. Wolfinger <sup>a\*</sup>

<sup>a</sup> Computer Science Department, University of Hamburg, Germany.

*Author's contribution*

*The sole author designed, analysed, interpreted and prepared the manuscript.*

*Article Information*

DOI: 10.9734/JAMCS/2024/v39i11860

**Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111735>

**Original Research Article**

*Received: 12/11/2023*

*Accepted: 16/01/2024*

*Published: 27/01/2024*

## Abstract

In this paper we tackle the challenging problem to determine, in a simple but reliable way, whether – for a given, arbitrary number  $x$ ,  $x \geq 2$  – the  $n$ -th root of  $x$  produces a rational or an irrational result, i.e. we determine whether  $\sqrt[n]{x} \in \mathbf{Q}$  or  $\sqrt[n]{x} \notin \mathbf{Q}$ . To solve this problem in a straightforward manner we make use of the prime factorization of  $x$ . As a main contribution we present a generally applicable algorithm to decide whether  $\sqrt[n]{x} \in \mathbf{Q}$  (for  $n, x \in \mathbf{N} \setminus \{1\}$ ) and if so, to determine the resulting value. Moreover, we design several tests which can be applied to determine, for which values of  $n$ ,  $\sqrt[n]{x} \in \mathbf{Q}$  if the natural number  $x$  satisfies a given set of properties. Quite often the tests proposed will allow us to answer the question “ $\sqrt[n]{x} \in \mathbf{Q}$ ?” in a matter of seconds. Finally, we demonstrate that, for a very high percentage of all natural numbers  $x$ ,  $x \geq 2$ , it is impossible to find even a single  $n \in \mathbf{N}$ ,  $n \geq 2$  such that  $\sqrt[n]{x} \in \mathbf{Q}$ .

\*Corresponding author: Email: [bernd.wolfinger@uni-hamburg.de](mailto:bernd.wolfinger@uni-hamburg.de);

*Keywords: Number theory; prime factorization; efficient tests for  $\sqrt[n]{x}$  ( $n, x \in \mathbf{N}, n, x \geq 2$ ) being a rational number; replacement of Newton's method to determine a rational value of  $\sqrt[n]{x}$ .*

## 1 Introduction

In [1] it has been demonstrated that it is astonishingly simple to answer the question “for which  $n \in \mathbf{N}, n \geq 2$  the  $n$ -th root of  $x$  yields to a rational result, i.e.  $\sqrt[n]{x} \in \mathbf{Q}$ ?” This question can be answered easily if one makes use of the (unique) prime factorization of  $x$  possessing the following structure:

$$x = p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_i^{k_i} \cdot \dots \cdot p_m^{k_m} \tag{1}$$

where  $p_i$  are prime numbers  $\forall i \in \{1, 2, \dots, m\}$  and  $p_i \neq p_j \forall i \neq j, m \geq 1, k_i \in \mathbf{N} \forall i \in \{1, 2, \dots, m\}$ .

*Note:*  $k_i$  to be read as  $k_i$ .

The result derived in [1] was:

For any  $n, x \in \mathbf{N}, n, x \geq 2$ :

$$\sqrt[n]{x} \in \mathbf{Q} \Leftrightarrow n \in \text{CD}(\{k_1, k_2, \dots, k_m\}), \tag{2}$$

where  $\text{CD}(M)$ , for  $M$  being a subset of  $\mathbf{N}$ , denotes the set of common divisors  $c, c \geq 2$ , of all elements of the set  $M$ .

Therefore, if  $m \geq 2$ :

$\text{CD}(\{k_1, k_2, \dots, k_m\}) := \{c \in \mathbf{N}, c \geq 2 \mid \forall k_i \exists v_i = v_i(k_i) \in \mathbf{N}: c \cdot v_i = k_i\}$ ,  
and, if  $m=1$ :

$$\text{CD}(\{k_1\}) := \{c \in \mathbf{N}, c \geq 2 \mid \exists v \in \mathbf{N}: c \cdot v = k_1\}.$$

It should be noted that for answering our question “ $\sqrt[n]{x} \in \mathbf{Q}$ ?” by applying eq. (2) the values of the  $p_i$  appearing in eq. (1) are of no interest but only the values of the exponents  $k_i$  are relevant.

The existence of  $\sqrt[n]{x}$  has been proven, e.g., in [2].

EXAMPLES:

- **Ex1.1:**  $x = p^{10}$ ,  $p$  denoting an arbitrary prime number

$$\Rightarrow \text{CD}(\{10\}) = \{2, 5, 10\} \text{ and, therefore, } \sqrt{p^{10}} \in \mathbf{Q}, \sqrt[5]{p^{10}} \in \mathbf{Q}, \sqrt[10]{p^{10}} \in \mathbf{Q}.$$

In [1] it has also been proven that, if  $\sqrt[n]{x} \in \mathbf{Q}$  and  $x \in \mathbf{N}$ , then this also directly implies that  $\sqrt[n]{x} \in \mathbf{N}$ . This will be taken into account by us in the following.

- **Ex1.2:**  $x = 2^{10}$ ,

$$\Rightarrow \text{CD}(\{10\}) = \{2, 5, 10\} \text{ cf. above and, therefore, } \sqrt{2^{10}} = 2^{10:2} = 2^5 = 32 \in \mathbf{N}, \sqrt[5]{2^{10}} = 2^{10:5} = 2^2 = 4 \in \mathbf{N}, \sqrt[10]{2^{10}} = 2^{10:10} = 2^1 = 2 \in \mathbf{N}.$$

- **Ex1.3:**  $x = 331776 = 2^{12} \cdot 3^4$ . Therefore, we have to determine  $\text{CD}(\{12, 4\}) = \{2, 4\}$ . This implies that  $\sqrt{331776} \in \mathbf{N}$  and  $\sqrt[4]{331776} \in \mathbf{N}$ . But,  $\sqrt[3]{331776} \notin \mathbf{Q} \forall n \notin \{2, 4\}$ .

One of the main goals of this contribution is to develop simple tests to determine – for arbitrary natural numbers  $n, x \in \mathbf{N}, n \geq 2$  and  $x \geq 2$  – whether  $\sqrt[n]{x} \in \mathbf{Q}$  and therefore even  $\sqrt[n]{x} \in \mathbf{N}$ . For our tests we need the prime factorization of  $x$ , according to eq. (1).

Algorithms solving the problem of prime factorization of natural numbers are well-known (for details see, e.g., [3,4,5]).

The prime factorization of  $x$  can be derived (in particular with the help of a computer) without much effort, if the numerical value of  $x$  is not very high. However, it is obvious that if  $x$  is the product of two extremely large prime numbers, prime factorization of  $x$  may become very difficult or even practically unfeasible. This fact is used, e.g. in the field of cryptography, in particular to construct very secure encryption algorithms, such as the RSA algorithm (see [6]).

To simplify our notation and argumentation throughout this paper let us introduce some basic notations and abbreviations.

We denote by:

$$\mathbf{N}_{\geq 2} := \{x \in \mathbf{N} \mid x \geq 2\}.$$

Moreover, for a natural number  $x \in \mathbf{N}_{\geq 2}$  we want to introduce notations to denote the set of natural numbers  $n \in \mathbf{N}_{\geq 2}$ , for which  $\sqrt[n]{x}$  results in a rational root, i.e.  $\sqrt[n]{x} \in \mathbf{Q}$ . So let denote

$$\mathbf{N}_{\text{rat}_R}(x) \text{ the set of natural numbers } n \in \mathbf{N}_{\geq 2} \text{ such that for given } x \in \mathbf{N}_{\geq 2} : \sqrt[n]{x} \in \mathbf{Q} \forall n \in \mathbf{N}_{\text{rat}_R}(x)$$

and – analogously – let denote

$$\mathbf{N}_{\text{irrat}_R}(x) \text{ the set of natural numbers } n \in \mathbf{N}_{\geq 2} \text{ such that for given } x \in \mathbf{N}_{\geq 2} : \sqrt[n]{x} \notin \mathbf{Q} \forall n \in \mathbf{N}_{\text{irrat}_R}(x).$$

Our notations  $\mathbf{N}_{\text{rat}_R}(x)$  and accordingly  $\mathbf{N}_{\text{irrat}_R}(x)$  are used to denote – for a specific  $x \in \mathbf{N}_{\geq 2}$  – the set of natural numbers  $x \in \mathbf{N}_{\geq 2}$  leading to rational or irrational results, respectively, of the  $n$ -th root  $\sqrt[n]{x}$ . We generalize this notation by replacing  $x$  by a set  $S \subseteq \mathbf{N}_{\geq 2}$ , e.g., in which the elements have common properties. Then,  $\mathbf{N}_{\text{rat}_R}(S)$  denotes the subset of  $\mathbf{N}_{\geq 2}$  such that  $\forall x \in S$  and  $\forall n \in \mathbf{N}_{\text{rat}_R}(S) : \sqrt[n]{x} \in \mathbf{Q}$ . And the meaning of  $\mathbf{N}_{\text{irrat}_R}(S)$  will be analogous.

The rest of this paper is structured as follows: In Section 2 we will start by giving a short survey of the methodology which underlies our investigation of  $n$ -th roots and which differs significantly from the methodology conventionally used up to now. Thereafter, in Section 3, we introduce simple tests which allow us, in a fast and efficient way, to precisely determine – for any given  $x \in \mathbf{N}_{\geq 2}$  – the sets  $\mathbf{N}_{\text{rat}_R}(x)$  and  $\mathbf{N}_{\text{irrat}_R}(x)$ . We will demonstrate the ease of use of our tests by various examples. In Section 4, investigations will be presented regarding the set of natural numbers the  $n$ -th roots of which all are irrational. The paper will conclude with a short summary and outlook.

## 2 Basic Methodology Used

Let us now shortly summarize the main methodology underlying this paper.

Basically, we will apply the following

**ALGORITHM:** Calculation of the value  $\sqrt[n]{x}$  (for  $x, n \in \mathbf{N}_{\geq 2}$  if  $\sqrt[n]{x} \in \mathbf{Q}$ )

- **Goals:** Given  $n, x$ . Test whether  $\sqrt[n]{x} \in \mathbf{Q}$  or  $\sqrt[n]{x} \notin \mathbf{Q}$ .

If  $\sqrt[n]{x} \in \mathbf{Q}$  then determine the value of  $\sqrt[n]{x}$ .

- **STEP 1:** For given  $x \in \mathbf{N}_{\geq 2}$  determine the prime factorization of  $x$ .

Result:  $x = p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m}$ .

- **STEP2:** Determine the common (integer) divisors for the set of exponents  $\{k_1, k_2, \dots, k_m\}$  occurring in the prime factorization of  $x$ , i.e. we determine  $CD(\{k_1, k_2, \dots, k_m\})$  using the definition of the set  $CD$  as introduced above.
- **STEP 3:** For  $\forall n \in CD(\{k_1, k_2, \dots, k_m\})$  we can conclude  $\sqrt[n]{x} \in \mathbf{Q}$ ; and for all other values of  $n \in \mathbf{N}_{\geq 2}$ , we now know that  $\sqrt[n]{x} \notin \mathbf{Q}$ .
- **STEP 4:** If  $\sqrt[n]{x} \in \mathbf{Q}$  then  $\sqrt[n]{x} = p_1^{k_1/n} \cdot p_2^{k_2/n} \cdot \dots \cdot p_m^{k_m/n}$ .

*Remark:* To the best of the knowledge of the author of this contribution, up to now, there does not exist any publication which presented an algorithm, oriented to a similar purpose, being as simple and efficient as the algorithm suggested by us.

### 3 Tests for $\sqrt[n]{x}$ Being an Irrational Number (for $n, x \in \mathbf{N}, n, x \geq 2$ )

In our introduction we have already demonstrated that for an arbitrary  $x \in \mathbf{N}_{\geq 2}$  possessing a prime factorization according to eq. (1) only the exponents  $k_1, k_2, \dots, k_m$  appearing in the factorization are relevant and the set  $\mathbf{N}_{\text{rat\_R}}(x)$  can be determined by  $\mathbf{N}_{\text{rat\_R}}(x) = CD(\{k_1, k_2, \dots, k_m\})$ .

And, similarly,  $\mathbf{N}_{\text{irrat\_R}}(x) = \mathbf{N}_{\geq 2} \setminus \mathbf{N}_{\text{rat\_R}}(x)$ .

Here, we make use of these elementary insights, to construct simple tests which are easy to apply and which will allow us to recognize immediately – for an adequately chosen subset  $S \subset \mathbf{N}_{\geq 2}$  – whether  $\sqrt[n]{x} \in \mathbf{N}$  or  $\sqrt[n]{x} \notin \mathbf{Q}$  for  $x \in S$  and a natural number  $n \in \mathbf{N}_{\geq 2}$ , being arbitrarily chosen.

#### ➤ Test T1:

Let us start by considering the set

$S_1 = \{ x \in \mathbf{N}_{\geq 2} \mid x = p_1 \cdot p_2^{k_2} \cdot \dots \cdot p_i^{k_i} \cdot \dots \cdot p_m^{k_m}; \text{ where } p_i \text{ are prime numbers } \forall i \in \{1, 2, \dots, m\} \text{ and } p_i \neq p_j \forall i \neq j, m \geq 1, k_j \in \mathbf{N} \forall j \in \{2, 3, \dots, m\} \}$ .

It should be noted that, without loss of generality, it is sufficient that for at least one of the  $k_i$  in eq. (1) the condition  $k_i=1$  to hold. Then, the form required by  $S_1$  can be achieved by simply renumbering the  $k_i$ .

**TEST T1:** If  $x \in S_1$  then  $CD(\{k_1, k_2, \dots, k_m\}) = \emptyset$ .

Therefore,  $\mathbf{N}_{\text{rat\_R}}(S_1) = \emptyset$  and  $\mathbf{N}_{\text{irrat\_R}}(S_1) = \mathbf{N}_{\geq 2}$ , i.e.  $\sqrt[n]{x} \notin \mathbf{Q} \forall x \in S_1, \forall n \in \mathbf{N}_{\geq 2}$ .

*Remark:* A special case for  $S_1$  is  $S_1^* = \{ p \in \mathbf{N}_{\geq 2} \mid p \text{ is a prime number} \}$  and, therefore, it is proven that  $\sqrt[n]{p} \notin \mathbf{Q} \forall n \in \mathbf{N}_{\geq 2}$  and  $\forall p, p$  being a prime number.

#### ➤ Test T2:

Let us now consider the set

$S_2 = \{ x \in \mathbf{N}_{\geq 2} \mid x = p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m}; \text{ where } p_i \text{ are prime numbers } \forall i \in \{1, 2, \dots, m\} \text{ and } p_i \neq p_j \forall i \neq j, m \geq 2, k_i \in \mathbf{N} \forall i, \text{ and } \exists \alpha, \beta \in \{1, 2, \dots, m\}: k_\alpha \text{ and } k_\beta \text{ having no common divisors } d > 1 \}$ .

**TEST T2:** If  $x \in S_2$  then  $CD(\{k_1, k_2, \dots, k_m\}) = \emptyset$ .

Therefore,  $\mathbf{N}_{\text{rat\_R}}(S_2) = \emptyset$  and  $\mathbf{N}_{\text{irrat\_R}}(S_2) = \mathbf{N}_{\geq 2}$ , i.e.  $\sqrt[n]{x} \notin \mathbf{Q} \forall x \in S_2, \forall n \in \mathbf{N}_{\geq 2}$ .

Remark: A special case for  $S_2$  is  $S_2^* = \{ p_1^{k_1} \cdot p_2^{k_2} \mid p_1, p_2, k_1 \text{ (resp. } k_{_1}) \text{ and } k_2 \text{ (resp. } k_{_2}) \text{ are all prime numbers and } p_1 \neq p_2 \text{ and } k_1 \neq k_2 \}$ . Consequently, it is proven – for this special case of  $S_2$  – that  $\sqrt[n]{x} \notin \mathbf{Q} \forall n \in \mathbf{N}_{\geq 2}$  and  $\forall x \in S_2^*$ .

➤ Test T3:

We want to close this section with a test leading to the result that, for  $x \in S$ ,  $\mathbf{N}_{\text{rat}_R}(x)$  is not an empty set as in tests T1 and T2 but in this case it contains just one element.

Thus, let us consider the set

$S_3 = \{ x \in \mathbf{N}_{\geq 2} \mid x = p_1^{k_1} \cdot y, y=1 \text{ or } y = p_2^{k_2} \cdot p_3^{k_3} \cdot \dots \cdot p_m^{k_m}, m \geq 2 ; \text{ where } p_i \text{ and } k_i \text{ (resp. } k_{_i}) \text{ are prime numbers } \forall i \in \{1, 2, \dots, m\} \text{ and } p_i \neq p_j \forall i \neq j, \text{ and, in the prime factorization of } x, \text{ if } m \geq 2, \text{ all exponents } k_i \text{ (resp. } k_{_i}), i \geq 1, \text{ share the common divisor } k_1, \text{ but no other} \}$ .

**TEST T3:** If  $x \in S_3$  then  $\mathbf{N}_{\text{rat}_R}(x) = \{k_1\}$  and  $\mathbf{N}_{\text{irrat}_R}(x) = \mathbf{N}_{\geq 2} \setminus \{k_1\}$

EXAMPLES:

- **Ex2.1:**  $x = p_1^{101}$ , e.g.  $p_1=211$  (being a prime number), i.e.  $x$  represents a number possessing more than 200 digits. Applying test T3 (where we set  $y=1$ ) we rapidly find out that  $\mathbf{N}_{\text{rat}_R}(x) = \{101\}$ , i.e.  $\sqrt[n]{x} \notin \mathbf{Q} \forall n \in \mathbf{N}_{\geq 2}, n \neq 101$  as well as that  $\sqrt[101]{x} = p_1 \in \mathbf{N}$ .
- **Ex2.2:** Let  $S_{12} = \{ p^{12} \mid p \text{ representing an arbitrary prime number} \} \Rightarrow \mathbf{N}_{\text{rat}_R}(S_{12}) = \text{CD}(\{12\}) = \{2,3,4,6,12\}$ , i.e.  $\sqrt[n]{x} \in \mathbf{N} \forall x \in S_{12}, \forall n \in \mathbf{N}_{\text{rat}_R}(S_{12})$ . And, therefore,  $\mathbf{N}_{\text{irrat}_R}(S_{12}) = \mathbf{N}_{\geq 2} \setminus \{2,3,4,6,12\}$ .

Last but not least, let us shortly discuss the interesting question of how to determine the value of  $\sqrt[n]{x}$  if we know that  $\sqrt[n]{x} \in \mathbf{N}$ , i.e.  $n \in \mathbf{N}_{\text{rat}_R}(x)$ . Assuming again that the prime factorization of  $x$  is given and is such as specified by eq. (1) then, of course, we immediately recognize that

$$\sqrt[n]{x} = p_1^{k_1/n} \cdot p_2^{k_2/n} \cdot \dots \cdot p_m^{k_m/n} .$$

Actually, this is an astonishingly simple way to determine the value of any  $n$ -th root of any natural number  $x \in \mathbf{N}_{\geq 2}$ , if  $\sqrt[n]{x}$  results in a rational (and in this case, in particular, even in a natural) number.

## 4 Investigating the Set of Natural Numbers Possessing Irrational $n$ -th Roots

Let us now introduce an additional notation to name the total set of natural numbers, which possess irrational  $n$ -th roots only:

$$\mathbf{SN}_{\text{irrat}_R} := \{ x \in \mathbf{N}_{\geq 2} \mid \sqrt[n]{x} \notin \mathbf{Q} \forall n \in \mathbf{N}_{\geq 2} \} .$$

Considering test T1 (cf. Section 3) it is evident, that the set  $S_4 = \{ x \in \mathbf{N}_{\geq 2} \mid x=4y-2, y \in \mathbf{N} \}$  characterizing the even numbers not being an integer multiple of 4 represent a strict subset of  $\mathbf{SN}_{\text{irrat}_R}$ . In particular, the elements  $x$  of  $S_4$  possess a prime factorization such that

$$x = 2 \cdot p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m}, p_i \text{ denoting an odd prime number } \forall i \in \{1, 2, \dots, m\} .$$

Therefore  $S_4 \subset S_1$ .

Testing whether a given (even) number is an integer multiple of 4 is certainly very simple. We just have to test whether the last two digits are a multiple of 4. Consequently, we realize immediately that all natural numbers ending with 02, 06, 10, 14, ..., 98 are part of set  $S_4$  and for all the other even natural numbers  $n : n \notin S_4$ .

By finding the set  $S_4$  we have proven the result that for  $\forall x \in \mathbf{N}$  a number  $z \in S_4$  can be found in the immediate neighborhood of  $x$  such that  $|x-z| \leq 2$ . Also, the fact  $\mathbf{N}_{\text{irrat}_R}(S_4) = \mathbf{N}_{\geq 2}$  directly implies that  $z \in \mathbf{SN\_irrat\_roots}$ . Thus, we can say that the set  $\mathbf{SN\_irrat\_roots}$  is a “dense coverage” of the set  $\mathbf{N}$ .

Considering now, e.g., the set  $\mathbf{N}_{\leq 1000} := \{n \in \mathbf{N} \mid n \leq 1000\}$  already the properties of set  $S_4$  imply that at least 25% of all numbers of  $\mathbf{N}_{\leq 1000}$  are part of  $\mathbf{SN\_irrat\_roots}$ . A more detailed view shows that – defining regions of size 100 to partition the set  $\mathbf{N}_{\leq 1000}$  – we find only rather few natural numbers  $x$ , being part of each partition, for which  $x \notin \mathbf{SN\_irrat\_roots}$ .

In detail, we observe that for

- Partition P1 = {2, 3, ..., 99}:

$$x \notin \mathbf{SN\_irrat\_roots} \Leftrightarrow x \in M_1 = \{4, 8, 9, 16, 25, 27, 32, 36, 49, 64, 81\}.$$

Thus, only for 11 numbers in P1:  $\exists n \in \mathbf{N}_{\geq 2}$  with  $\sqrt[n]{x} \in \mathbf{N}, x \in M_1$ .

- Partition P2 = {100, 101, ..., 199}:

$$x \notin \mathbf{SN\_irrat\_roots} \Leftrightarrow x \in M_2 = \{100, 121, 125, 128, 144, 169, 196\}.$$

Thus, only for 7 numbers in P2:  $\exists n \in \mathbf{N}_{\geq 2}$  with  $\sqrt[n]{x} \in \mathbf{N}, x \in M_2$ .

- Partition P3 = {200, 201, ..., 299}:

$$x \notin \mathbf{SN\_irrat\_roots} \Leftrightarrow x \in M_3 = \{216, 225, 243, 256, 289\}.$$

Thus, only for 5 numbers in P3:  $\exists n \in \mathbf{N}_{\geq 2}$  with  $\sqrt[n]{x} \in \mathbf{N}, x \in M_3$ .

And in all the other partitions covering neighboring numbers (with a size of 100) between 300 and 1000 at least 97% of those 100 natural numbers are part of  $\mathbf{SN\_irrat\_roots}$ . This implies that, considering an arbitrary natural number  $x, 300 \leq x \leq 1000$ , in nearly all cases we will find a number  $x^*$  directly neighboring to  $x$  (distance=1) such that  $x^* \in \mathbf{SN\_irrat\_roots}$ . In almost all cases, however, even “ $x \in \mathbf{SN\_irrat\_roots}$ “ would already be satisfied for  $x$  itself.

## 5 Conclusions and Outlook

This contribution showed that understanding the properties of  $n$ -th roots  $\sqrt[n]{x}$  of natural and of positive rational numbers can be improved significantly if the argumentation directly relies on the prime factorization, resulting for  $x$ , as opposed to the usual argumentation based on polynomials. We demonstrated that, using the prime factorization of  $x$ , it is possible to design numerous tests which allow one to recognize in an astonishingly simple and efficient manner whether  $\sqrt[n]{x}$ , for arbitrary values  $x \in \mathbf{N}_{\geq 2}$  and  $n \in \mathbf{N}_{\geq 2}$ , result in a rational or an irrational value. In case  $\sqrt[n]{x} \in \mathbf{Q}$  (with the consequence  $\sqrt[n]{x} \in \mathbf{N}$ ), by means of an innovative calculation method, it is usually feasible to determine the exact result of  $\sqrt[n]{x}$  in a matter of seconds. Calculation methods to determine the result of  $\sqrt[n]{x}$  which also allow one to approximate the value of  $\sqrt[n]{x}$  (such as, e.g., Newton’s method [7 or 8,9,10]) can thereby be limited to the approximation of  $\sqrt[n]{x}$ , if  $\sqrt[n]{x} \notin \mathbf{Q}$ . Similarly, usage of the Lemma of Gauß [11] can be restricted to some cases in which we want to prove that  $\sqrt[n]{x} \notin \mathbf{Q}$  and, e.g., prime factorization of  $x$  could be practically infeasible.

The tests presented by us to cover the cases  $\sqrt[n]{x}$ , for  $x \in \mathbf{N}_{\geq 2}$ , can be extended in a straight-forward manner to cases  $\sqrt[n]{y}$ , in which  $y \in \mathbf{Q}^+ \setminus \mathbf{N}$ , for given  $n \in \mathbf{N}_{\geq 2}$  (where  $\mathbf{Q}^+$  denotes the set of positive rational numbers). Those modified tests could be used to prove that  $\sqrt[n]{y} \in \mathbf{Q}$  (still for  $y \in \mathbf{Q}^+ \setminus \mathbf{N}$ ) and, if so, to determine the value of  $\sqrt[n]{y}$  (which, typically, will no longer be a natural number).

As it has been proven by the author earlier [1]:  $\sqrt[n]{x} \notin \mathbf{Q} \quad \forall x \in \mathbf{R}^+ \setminus \mathbf{Q}, \forall n \in \mathbf{N}_{\geq 2}$ .

Therefore, evidently, for any given  $n \in \mathbf{N}_{\geq 2}$  it is not (!) possible that:  $\sqrt[n]{x} \in \mathbf{Q}$  for any  $x \in \mathbf{R}^+ \setminus \mathbf{Q}$ .

This contribution clearly shows that the decision whether the  $n$ -th root  $\sqrt[n]{x}$  of a natural number  $x$  possesses a rational value becomes rather trivial, if the prime factorization of  $x$  is available. In the particular case that a given natural number  $x$  is relatively small, e.g.  $x \leq 1000$ , then obtaining the prime factorization of  $x$  is actually a very simple task.

## Competing Interests

Author has declared that no competing interests exist.

## References

- [1] Wolfinger BE. Simple Criteria for  $\sqrt[n]{x}$  ( $n \in \mathbf{N}$ ,  $n \geq 2$ ,  $x \in \mathbf{R}$ ) Being a Rational or an Irrational Number. J. Advances in Mathematics and Computer Science. 2023;38(9):23-30.
- [2] Alvaro H. Salas S. Proving the existence of the  $n$ th root by induction. arXiv: General Mathematics; May 21, 2008.
- [3] Bressoud DM. Factorization and Primality Testing. Springer; 1989.
- [4] Odlyzko AM. The complexity of computing discrete logarithms and factoring integers. "§4.5 in: Open Problems in Communication and Computation (eds. Cover TM, Gopinath B.) Springer. 1987;113-116.
- [5] Pomerance C. Analysis and comparison of some integer factorization algorithms. in: Discrete Algorithms and Complexity (eds. Johnson DS, Nishizeki T, Nozaki A, Wilf HS.) Academic Press. 1987;119-143.
- [6] Rivest RL, Shamir A, Adleman L. A Method for Obtaining Digital Signatures and Public-key Cryptosystems. C.ACM. 1978;21(2):120-126.
- [7] Hubbard JH, Schleicher D, Sutherland S. How to Find All Roots of Complex Polynomials by Newtons Method. Inventiones Mathematicae. 2021;146:1-33.
- [8] Kelley CT. Solving Nonlinear Equations with Newton's Method, no1 in Fundamentals of Algorithms, SIAM; 2003.
- [9] Murugesan N, Ramasamy AMs. A numerical algorithm for nth root. Malaysian Journal of Fundamental and Applied Sciences, Penerbit UTM Press. 2014;7(1).
- [10] Özban AY. New methods for approximating square roots. Applied Mathematics and Computation. 2006; 175: 532-540.
- [11] Weisslein EW. Rational Zero Theorem. From MathWorld – A Wolfram Web Resource, <https://mathworld.wolfram.com/RationalZeroTheorem.html>

---

© 2024 Wolfinger; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/111735>