



# Characterization of Diverse Carrot (*Daucus carota* L.) Genotypes Using Long Noncoding RNA Based Microsatellite Markers

Aishwarya Hundekar <sup>a++</sup>, Sarvamangala S C <sup>b#\*</sup>,  
Manjunath H <sup>c#</sup>, Mahantesha B. N. Naika <sup>d#</sup>  
and Bapurayagouda P <sup>e#</sup>

<sup>a</sup> Department of Genetics and Plant Breeding, College of Horticulture, Bagalkot, Karnataka -587104, India.

<sup>b</sup> Genetics and Plant Breeding, Department of Biotechnology and Crop Improvement, College of Horticulture, Bagalkot, Karnataka -587104, India.

<sup>c</sup> Department of Plant Pathology, College of Horticulture, Bagalkot, Karnataka -587104, India.

<sup>d</sup> Department of Biotechnology and Crop Improvement, Kittur Rani Channamma College of Horticulture, Arabhavi, Karnataka -591306, India.

<sup>e</sup> Department of Seed Science and Technology, College of Horticulture, Bagalkot, Karnataka -587104, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i81255>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121173>

Original Research Article

Received: 04/06/2024  
Accepted: 06/08/2024  
Published: 10/08/2024

<sup>++</sup> PG Research Scholar;

<sup>#</sup> Assistant Professor;

<sup>\*</sup>Corresponding author: E-mail: sarucholin@gmail.com;

**Cite as:** Hundekar, Aishwarya, Sarvamangala S C, Manjunath H, Mahantesha B. N. Naika, and Bapurayagouda P. 2024. "Characterization of Diverse Carrot (*Daucus Carota* L.) Genotypes Using Long Noncoding RNA Based Microsatellite Markers". *Journal of Advances in Biology & Biotechnology* 27 (8):1316-24. <https://doi.org/10.9734/jabb/2024/v27i81255>.

## ABSTRACT

In the pursuit of enhancing carrot breeding programs, our study focused on the experimental validation of selected long non-coding RNAs (lncRNAs) using end-point polymerase chain reaction (PCR). Twenty-seven carrot (*Daucus carota* L.) genotypes, including Asiatic and Western varieties, were grown in RCBD design and evaluated for root traits at UHS Bagalkot, Karnataka, India, during 2022-2023. These genotypes were used to validate five SSR markers developed for lncRNAs in carrots. From an initial set, five lncRNAs were chosen for further experimental validation based on their bit score and the presence of simple sequence repeat (SSR) motifs. Our novel approach involved the development of 27 imperfect SSR markers and five perfect SSR markers specifically targeting these lncRNAs. The integration of SSR markers with lncRNA studies is unprecedented in carrot breeding, opening new avenues for genetic improvement. Among the five lncRNA-derived SSR (lncSSR) markers developed, *DcLNC55* and *DcLNC62* exhibited polymorphisms across 27 diverse carrot genotypes. In conclusion, our study highlights the innovative use of lnc-SSR markers in carrot breeding, demonstrating their utility in genetic diversity studies and their potential role in uncovering the functional importance of lncRNAs in crop phenotypes. The findings present significant implications for future carrot breeding strategies, providing a foundation for integrating molecular markers and lncRNA research in carrot and other related horticultural crop improvement programs.

**Keywords:** Carrot; long non-coding RNA; polymerase chain reaction; simple sequence repeat; polymorphism.

## 1. INTRODUCTION

Carrot (*Daucus carota* L.) is a biennial herb with a chromosome number of  $2n = 2x = 18$  with a genome size of 473Mb and belongs to the Apiaceae family [1]. Fewer studies are available on lncRNAs in carrots compared to model organisms like *Arabidopsis* despite the availability of high-quality genome sequences in carrots as large numbers of lncRNAs have been found in nature [2].

Long non-coding RNAs (lncRNAs), which exceed 200 nucleotides in length and do not code for proteins, play a crucial role in the regulation of gene expression at multiple levels [3]. To classify a transcript as "noncoding," it is assessed based on the absence of an extensive open reading frame (typically over 100 codons) and the lack of codon conservation [4]. In the nucleus, lncRNAs regulate gene expression through transcriptional and epigenetic mechanisms, including the modulation of chromatin compaction and the establishment of chromosomal connections [5]. They also influence gene expression by regulating the loading of histone or chromatin modifiers to chromatin [6]. Additionally, lncRNAs can directly affect transcription through the formation of R-loops, interference with RNA polymerase machinery, and transcription of the lncRNA locus. In the cytoplasm, a subset of lncRNAs is involved in controlling mRNA turnover and translation, highlighting their

multifaceted roles in gene expression regulation [7].

Plant lncRNAs, transcribed by RNA polymerases PolI, PolII, and PolIV, originate from various genomic regions, including enhancers, promoters, gene introns, pseudogenes, and as antisense transcripts to other genes [8,9]. These transcripts can be classified into different types based on their genomic position relative to protein-coding genes, such as natural antisense transcripts (lncNATs), intronic lncRNAs, intergenic lncRNAs (lincRNAs), and sense lncRNAs. lncNATs align with or diverge from sense strand transcripts, potentially modulating gene regulation, while intronic lncRNAs are transcribed from within introns of protein-coding genes without exon overlap. Sense lncRNAs share promoters with protein-coding genes and transcribe from regions overlapping exons [5,10]. Recent studies have revealed the involvement of lncRNAs in various biological processes, including flowering, reproduction, photomorphogenesis, vernalization, organ development, cell cycle control, and responses to biotic and abiotic stresses. The experimental validation of lncRNAs is essential to elucidate their interactions with transcriptional and translational components, providing a strong rationale for exploring their regulatory roles in different metabolic pathways in plants [11].

Simple Sequence Repeat (SSR) markers, also known as microsatellites, are highly polymorphic

and co-dominant markers widely used in genetic research. They consist of short, repetitive DNA sequences that are scattered throughout the genome, making them ideal for assessing genetic diversity and mapping genetic traits. Due to their high reproducibility and ease of analysis, SSR markers are invaluable tools in plant breeding and molecular biology [12]. In reality, there are few SSR markers unique to the carrot genome, and the earliest accounts of the developing SSRs date back only a few years [13,14]. Cavagnaro et al., 2011 used BACend sequences and an SSR-enriched genomic collection to create 300 SSR markers in carrots. SSRs were discovered by Iorizzo et al. in an assembly of the carrot transcriptome from four genotypes and provided a collection of 114 markers they identified by *in silico* investigations as polymorphic. Considering since then, most research on the genomics of carrots have employed these SSR markers [15].

## 2. MATERIALS AND METHODS

### 2.1 Plant Material

The present study was conducted at the Department of Biotechnology and Crop Improvement, University of Horticultural

Sciences, Bagalkot. Total of 27 genotypes carrot (*Daucus carota* L.) germplasm lines were grown, inclusive of Asiatic/Eastern and European/Western were used. The detailed names and colour description of the 27 carrot genotypes are listed in Table 1. The vegetative phase carrot root evaluation experiment was conducted at sector 1 farm (16°12'N, 75°12'45 E), University of Horticultural Sciences (UHS), Bagalkot, Karnataka, India during 2022-2023. It is part of a semi-arid tropical region with a mild climate, with typical temperatures between 23 °C and 35 °C and receives an average rainfall of 318 mm.

### 2.2 Primer Designing for lncRNAs

The lncRNAs were identified in the Alnc database and Primers were designed by searching microsatellite regions in the lncRNA transcripts. Sequences flanking the microsatellite regions in lncRNA were targeted for primer design using the Krait tool [16] by targeting the microsatellite flanking regions. The primer pairs were designed based on standard criteria, such as predicted melting temperature of 50-60°C and Guanine-Cytosine (GC) content of 45-55 percent for designing the primer pairs to obtain PCR amplicon length of 100-200 bp (Table 2).

**Table 1. List of carrot (*Daucus carota* L.) genotypes used for evaluation of genetic variability**

Sl. No.	Genotypes	Description
1.	UHSBC-34-2	Orange
2.	UHSBC-17	Orange
3.	UHSBC-31-2	Orange
4.	UHSBC-31-1	Orange
5.	UHSBC-32-2	Orange
6.	UHSBC-65	Released variety, Red colour (Pusa Rudhira)
7.	UHSBC146	Orange (Improved-population-2)
8.	UHSBC-23-1-1	Orange-red
9.	UHSBC-66	Black, released (Pusa Asita)
10.	UHSBC-31-4	Orange
11.	UHSBC-34-2-1	Orange
12.	UHSBC-100	Dark Orange (Western)
13.	UHSBC-46-1-1	Orange
14.	UHSBC-151	Reddish orange
15.	UHSBC-23-1	Orange
16.	UHSBC-150	Reddish orange
17.	UHSBC-1	Pale orange
18.	UHSBC-152	Orange
19.	UHSBC-160	Orange
20.	UHSBC-17-2	Orange
21.	UHSBC-59-66	White
22.	UHSBC-59	White
23.	UHSBC-17-1	Orange
24.	UHSBC-117	Orange
25.	UHSBC-150-1	Orange

### 2.3 Genomic DNA Extraction and Amplification of IncSSR Markers

The genomic DNA was extracted from carrot leaves using CTAB extraction method [17]. DNA amplification was carried out using IncRNA-SSR primers for 27 genotypes of carrot to check their utility and functional validation. Amplification of DNA for the respective target IncRNA was performed using polymerase chain reaction (PCR) in a 10 µl reaction mixture for IncRNA validation each containing 2 µl template DNA (50 ng/µl), 0.5 µl forward primers (5 µmol/l), 0.5 µl reverse primers (5 µmol/l), 2 µl ddH<sub>2</sub>O and 5 µl 2 × Taq PCR Master mix. The PCR program was as follows: 1 min at 95 °C, 20 s denaturing at 95 °C, 45 s annealing at 47-53 °C (Annealing temperature was optimised for each primer set using gradient PCR before actual amplification) and 30 s elongation at 72 °C followed by a final step at 72 °C for 2 min. The details of primer sequences used for amplification of IncRNA is presented in the Table 3. After the PCR, fractionation was done using agarose gel electrophoresis to assess the amplification of target genes followed by EtBr staining. 3.5 % agarose gel was used for separation of Inc-SSR markers in 1 × TBE buffer at initial 85V for 15 minutes followed by 120 V for 1 h. Subsequently, the banding pattern was visualized using a gel documentation system and clear images of the bands were captured and scoring was performed. The clear and unambiguous bands of all the polymorphic IncSSR markers were scored for 27 carrot genotypes and calculated manually.

## 3. RESULTS AND DISCUSSION

### 3.1 Identification of SSR Regions in IncRNA and Marker Development

The identified sets of transcripts from the Alnc database were used to search for simple sequence repeats (SSR) in the Krait tool. 32 sequences were considered to contain microsatellites or SSR motifs. The identified SSRs have mono, di, tri, tetra, and hexa nucleotide repeat motifs. Among 32, perfect SSRs were five (Table 2a) and the remaining 27 were imperfect SSRs (Table 2b). From the perfect SSRs- two SSRs contain mono- a nucleotide repeat motif, two SSRs contain di- a nucleotide repeats motif and the remaining one has hexa- nucleotide repeat motif. Among the imperfect SSRs- five SSRs contain mono-

nucleotide repeat motif, eleven SSRs contain di- a nucleotide repeat motif, six SSRs contain tri- nucleotide repeat motif, four SSRs contain quad- nucleotide repeat motif and the remaining one has hexa- nucleotide repeat motif. One perfect SSR (STRG.55.1) was chosen for validation as it consists of a hexa-nucleotide repeat motif. Four among the imperfect SSRs, STRG.25.1 (di- nucleotide repeat motifs), STRG.62.1 and STRG.70.1 (tri-nucleotide repeat motifs), and STRG.91.1 (tetra nucleotide repeat motifs) were chosen for validation because of the presence of a greater number of repeats and higher chances of polymorphism.

### 3.2 Validation of IncRNA by PCR

A total of five SSR markers were used for the validation of 27 genotypes which yielded clear and scoreable bands. Among the five SSR markers considered for validation *DcLnc91*, *DcLnc25* and *DcLnc70* showed monomorphic bands of 140, 100, and 140bp respectively. While markers *DcLNC55* and *DcLNC62* showed polymorphism with band sizes ranging from 100-125 bp and 200-480 bp respectively (Table 4) and gel profile (Fig. 1) indicating their suitability in the carrot genetic diversity assessment. For validation of IncRNA, 5 Inc-SSR markers were used. Among them, 3 Inc-SSR markers showed monomorphic bands, while 2 markers showed polymorphism among 27 diverse carrot genotypes. Polymorphic SSR markers can reveal genetic diversity or variations in IncRNA sequences among different genotypes and would be further useful for mapping and understanding its role in phenotypic expression of economic traits in carrots.

### 3.3 Genotypic Scores for Inc-SSR Markers

For Inc-SSR allele sizing, the genotypes were analysed by comparing the sizes of the amplified bands against a 100bp reference ladder. Each genotype's band size was measured and recorded manually by comparing the DNA fragment lengths to a molecular weight marker on an agarose gel ensuring accurate identification of allele variations. The use of the 100bp ladder provides a reliable standard for consistent and reproducible results across different samples. The observation of monomorphic bands in 3 and-SSR markers and polymorphic bands in two markers suggests potential genetic variation.

**Table 2a. List of lncRNA perfect SSRs designed**

Sl. No.	Sequence ID	Entry	Product	Forward	Tm1	GC1	Stability1	Reverse	Tm2	GC2	Stability2
1	STRG.36.1	1	122	ACAAAGCTCCCCTCCC	54.77	62.5	4.3	CGGTGGATAACAGTGTGC	55.8	55.56	4.57
2	STRG.45.1	1	114	CCTTAATTCTATGATAAAGGCTGG	55.12	37.5	4.85	ACATGCAACCCCTAAAAGc	56.07	47.37	3.51
3	STRG.48.1	1	134	CCGGCCATGACCAAGTCC	60.44	66.67	3.85	GAGGGGGAGAGAGAGCGG	60.52	72.22	5.52
4	STRG.55.1	1	122	TTTCTTGGGTAAAGCAAGG	53.01	42.11	3.61	GACTTCTTAAAGGCGATACC	54.41	42.86	2.73
5	STRG.70.1	1	145	AAAAGGGGAAAAGTGCGGC	58.54	55.56	6.53	CAATGACCATTTTTATCAAACCC	57.02	37.5	4.95

**Table 2b. List of lncRNA imperfect SSRs**

Sl. No.	Sequence ID	Forward	Tm1	GC1	Stability1	Reverse	Tm2	GC2	Stability2
1	STRG.8.1	CCTCTGCTCTGCTAAGCTCC	59.9	60	4.7	ATCGGAATCTGAGCGGCG	59.97	61.11	6.46
2	STRG.8.1	CCTCTGCTCTGCTAAGCTCC	59.9	60	4.7	ATCGGAATCTGAGCGGCG	59.97	61.11	6.46
3	STRG.24.1	AAACAACAATGCCTGGGC	56.48	50	5.36	CTCCAGGAGAGTTTTAGTAGG	54.49	47.62	3.18
4	STRG.25.1	TGCTGTTTGCTTTCCCGG	58.56	55.56	5.73	GCGTGTATTCCAAAATGGCC	59.33	45.45	5.36
5	STRG.27.1	TCCAACCCAGCGATTCCGG	59.73	61.11	4.3	TCTCCGATGAACACCCGCG	59.82	61.11	6.46
6	STRG.34.1	CACGGGCGTTTGAATGGC	60.13	61.11	4.4	TCACACGTTCCGAAGAGCC	60.01	57.89	4.7
7	STRG.36.1	ACAAAGCTCCCCTCCC	54.77	62.5	4.3	CGGTGGATAACAGTGTGC	55.8	55.56	4.57
8	STRG.40.1	CAATCCTATTATTCAACCCCC	53.68	42.86	5.4	TTCTCGGTGTTGTGTGC	54.94	52.94	4.57
9	STRG.45.1	CCTTAATTCTATGATAAAGGCTGG	55.12	37.5	4.85	ACATGCAACCCCTAAAAGC	56.07	47.37	3.51
10	STRG.48.1	CCGGCCATGACCAAGTCC	60.44	66.67	3.85	GAGGGGGAGAGAGAGCGG	60.52	72.22	5.52
11	STRG.48.1	CCGGCCATGACCAAGTCC	60.44	66.67	3.85	TCAAGAGTGGGAGAGAGAGGG	59.99	57.14	4.3
12	STRG.48.1	TTATCTCTCTCCCTCTCC	50.64	50	3.71	AGAGAGAGAGAGTGGG	49.42	56.25	4.61
13	STRG.49.1	GCACCAAGCAAATTTTCGG	56.29	47.37	4.3	GGCGGATCTAGGAAAAGGC	57.99	57.89	4.35
14	STRG.50.1	GTGAGGTCTGCGGCGG	60.15	75	6.13	ACCCTCGACAAATTCAAATAGGC	59.31	43.48	3.93
15	STRG.51.1	TGTCAAGGCAGGACAGAAGC	60.25	55	3.86	GTAAAGGGCAAGGCAGGC	58.71	61.11	4.85
16	STRG.53.1	TGAATAGGACTGCGAGAGAAAGG	59.87	47.83	3.11	AGTAAGGAGGGGCGTACG	58.07	61.11	3.67
17	STRG.55.1	TGATTTTCTTGGGTAAAGCAAGG	57.53	39.13	3.61	AGCTTAAGACACATCCAATCCC	58.11	45.45	3.85
18	STRG.62.1	CTCCTTTATAATTTAACAGGTGGG	55.23	37.5	4.61	CCTCAAAGTCCAAAGTGC	56.42	52.63	4.4

Sl. No.	Sequence ID	Forward	Tm1	GC1	Stability1	Reverse	Tm2	GC2	Stability2
19	STRG.62.1	TCTTCCTTTGATTATCCCACAGG	57.88	43.48	4	TGTGCAAACCTCCAAAGTCC	58.6	50	3.85
20	STRG.66.1	CCCTCTTTCTCAGCGCC	57.3	64.71	6.53	CGACGATCTCGAAATCATGG	56.53	50	3.66
21	STRG.70.1	AAAAGGGGAAAGTGCGGC	58.54	55.56	6.53	CAATGACCATTTTTATCAAACCCC	57.02	37.5	4.95
22	STRG.70.1	AATTGTGGGGTTTGATAAAAATGG	56.61	33.33	3.16	AATTTTGCCCCCACCACC	58.1	55.56	4.61
23	STRG.75.1	TCACTACACTAGTCTCTCTCTCC	57.65	47.83	3.71	AGGTGTATGCAGAGAGGCC	58.78	57.89	5.19
24	STRG.78.1	CCAACAACCCAGCAGCCC	61.31	66.67	5.19	GGTGTGTCTCCCCAGTAATGG	60.07	57.14	3.16
25	STRG.84.1	AGCTCGCTCACTTCTGGC	59.74	61.11	4.85	GCGACAATGATTTCTCCGGC	59.97	55	6.13
26	STRG.91.1	GGGAGGGAGGGAAGATCC	57.68	66.67	3.36	ATTATCACAATGCTTTTTCTTCC	55.74	33.33	3.46
27	STRG.91.1	GTGATAATGTAAGTAAGTAGCTGC	55.46	37.5	5.25	TTTGTTCCTCCCTCCTTGC	54.99	50	4.01




Fig. 1. Validation of IncSSR markers

**Table 3. The details of Inc-SSR markers used for validation in endpoint PCR**

Sl. No.	Primer ID	Forward primer sequence (5'-3')	Reverse primer sequence (5'-3')
1.	DcLNC55	TTTCTTGGGTAAAGCAAGG	GACTTTCTTAAAGGCGATACC
2.	DcLNC62	CTCCTTTATAATTTAACAGGTGGG	CCTCCAAAGTCCAAAGTGC
3.	DcLNC91	GTGATAATGTAAGTAAGTAGCTGC	TTTGTTCCTCCTTGC
4.	DcLNC25	TGCTGTTTGCTTTCCCGG	GCGTGTTTATTCCAAAATGGCC
5.	DcLNC70	AAAAGGGGAAAGTGCGGC	CAATGACCATTTTTATCAAACCCC

**Table 4. List of the Inc-SSRs amplification data in carrot (*Daucus carota* L.) genotypes**

Sl. No.	Primer ID	Annealing temperature (°C)	Observed amplicons size (bp)
1.	DcLNC55	47	100-125
2.	DcLNC62	48	200-480
3.	DcLNC91	48	140
4.	DcLNC25	53	100
5.	DcLNC70	50	140

#### 4. CONCLUSION

The identification of polymorphic bands in two out of five lncRNA SSR markers used for validation indicates genetic variation in the genotypes. This study underscores the pivotal role of long non-coding RNAs (lncRNAs) in the genetic and phenotypic diversity of carrot genotypes. By developing novel lncRNA-derived SSR markers, we have provided useful tools for assessing genetic variation in carrots. The polymorphism observed in these markers among diverse carrot genotypes highlights their potential application in carrot breeding programs. Our findings open new avenues for integrating molecular markers and lncRNA research in crop improvement strategies.

#### FUTURE SCOPE

Generating genome-wide lncRNAs and utilizing the polymorphic markers in future analyses will enhance insights into the genetic diversity of regulatory roles of these non-coding sequences. They help breeders to understand the functional roles and regulation of important genes influencing carrot morphology and other phenotypic characteristics. This knowledge can be leveraged to develop new carrot varieties with desirable traits, enhancing breeding efficiency and outcomes.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Iorizzo M, Ellison S, Senalik D, Zeng P, Satapoomin P, Huang J, Bowman M, Iovene M, Sanseverino W, Cavagnaro P, Yildiz M. A high-quality carrot genome assembly provides new insights into carotenoid accumulation and asterid genome evolution. *Nat. Genet.* 2016;48(6): 657-666.
- Rai MI, Alam M, Lightfoot DA, Gurha P, Afzal AJ. Classification and experimental identification of plant long non-coding RNAs. *Genomics.* 2019;111(5):997-1005.
- Budak H, Kaya SB, Cagirici HB. Long non-coding RNA in plants in the era of reference sequences. *Frontiers in Plant Science.* 2020;11:276.
- Morris KV, Mattick JS. The rise of regulatory RNA. *Nat. Rev. Genet.* 2014; 15(6):423-437.
- Ma X, Zhao F, Zhou B. The characters of non-coding RNAs and their biological roles in plant development and abiotic stress response. *Int. J. Mol. Sci.* 2022;23(8): 4124.
- Cui J. lncRNA in plants: Function, mechanisms and applications. *Frontiers in Plant Science.* 2023;14:1238185.



7. Waititu JK, Zhang C, Liu J, Wang H. Plant non-coding RNAs: Origin, biogenesis, mode of action and their roles in abiotic stress. *Int. J. Mol. Sci.* 2020;21(21):8401.
8. Axtell MJ. Classification and comparison of small RNAs from plants. *Annu. Rev. Plant Biol.* 2013;64:137-159.
9. Dekeba DG. Advances on the application of non-coding RNA in crop improvement. *Afr. J. Biotechnol.* 2021;20(11):440-450.
10. Meng X, Li A, Yu B, Li S. Interplay between miRNAs and lncRNAs: Mode of action and biological roles in plant development and stress adaptation. *Comput. Struct. Biotechnol. J.* 2021;19:2567-2574.
11. Chekanova JA. Long non-coding RNAs and their functions in plants. *Current opinion in plant biology.* 2015;27:207-216.
12. Taheri S, Lee Abdullah T, Yusop MR, Hanafi MM, Sahebi M, Azizi P, Shamshiri RR. Mining and development of novel SSR markers using next generation sequencing (NGS) data in plants. *Molecules.* 2018;23(2):399.
13. Iorizzo M, Senalik DA, Grzebelus D, Bowman M, Cavagnaro PF, Matvienko M, Ashrafi H, Van Deynze A, Simon PW. De novo assembly and characterization of the carrot transcriptome reveals novel genes, new markers, and genetic diversity. *BMC genomics.* 2011;12:1-14.
14. Cavagnaro PF, Chung SM, Manin S, Yildiz M, Ali A, Alessandro MS, Iorizzo M, Senalik DA, Simon PW. Microsatellite isolation and marker development in carrot-genomic distribution, linkage mapping, genetic diversity analysis and marker transferability across Apiaceae. *BMC genomics.* 2011;12:1-20.
15. Uncu AO, Uncu AT. High-throughput simple sequence repeat (SSR) mining saturates the carrot (*Daucus carota* L.) genome with chromosome-anchored markers. *Biotechnology & Biotechnological Equipment.* 2020;34(1):1-9.
16. Du L, Zhang C, Liu Q, Zhang X, Yue B. Krait: an ultrafast tool for genome-wide survey of microsatellites and primer design. *Bioinformatics.* 2018;34(4):681-683.
17. Doyle J. DNA protocols for plants. In *Molecular techniques in taxonomy* Berlin, Heidelberg: Springer Berlin Heidelberg. 1991;283-293.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/121173>