



The Use of Bio-control Agents in Mulberry Pest Management: Successful Techniques and Important Issues

Naveen Chandra Reddy ^{a++*}, Thrilekha D ^{a++},
Dukare Pradip Gulabrao ^{a++}, P. Harshita Mala ^{a#},
Ashish S Karur ^{a++}, Bharath Kumar B. M ^{a++},
Pavithra M R ^{b++} and Ashrith S ^{b++}

^a Department of Sericulture, University of Agricultural Sciences, College of Agriculture, GKVK, Bengaluru -560065, India.

^b Post Graduate Studies, University of Agricultural Sciences Bangalore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i94416>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122416>

Review Article

Received: 27/06/2024

Accepted: 29/08/2024

Published: 01/09/2024

ABSTRACT

Since bio-control agents target specific pests without affecting the usefulness insects or/and damaging the soil and water resources, it plays a role in conservation of ecological balance in the environment. A variety of insect and non-insect pests, as well as hosts of parasites (fungi, insect,

⁺⁺ M.Sc. (Agri.) in Sericulture;

[#] Ph.D in Sericulture;

^{*}Corresponding author: E-mail: ncreddy006@gmail.com

Cite as: Reddy, Naveen Chandra, Thrilekha D, Dukare Pradip Gulabrao, P. Harshita Mala, Ashish S Karur, Bharath Kumar B. M, Pavithra M R, and Ashrith S. 2024. "The Use of Bio-Control Agents in Mulberry Pest Management: Successful Techniques and Important Issues". *International Journal of Environment and Climate Change* 14 (9):330-37. <https://doi.org/10.9734/ijecc/2024/v14i94416>.

and mite parasites), occur in silkworm sericulture. This method does not in any way influence the sustainability of environment through pest control as required by sustainable agriculture. Since bio-control agents are known to be harmless to human beings and other organisms that may be of equal importance in the ecosystem such as pollinators, wildlife etc., they also have safety advantages. This is a big step forward to eradicating pests as compared to the chemical pesticides which have been known to affect the human lives and other wildlife. In the meantime, the bio-control agents may be cost effective in the long run. Because bio-control agents could offer continuous control of pests with minimal extra cost, it is possible, even when the outlay of obtaining and developing such organisms is more expensive than straightforward chemical applications. Further, the use of bio-control agents may reduce the reliance on traditional chemical pesticides and hence cost outlay and pest control costs and also reduce the probabilities of pesticide resistance. In the farming of mulberry, bio control agents are one the most important components of integrated pest management and have proved be an effective substitute to chemical control. Bio-control agents offer long-term pest suppression in comparison to chemical control methods, whose impacts tend to weaken after several generations of treatment. The continuous use of chemicals may result in pest resistance to the pesticides used. Spraying synthetic pesticides leads to the formation of pesticide residues on mulberry foliage, which affects food products and environmental health.

Keywords: Bio-control; bio-control agents; challenges; mulberry; pest management; strategies.

1. INTRODUCTION

Bio-control of pests and diseases is quite successful in modern agriculture. Recent developments in the fields of scientific pest management technologies have enabled researchers to find alternative solutions to the synthetic limitations in sericulture. Thus, the time is ripe to discuss the innovative growth-promoting and biological control agents, including experiences with commercial products employed for the efficient management of harmful insects in mulberry cultivation [1]. A variety of insect and non-insect pests, as well as hosts of parasites (fungi, insect, and mite parasites), occur in silkworm sericulture. Mulberry is the sole host plant for silkworms, and it is grown in over 70 countries across the world. Over 96% of the world's silk is produced from mulberry grown in India. The management of pests certainly requires artificial measures, such as the application of pesticides. Mulberry is subjected to numerous pests, including mosquitoes, dipterous larvae, lepidoptera, mealybugs, and silkworms. These insects can have an impact on the quality and quantity of leaf yield crop. The deciduous spray of broad-spectrum pesticides may greatly promote crop production, but pests can become immune after repeated treatments. The sting of sucking pests and broad-spectrum pesticides has many harmful effects on unusual fauna and animals [2]. The alteration and fragmentation of non-target organisms, including pests of crops and soil microorganisms, in response to pesticides.

Aviation could also harm humans, as well as undesirable chemicals induced by broad-spectrum pesticides [3].

Human can decrease the adverse effects of long-term use of broad-spectrum pesticides by developing appropriate non-chemical procedures, mechanical methods, and biological control. A number of researchers have also worked to control pests by utilizing non-chemical techniques around the globe. Biopesticides include neem products (*Azadirachta indica* oil, *Azadirachta indica* kernel liquid, etc.), tobacco products (*Nicotiana tabacum* extracts, etc.), Mexican marigold (*Tagetes patula* extract), gliricidia (*Giricidia* extract), contact poison (Magnesium fluoride, K_2SiF_6 , etc.) [4,5], and biological control agents. Bio, botanical, and contact actives are used in the control of pink mealybug. Bio-control agents or biocontrol are mechanisms to avoid large increases in pest populations due to competition or predation [6,7].

The mulberry or *Morus* sp. is an important agricultural product used in the production of sericulture and is renowned for being used in the making of silk [8]. Leaves are the sole feed of silkworms (*Bombyx mori* L.), which spins a cocoon out of fibers and produces silk, an article of immense cultural and market value on the global level [9]. Apart from their use in silkworm farming, the plant is used in traditional medicine, fruit production and as ornamental trees in landscaping. The demand for mulberries and

their products worldwide indicates how important it is to sustain a healthy, high-yielding type [10].

Nevertheless, like all similar crops, mulberry has many threats in the form of pests and diseases that pose danger to the trees' health and productivity. Some of the common pests include Aphids, Scales, Borers while the diseases include fungal, bacterial, viral, etc. These threats can cause significant economic losses and impact the quality of the mulberry leaves used in the sericulture [11]. In response to these challenges, traditional pest management methods have mainly focused on the use of chemical pesticides. These chemicals although efficient in pest control have the following impacts; pollution of the environment, toxicity to organisms that are not pests and potential harm to man. Additionally, excessive reliance on chemical pesticides can result in pest resistance thus their effectiveness declines over time and require more of it or a stronger chemical [12].

These problems have been met by a shift towards more sustainable organic farming methods that include reduced use of chemicals and enhanced resource conservation. Biological control agents have been found promising when applied as a replacement to synthetic pesticides [13]. These agents, which play a part in managing the biological balance of pests, are either living organisms or substances that are extracted from live organisms. Using bio-control agents in pest management is a significant difference in pest management aiming at reducing the excessive reliance on artificial substances that are not friendly to the ecosystem [14].

Bio-control agents are typically categorized into three main groups: These include predators, parasitoids, and pathogens. In this case, predators are living things that have a direct control on other pests through feeding on them. Parasitoids deposit their eggs usually on or inside the pest insects and the larvae consume and destroy the host [15]. These are organisms that invade and destroy pests, and normally target specific pests or stages of pest's life cycle. The various forms of bio-control agents have different modes of action for an integrated approach to pest control [16].

However, the use of bio-control agents in mulberry has its difficulties involved in the process. Since most bio-control agents are species specific, one cannot completely eliminate

pest problems by using these agents alone; a precise choice and application of the bio-control agents are required for optimum pest control [17]. Temperature and humidity also pose a challenge influencing the efficacy of bio-control agents and thus the need to make adjustments. Furthermore, the integration of biocontrol agents to other pest management practices such as cultural and physical control entails proper planning and management [18].

The use of bio-control agents and products in mulberry farming is still growing due to evolving research and technology. The following is likely to improve the effectiveness and flexibility of bio-control agents through genetic improvements and different formulations: More extensive ecological study of bio-control agents, pests and their environment is therefore required to enhance bio-control efficacy in various conditions.

2. UNDERSTANDING BIO-CONTROL AGENTS

Bio-control agents refer to those living organisms that are used in the control of pests and diseases through efficient natural processes. They can be broadly categorized into three types: predators, parasitoids and pathogens and include organisms such as insects, spiders, nematodes and fungi [19].

Predators: These are organisms that feed on other pests as opposed to plants, animals and other food materials that are obtainable in the environment. The parasitic insects including lady beetles, the lacewings and spiders serve as biological control agents to suppress pests in mulberry production. Predatory agents are animals that can capture and feed on prey, limiting pest populations.

Parasitoids: Parasites are usually oviposit on or in another insect (host). Its larvae which are in the process of developing feed on the host and finally kills it. Some of the commonly employed parasitoids include the wasps that are selective in the pests they attack in mulberry plant [20].

Pathogens: These are independent causal organisms and include bacteria, fungi, viruses and nematodes that infect and kill pests. Some bioagents in the form of pathogens have been found to provide significant level of suppression of certain pests and diseases in mulberry [21]. Therefore, pathogens are utilized to reduce

hemolymph cellularity and in some cases to induce apoptosis in insect hemocytes. Pathogens are also developed in the fat bodies. The released spores hence multiply in numbers to suppress hemocytes, facilitating various beneficial reactions. However, ingestion of toxins (spores) is essential [22,23]. Once the enterotoxic spores exert their action, infection in the hemocytes takes over to complete the season of suppression. Pathogens due to these versatile activities are considered as successful bio-control agents possessing the potential to manage crop pests sustainably. Moreover, pathogens are not specific in infecting only a few CNA, unlike parasites that have to be hosted by one or two CNA and, therefore, have suggesting better applicability in pest managing tropical crop ecosystem. For addressing the issue of critical concern, we have to look at more information including the exclusion of other infection routes and immune protective strategies if developed [24].

3. KEY BIO-CONTROL AGENTS USED IN MULBERRY PLANTATION

3.1 Predators

Lady Beetles (Coccinellidae): They are most suitable for agrophotous arthropods such as aphids, scale insects and other pests of soft-bodied nature. This is a plus since they feed aggressively and ensure that pest infestations are minimal at all times.

Lacewings (Chrysopidae): Lacewing larvae that are often referred to as 'aphid lions', because they feed on aphids, whiteflies and other similar insects. They are very beneficial in that they can greatly minimize pest populations in mulberry orchards.

Spiders: Orb-web and jumping-spiders are only some of the favoured spider species that feed on different pests. These are also useful in the general pest management involved in mulberry farming [25].

3.2 Parasitoids

Trichogramma Wasps: These are small parasitic wasps that lay their eggs into those of other moths and butterflies possibly including the mulberry. *Trichogramma* sp. is first of all parasitic at the egg stage of pests. it plays a role of pest control in early stages to avoid occurrence of a big crisis.

Braconid Wasps: These parasitoids are known to parasite the caterpillars and other larval stages of the pests. They are useful at controlling pests including the mulberry borers [26].

3.3 Pathogens

Bacillus thuringiensis (Bt): These toxins are aimed at caterpillars and other insect larvae that are invulnerable to other kinds of bacteria. Formulations of Bt are typically employed in their organic mulberry plantation for the management of pests such as borers [27].

Beauveria bassiana: This fungus parasitize and causes death to a number of insect pests. At present it is applied to control pests including whitefly and spider mites in mulberry [25].

Metarhizium anisopliae: The other useful fungus is *M. anisopliae* that can be useful in controlling pests that dwell on the soil and feed on mulberry plants; this control is particularly helpful in controlling beetles and other ground feeding insects [28].

4. EFFECTIVE STRATEGIES FOR IMPLEMENTING BIO-CONTROL AGENTS

The application of the bio-control agents applies an overall method of pest control where bio-control agents are used as one of the segments. Several key strategies are essential for effective use: Several key strategies are essential for effective use:

Choice of Effective Bio-Control Agents: The effectiveness of a number of these bio-control agents depends on the choice of correct agents for specific pests and conditions of the environment. It is made depending on the type of the target pest, effectiveness of the agent used and the other conditions prevailing from time to time. Such parasitoids as *Trichogramma* sp. To control the moth and butterfly, the wasps preferably seek the egg stage of the insects; lady beetles (*Coccinellidae*), predators work effectively on aphids and scale insects. Entomopathogenic fungi such as *Beauveria bassiana* intends to kill a broad spectrum of insect pest whereas Bacterial pathogens as formerly mentioned, for example *Bacillus thuringiensis* (Bt) is applied against caterpillar pests. To carry out this selection it is crucial to note the correct identification of pests and also to understand the life cycle and behaviour of the bio-control agent [29].

Timing and Application: More generally, the action time and the way of application of the bio-control agents must be considered. The use of bio-control agents in a way that optimize the timing of a pest population, is scientifically known as sensitive timing [30]. Suppose, to guarantee that the parasitoids are able to parasitize the eggs before they hatch, for example, introduce *Trichogramma* sp. wasps at the time of the moths when the female is laying eggs. The movement of the bio-control agents throughout the crop should have been ensured by the application techniques employed [31]. To completely cover contaminated areas this may need to involve using agents with specific methods of coverage, or to let the agents seep out bit by bit in specific time frames.

Monitoring and Evaluation: Evaluations are needed at a more frequent basis to identify the extent of the effectiveness of bio-control agents. The bio-control agents inhabiting the area and the sporadic surveys of pest populations help in ascertaining whether the bio control agents are having an effectiveness. Monitoring also shows that it is possible to adjust the extent of the effort tactically whether by increasing the frequency of the releases or changing the application techniques due to the initial efforts being insufficient. The decisions are based on information drawn from monitoring endeavours; these also help in the progressive enhancement of bio-control strategies [32].

4.1 Benefits of Using Bio-control Agents

Use of successful bio-control technological intervention in mulberry agro-ecosystem offers distinct advantages compared to the conventional chemical interventions. Biopesticides containing living microorganisms such as bacteria, fungi, viruses, protozoa, and beneficial nematodes are potential biological control agents to suppress a wide variety of pests and plant pathogens using multiple modes of action. The use of these kinds of bio-control offers great potential to control the various pests at mulberry and have several advantages. The major advantage is that they leave very few or no residues in both the host plants and environment. The microorganisms are destroyed rapidly by environmental factors. The requirement of beneficial microorganisms for bio-control agents for such requirements is lower. These agents offer attractive environment restoration. As the incidence of pests and pathogens is considerably reduced, there is no longer a need to use large quantities of chemicals. Consequently, toxic

chemicals do not spread throughout the environment. Bio-control agents offer long-term pest suppression in comparison to chemical control methods, whose impacts tend to weaken after several generations of treatment. The continuous use of chemicals may result in pest resistance to the pesticides used. Spraying synthetic pesticides leads to the formation of pesticide residues on mulberry foliage, which affects food products and environmental health. In the medium and long-term, the use of conventional pesticides results in inflation and reduced efficiency [33,34]. Bio-control agents or bio-pesticides offer a cheaper alternative to pesticides. Using natural predators to minimize pests and harmful insects avoids the direct costs associated with pesticides and their application. In fact, the reduction in the use of chemicals helps to minimize the risk of pest resurgence.

The following are benefits of using bio-control agents:

Environmental Benefits: Bio-control agents are better than chemical pesticides from the point of view of environmental conservation. They decline on the use of synthetic chemicals, the possibility of pesticide residue in food, water pollution and soil pollution is also eliminated [35].

Sustainability: Bio-control agencies have been found to help in the sustainable production of mulberry since they foster the natural pest control. They have minimal effects on the environment retaining bio-diversity and are far less probable to cause pest resistance as is the case with chemical pesticides [36].

Safety: Compared to chemical pest killers, this is one way of pest control that has little impacts on health hazards. They are normally biodegradable, as well as having low toxicity to humans and other species of beneficial insects and wildlife [37].

Cost-Effectiveness: When it has been set up, bio-control agents for pest management are long-term. Because they may have setup costs in their introduction and the costs involved in pest management if reduced through the use of these organisms may prove cheaper in the long run mainly because of little or no use of chemicals [38].

5. CONCLUSION

In conclusion, bio-control agents are an important significant step in the sustainable

control of diseases and pests in mulberry production system. Their respective roles in the current agriculture are well demonstrated by the level of efficiency they have in providing pest control but at the same time minimizing on the rate of harm that they cause on the environment, the level of economic benefits that they bring in addition to the health and safety of the people who are involved in the farming process. The use of bio-control agents in the integrated pest management system will indeed become paramount with improvements in science and technology so as to retain the health of the mulberry crops as well as the yield. As the present problems addressed in the production of mulberry and with the advancement of the upcoming technologies attach the solution to it, the agricultural method assuring more stability and sustainability to this important crop will not only build the future but retains the prosperity and viability it needs. In the farming of mulberry, bio control agents are one the most important components of integrated pest management and have proved be an effective substitute to chemical control. As the global demand for mulberry products especially in sericulture production improves it is strategic that there be strong and improved yielding crops. Introducing parasitoids, diseases and predators is consistent with methods of modern agriculture that first of all preserves crop health, has a positive balance of cost and income, and does not have a detrimental effect on the environment.

6. FUTURE PROSPECTS AND RESEARCH DIRECTIONS

In the future, we believe that molecular and genetic advancements and new technologies with user-friendly approaches should be developed for isolating, producing, and enhancing the efficacy of these bio-priming agents and biopesticides. In addition, the use of other sources for obtaining or isolating novel microbial isolates or their metabolites for the control of multiple pests attacking the mulberry fields has potential and is an area that needs to be explored. Considering the advantages and future growth, integrated use of bio-priming agents or bio-inputs could be a trend that would be widely accepted in the management of insect pests attacking crops and would be a promising innovation in total. More biochemical and molecular research is needed to be conducted to reveal important genetic or transcription regulation, payment pathways stimulated by these BCA compounds that are related to

defense systems. Research needs to be initiated on various aspects, including: (1) BCA sprayed mulberry leaves expecting that applied at one concentration (100 mg/mL⁻¹) to determine dose minimal important application (2-10 mg/mL⁻¹), even from Bt formulations derived from different *Aspergillus* isolates their treatments against different *A. flavus* strains. Therefore, a kind of performance evaluation, or additional comparison research, or an examination of the potential BCA versus others, is not restricted to *A. flavus* or mulberry fruits.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Baruah JP, Kalita C. Integrated pest management of uzi fly (*Exorista sorbillans*) in Muga silkworm *Antheraea assamensis* Helfer (Lepidoptera: saturniidae): A review. *Journal of Entomology and Zoology Studies*. 2020;8(4):341-3.
2. Bawaskar DM, Chowdary NB, Kedar SC, Reddy BT, Selvaraj C, Rathore MS, Srinivas C, Navik O. Traditional and innovative technologies for pest management of tropical tasar silkworm, *Antheraea mylitta* (Drury) by the tribes of Eastern-Central India. *International Journal of Tropical Insect Science*. 2022 Apr;42(2): 1737-48.
3. Saini P, Rohela GK, Kumar JS, Shabnam AA, Kumar A. Cultivation, utilization, and economic benefits of Mulberry. In *The Mulberry Genome*. Cham: Springer International Publishing. 2023; 13-56.
4. Dandin SB, Kumari V. Mulberry (*Morus* sp.) cultivation for sustainable sericulture. CRC Press. 2021 Jul 25;188-207.
5. Gupta SK, Mukhopadhyay SK, Bhattacharyya H, Modak BK. Integrated management of diseases and pests of silkworm. *Journal of Environment and Sociobiology*. 2016;13(2):147-55.

6. Elango K, Sobhana E, Sujithra P, Bharath D, Ahuja A. Traditional agricultural practices as a tool for management of insects and nematode pests of crops: An overview. *Journal of Entomology and Zoology Studies*. 2020; 8(3):237-245.
7. Koshariya AK, Jadhav SM. Chapter-4 organic farming and integrated nutrient management (INM) in horticulture. *Microbial Flocculants as an Alternate to Chemical Flocculants for Wastewater Treatment*. 2021;49.
8. Karthick Mani Bharathi, B., S. Susikaran, K.T. Parthiban, K.A. Muruges and K. Chozhan: The economics of commercial mulberry saplings production using mini clonal technology over conventional method. *The Pharma Innovation Journal*. 2022;11:1236-1241.
9. Bindroo BB. Breakthroughs and betterments in silkworm improvement. *Beneficial Insect Farming-Benefits and Livelihood Generation*. 2016:163.
10. Karthick Mani Bharathi B, Susikaran S, Parthiban KT. A comparative biochemical study of mulberry (*Morus* spp.) mini clones over conventional stem cuttings. *International Journal of Plant & Soil Science*. 2024;36:975-983.
11. Karthick Mani Bharathi B, Susikaran S, Parthiban KT, Vasanth V, Vijay S. Influence of different transplanting days on yield attributes of mini clones under field conditions for *Morus indica* (V1). *Madras Agricultural Journal*. 2023;111:1-3.
12. Madyarov Shukhrat R. Biotechnological approaches in sericultural science and technology of Uzbekistan. *International Journal of Industrial Entomology*. 2005; 11(1):13-9.
13. Singh RN, Maheshwari M, Saratchandra B. Sampling, surveillance and forecasting of insect population for integrated pest management in sericulture. *International Journal of Industrial Entomology*. 2004; 8(1):17-26.
14. Poveda J. *Trichoderma* as bio-control agent against pests: New uses for a mycoparasite. *Biological Control*. 2021 Aug 1;159:104634.
15. Singh RN, Saratchandra B. Biological control strategy of uzi fly in sericulture. *International Journal of Industrial Entomology*. 2003;6(2):125-32.
16. Sakthivel N, Kumaresan P, Qadri SM, Ravikumar J, Balakrishna R. Adoption of integrated pest management practices in sericulture-A case study in Tamil Nadu. *Journal of Biopesticides*. 2012;5:212.
17. Unni BG, Devi B, Kakoty Y, Wann SB, Borah A, Dowarah P. Role of Plant: Microbe interactions in the sustainable development of Muga sericulture. *Bacteria in Agrobiolgy: Plant Probiotics*. 2012:213-25.
18. Singh A, Bhardwaj R, Singh IK. Bio-control agents: Potential of biopesticides for integrated pest management. *Biofertilizers for Sustainable Agriculture and Environment*. 2019:413-33.
19. Rahman S, Biswas SK, Barman NC, Ferdous T. Plant extract as selective pesticide for integrated pest management. *Biotechnological Research*. 2016 Jan 31;2(1):6-10.
20. Singh A, Kumar V, Majumdar M, Guha L, Neog K. A comprehensive review of insect pest management in Muga silkworm (*Antheraea assamensis* Helfer): Current scenario and future prospects. *Journal of Experimental Agriculture International*. 2024 Mar 7;46(5):47-55.
21. Van Zyl C, Malan AP. The role of entomopathogenic nematodes as biological control agents of insect pests, with emphasis on the history of their mass culturing and in vivo production. *African Entomology*. 2014 Jul 1;22(2):235-49.
22. Singh RN, Maheshwari M, Saratchandra B. Biocoenology and control of whiteflies in sericulture. *Insect Science*. 2005 Dec; 12(6):401-12.
23. Singh S, Singh A, Kumar S, Mittal P, Singh IK. Protease inhibitors: Recent advancement in its usage as a potential bio-control agent for insect pest management. *Insect science*. 2020 Apr; 27(2):186-201.
24. El-Wakeil N, Saleh M, Abu-hashim M. Cottage industry of biocontrol agents and their applications. *Cottage Industry of Biocontrol Agents and Their Applications*; 2020.
25. Singh D, Raina TK, Singh J. Entomopathogenic fungi: An effective bio-control agent for management of insect populations naturally. *Journal of Pharmaceutical Sciences and Research*. 2017 Jun 1;9(6):833.
26. Singh RN, Rao JK, Sampson MV. Role of parasitoids in pest management in Tasar culture. In *biocontrol potential and its exploitation in sustainable agriculture*.

- Boston, MA: Springer US. Insect Pests. 2001 Nov 30;2:379-387.
27. Liu SS, Rao A, Vinson SB. Biological control in China: Past, present and future -An introduction to this special issue. Biol. Control. 2014 Jan 1;68(1):5.
28. Ballal CR. Biological control for sustainable plant protection. Ensuring Food Safety, Security and Sustainability through Crop Protection. 2021;5:20-26.
29. Nadaf HA, Vishaka GV, Sathyanarayana K, Chandrashekharaiah M, Rathore MS, Balaji Chowdary N, Reddy BT, Selvaraj C. Integrated farming system—a key to sustainable livelihood in tasar sericulture. Journal of Experimental Zoology India. 2022 Jul 1;25(2).
30. Bhat A, Fatima A and Dar KA. Biocontrol and its role in sericulture: A review paper. International Journal of Advance Research in Science and Engineering. 2018;7: 1881-5.
31. Singh RN, Saratchandra B. An integrated approach in the pest management in sericulture. International Journal of Industrial Entomology. 2002;5(2):141-51.
32. Cappellozza S, Casartelli M, Sandrelli F, Saviane A, Tettamanti G. Silkworm and Silk: Traditional and Innovative Applications. Insects. 2022 Nov 3;13(11): 1016.
33. St. Leger RJ, Wang C. Genetic engineering of fungal bio-control agents to achieve greater efficacy against insect pests. Applied Microbiology and Biotechnology. 2010 Jan;85:901-907.
34. Sujatha GS, Kumar GA, Teja KS, Devi DL, Panda A, Rupali JS, Gautam SK. A comprehensive review of the effect and mitigation of climate change on sericulture. International Journal of Environment and Climate Change. 2024 Jul 25;14(7):776-88.
35. Sabbahi R, Hock V, Azzaoui K, Saoiabi S, Hammouti B. A global perspective of entomopathogens as microbial bio-control agents of insect pests. Journal of Agriculture and Food Research. 2022 Dec 1;10:100376.
36. Singh RN, Maheshwari M. Biological control of pests of Non-mulberry silkworms and its host plants in India. International Journal of Industrial Entomology. 2002; 4(2):83-91.
37. Sharma K, Kapoor B. Sericulture as a profit-based industry—A review. Indian Journal of Pure and Applied Biosciences. 2020;8(4):550-62.
38. Wani MY, Mir MR, Baqual MF, Mehraj K, Bhat TA, Rani S. Role of foliar sprays in sericulture industry. Journal of Pharmacognosy and Phytochemistry. 2017;6(4):1803-6.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/122416>