

Effect of Different Seasons on Pregnancy Gender Outcome in Sample of Iraqi Women

Israa, F. Jaafar ^{a*≡}, Safa Muntadher Fawzi ^{b⊖} and Islam Muntadher Fawzi ^{b⊖}

^a Al-Esraa University College, Baghdad, Iraq.

^b Ministry of Health, Baghdad Al - Russafa Health, Republic of Iraq.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Asmaa Fathi Moustafa Hamouda, Jazan University, Saudi Arabia.

Reviewers:

(1) Ihor Atabiekov, Moscow Regional Oncological Dispensary, Russia.

(2) Miljana Z Jovandaric, University Clinical Centre of Serbia, Serbia.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/77198>

Original Research Article

Received 13 September 2021

Accepted 27 November 2021

Published 07 December 2021

ABSTRACT

Background: Sex ratio is important for human life since its creation, some families prefer male than females, also its related to some medical problems that affect certain sex than another. Studies showed that males are more fragile, and might die earlier than females. Climate has a certain effect on male to female ratio of the pregnancy outcome and certain seasons are in favor of male than female reproduction.

Objectives: To evaluate the effect of climate and seasonal variation on the sex of live births in a sample of pregnant Iraqi women.

Subjects and Methods: This study was carried out on 2424 live babies (apart from 225 miscarriages) of women attending Al-Numan teaching hospital and Al-Husaynia health care center, from January 2017 till January 2019. The time of last menstrual period recall was taken from each mother of this study. An early ultrasound test was done to confirm the accuracy of pregnancy duration. The gender of live babies, maternal age and parity were recorded.

Results: The highest birth rate were in Autumn (721) then Winter (662), Spring (577) and the least in summer (521).

The male to female ratio increased with maternal age. Mothers of 30 years and over have 1.09 male female ratio as compared with those under the age of 30 years (0.929).

[≡] MBChB, MSc, PhD;

[⊖] MBChB;

*Corresponding author: Email: pro.israa@esraa.edu.iq;

The male female ratio was highest in Autumn 1.06 compared with other seasons; 0.97, 0.9233, 0.887 Winter, Spring and Summer respectively. The abortion rate was only (0.09) from total births.

Conclusion: The best season for conception in Iraq was Autumn. The total male female ratio regardless the season was 0.9659. The male female ratio was best in Autumn then Winter, Spring, lastly Summer. Sex differences shows that; males are conceived in seasons of more favorable conditions for reproduction.

Keywords: Birth rate; climate; Iraqi women; gender; maternal age; pregnancy; seasons; sex ratio.

1. INTRODUCTION

Sex ratio is important for human life since his creation . the gender of a pregnancy outcome has many implications like socioeconomic, some families prefer male babies than females, some health problems affect certain sex more than the other etc. most studies had shown that males are more susceptible to be affected by harsh environmental conditions like heat i.e. they are more delicate and might die earlier than female babies [1]. Human studies suggested that climate has certain effect on male to female ratio of normal pregnancies and certain seasons are in favor of male than female reproduction. Mothers usually get pregnant with male babies at optimal conditions than females.

The male to female ratio (i.e. sex ratio) decreases, as a results of changes in conditions of environment like; contaminated environment [2,3], destructive earthquakes , smoking parents[4,5], and aged mothers or fathers [6].

Paternal age is of importance especially in mothers with multiple pregnancies, because the nutrients and energy available for their babies are less [5,6]. Hopefully the diurnal pattern and changes in temperature and environments in different seasons are important for conservation of species. The plan of people using certain seasons for breeding by designing the time of reproduction and hence delivery to seasons that are more favorable for the survival of newborn. Cagnacci and Volpe, claimed that human being are not seasonal breeders. but if human follows certain rules for conception as changes in climates according to seasonal variation [7], This would have good applicants for babies to be conceived and then delivered in better situations so give us a good environment for more male babies in more favorable conditions .Melnikov *et al.* reported an excellent percentage of mother conception with male babies in western parts of Siberia during summer time, but less male ratio in winter [8].

Since the length between conception and delivery i.e gestational period might vary, so the Sex ratio at birth may not be very accurate [9]. So in this study evaluation of the effect of variation between seasons and its effect on reproduction rates and the sex ratio was detected as accurate as possible

2. SUBJECTS AND METHODS

This study was carried out on 2481live birth (apart from 225 miscarriages) from women attending Al –Numan teaching hospital and Al - Husaynia health care center in Baghdad, from January 2017 till January 2019.The pool of mothers was from Baghdad center and rural areas around it.

Information on the time of conception was taken from mothers of the babies involved in the study. For every single pregnancy, the time of conception had been taken according to the date of the last menstrual period and was confirmed by gynecological examination.

To be accurate in calculating the duration of pregnancy , an early ultrasound test was done to confirm the accuracy of pregnancy. Women ageranges between (15–45 years), and only live healthy births involved in the season variation of male female ratio, while the miscarriages were calculated independently.

The ratio of the sex was assessed to evaluate both the month at which pregnancy occurred and the month at which they have their babies. The duration of pregnancy of every single delivery, and the sex of babies i.e male to female ratio of pregnancies that was ended at different weeks of gestation was assessed. The sex ratio was calculated according to the month and the season of delivery.

Observation of variation in sex ratio at different times of gestation were assessed, and at different months and so in different seasons of the year.

3. RESULTS

This study included 2736 pregnancies :2481 normal live birth from Iraqi women and 255 are aborted early in pregnancy. The study carried out from January 2017 till January 2019.

Table 1 shows the demographic data of the studied group; the age of mothers was (15-45 year) with a mean± SD of (30 ± 5.6 years).

The differences in number births between seasons are shown in Fig. 1.

This figure shows that the highest birth rate was in Autumn (29.1%) then Winter (26.7%), Spring (23.2%) and least in summer (21%).

The distribution of pregnancy outcomes as male and female are shown in Fig. 2. The highest number. of males were in autumn then winter spring and summer respectively.

Table 1. The demographic data of all women with their new born babies participating in this study

Total no.	2736
Mother age	15-45
Number of live births	2481
Number of male babies	1219
Number of female babies	1262
No of miscarriages	225




Fig. 1. seasonal variation of total birth rates




Fig. 2. The differences between male and female outcome of pregnancy according to season(left side male ;right side female)

Table 2. Characteristics of the birth-sex ratios according to age and season

Characteristic	Male	Female	Sex Ratio (M/F)	p-value
Mother Age				
≥30 total 600	313	287	1.090	0.05
<30 1881	906	975	0.929	0.05
Total male&female2481	1219	1262	0.9659	0.05
Pregnancy Season				
Summer	245	276	0.887	0.05
Spring	277	300	0.9233	0.05
Autumn	371	350	1.06	0.01
Winter	326	336	0.97	0.05

P ≤ 0.05 regarded as significant

Table 3. Miscarriages during seasons

Miscarriage history	Total of seasons	Winter	Spring	Summer	Autumn	P value
<30 year	124	41	4	35	44	0.01
≥30 year	101	36	3	18	44	0.01
total	225	77	7	53	88	0.01

P ≤ 0.05 regarded as significant

The total male female ratio in the studied group is shown in (Table 2) . The male to female ratios was 0.9659.

The male female ratio increased with mothers age. At the age of 30 years and over the male to female ratio was 1.09 as compared with those under the age of 30 years which was 0.929.

While the best season for males was Autumn as shown by male female ratio of 1.06. then Winter, Spring and Summer shows; 0.97,0.9233 ,0.887 respectively.

Table 3 shows miscarriages .The total abortion rate was only 0.09, which was more under the age of 30 ,and less over 30 years.

Highest number of miscarriages were in Autumn then in Winter, Summer and the least were in Spring.

4. DISCUSSION

The results of this study showed that pregnant females in summer or spring are more likely to deliver girls than boys, when they are compared with pregnant in autumn then winter . Our results were partly aligned with previous studies that stated there is a seasonality variation of birth-sex ratio they pointed to variation in sex with different seasons [10].

Another study conducted In Siberia showed that summer season in Siberia is the season of male babies [8].Male babies are more fragile than female so it requires cool weather ,in Iraq summer has long day 16 hours and very hot temperature ranges between 47- 52 ° C. that' s why summer is not the season for male in Iraq.

This is the 1st study done in Baghdad , the change in male ratio can be attributed to effect of melatonin since it increases ovulation and it's a survival factor for sperms as well [11], when the weather in Iraq transferred from summer to Autumn day night cycle changed: day hours12 hour especially in October and November. Melatonin release is more this means more antioxidant and good fertility environment hence good fertility rates [12,13,14] and so it's a good climate for male babies since melatonin level is higher [15].

The abortion rates were highest in autumn due to the large number of pregnancies so in comparison with another seasons it seems more. The male to female ratio increased with maternal age it was more at 30 years and over (1.09) as compared with those under the age of 30 years (0.929).This can be attributed to the increased age of marriage in Iraq and can be explained that the more mature the mother is it takes more care of herself, as good nutrition as well as good sleeping hours.

5. CONCLUSIONS

From this study we concluded that the best season for conception in Iraq was Autumn. The total male to female ratio regardless the season was 0.9659. The male to female ratio was best in Autumn then Winter , Spring and lastly Summer. This supports the idea that more males than females are conceived in seasons with more favorable reproductive conditions. The male to female ratio increased with maternal age it was more at 30 years and over (1.09) as compared with those under the age of 30 years (0.929).

6. RECOMMENDATIONS

1. We recommend to do year to year comparison since the whole environmental factors are changing yearly.
2. Add number of women participating in the study and in different areas of Iraq.
3. Study the effect of socioeconomic class on gender outcome.

CONSENT

It is not applicable.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected from Republic of Iraq, Ministry of health, Baghdad Al Russafa Health and preserved by the authors.

COMPETING INTEREST

Authors have declared that no competing interests exist.

REFERENCES

1. Cagnacci A, Renzi S, Arangino C, Alessandrini A, Volpe A .The male disadvantage and the seasonal rhythm of sex ratio at the time of conception Human Reproduction. 2003;18(4): 885–887.
2. Astolfi P, Zonta LA. Reduced male births in major Italian cities: Human Reproduction.1999;14:3116–3119.
3. Mocarelli P, Gerthoux PM, Ferrari E, Patterson DJ, Kieszak SM, Brambilla P, et al. Paternal concentration of dioxin and sex ratio of offspring: Lancet. 2000;355: 1858–1863.
4. Fukuda M, Fukuda K, Shimizu T, Andersen CY, Byskov AG. Parental periconceptional smoking and male: female ratio of newborn infants: Lancet. 2002;359:1407–1408.
5. Orvos H, Kozinszky Z, Bartfai G. Natural variation in the human sex ratio. Hum. Reprod. 2001;16:803.
6. Nicolich MJ, Huebner WW, Schnatter RA. Influence of parental and biological factors on the male birth fraction in the United states: an analysis of birth certificate data from 1964 through 1988: Fertil. Steril. 2000;73:487 –492.
7. Cagnacci A, Volpe A. Influence of melatonin and photoperiod on animal and human reproduction: J. Endocrinol. Invest. 1996;19 :382 –411.
8. Melnikov VN, Grech V. Seasonality of live birth sex ratio in south western Siberia, Russia, 1959–2001: J. Epidemiol. Community Health, 2003;57: 471–472.
9. James, W.H. Why are boys more likely to be preterm than girls? Plus other related conundrums in human reproduction: Hum. Reprod. 2000;15: 2108 –2111.
10. James W.H. Interpregnancy intervals, high maternal age and seasonal effects on the human sex ratio. Hum. Reprod. 1996;11:7–8.
11. Tong J, Sheng S, Sun Y, Li H, Li WP, Zhang C. et al. Melatonin levels in follicular fluid as markers for IVF outcomes and predicting ovarian reserve: Reproduction. 2017;153: 443–451.
12. Shavi Fernando, LukRombauts. Melatonin: shedding light on infertility? - a review of the recent literature: J .ovarian Res. 2014;7:98.
13. Javier Espino, MaríaMacedo, Graciela Lozano, Águeda Ortiz , Carmina Rodríguez , Ana B. et al. Impact of Melatonin Supplementation in Women with Unexplained Infertility Undergoing Fertility Treatment: Antioxidants. 2019; 8:338.
14. Zhang HM, Zhang Y. Melatonin: A well-documented antioxidant with conditional pro-oxidant actions: J. Pineal Res. 2014;57:131–146.
15. Chao Song, Wei Peng, Songna Yin, Jiamin Zhao, Beibei Fun , Jingcheng

Zhang, et al . Melatonin improves oxidative stress in mice =:
age-induced fertility decline and www.nature.com/ Scientific Reports. 2016;
attenuates ovarian mitochondrial 1- 15.

© 2021 Jaafar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77198>